



# wwPDB X-ray Structure Validation Summary Report

Oct 3, 2023 – 04:26 AM EDT

PDB ID : 6O41  
Title : Crystal structure of the unbound PGZL1 germline Fab fragment  
(PGZL1\_gVmDmJ)  
Authors : Irimia, A.; Wilson, I.A.  
Deposited on : 2019-02-27  
Resolution : 2.46 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : **FAILED**  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : **FAILED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1

## 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.46 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 11744 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called germline PGZL1\_gVmDmJ light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	213	Total 1668	C 1037	N 291	O 335	S 5	0	4	0
1	A	213	Total 1635	C 1019	N 279	O 332	S 5	0	1	0
1	C	213	Total 1635	C 1019	N 279	O 332	S 5	0	1	0

- Molecule 2 is a protein called germline PGZL1\_gVmDmJ heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	223	Total 1652	C 1044	N 275	O 324	S 9	0	4	0
2	B	222	Total 1643	C 1039	N 274	O 321	S 9	0	3	0
2	D	222	Total 1639	C 1036	N 273	O 321	S 9	0	3	0

- Molecule 3 is a protein called Immunoglobulin G-binding protein G (DOMAIN III).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	M	60	Total 465	C 292	N 76	O 97	0	0	0
3	N	60	Total 473	C 296	N 78	O 99	0	1	0
3	O	59	Total 460	C 289	N 75	O 96	0	0	0

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
M	1	MET	-	initiating methionine	UNP P19909
M	62	HIS	-	expression tag	UNP P19909

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
M	63	HIS	-	expression tag	UNP P19909
M	64	HIS	-	expression tag	UNP P19909
M	65	HIS	-	expression tag	UNP P19909
M	66	HIS	-	expression tag	UNP P19909
N	1	MET	-	initiating methionine	UNP P19909
N	62	HIS	-	expression tag	UNP P19909
N	63	HIS	-	expression tag	UNP P19909
N	64	HIS	-	expression tag	UNP P19909
N	65	HIS	-	expression tag	UNP P19909
N	66	HIS	-	expression tag	UNP P19909
O	1	MET	-	initiating methionine	UNP P19909
O	62	HIS	-	expression tag	UNP P19909
O	63	HIS	-	expression tag	UNP P19909
O	64	HIS	-	expression tag	UNP P19909
O	65	HIS	-	expression tag	UNP P19909
O	66	HIS	-	expression tag	UNP P19909

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	L	1	Total	C O	0	0
			6	3 3		
4	L	1	Total	C O	0	0
			6	3 3		
4	L	1	Total	C O	0	0
			6	3 3		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	L	1	Total 6	C 3	O 3	0	0
4	L	1	Total 6	C 3	O 3	0	0
4	H	1	Total 6	C 3	O 3	0	0
4	H	1	Total 6	C 3	O 3	0	0
4	A	1	Total 6	C 3	O 3	0	0
4	B	1	Total 6	C 3	O 3	0	0
4	C	1	Total 6	C 3	O 3	0	0
4	C	1	Total 6	C 3	O 3	0	0
4	D	1	Total 6	C 3	O 3	0	0
4	O	1	Total 6	C 3	O 3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	L	79	Total 79	O 79	0	0
5	H	58	Total 58	O 58	0	0
5	A	59	Total 59	O 59	0	0
5	B	47	Total 47	O 47	0	0
5	C	72	Total 72	O 72	0	0
5	D	42	Total 42	O 42	0	0
5	M	16	Total 16	O 16	0	0
5	N	13	Total 13	O 13	0	0
5	O	10	Total 10	O 10	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.

### 3 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.84Å 107.49Å 131.96Å 90.00° 94.61° 90.00°	Depositor
Resolution (Å)	39.67 – 2.46	Depositor
% Data completeness (in resolution range)	97.8 (39.67-2.46)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.68 (at 2.48Å)	Xtrriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, $R_{free}$	0.169 , 0.219	Depositor
Wilson B-factor (Å <sup>2</sup> )	50.0	Xtrriage
Anisotropy	0.211	Xtrriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	11744	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 4 Model quality [i](#)

### 4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.3 Torsion angles [i](#)

#### 4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 4.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GOL	L	304	-	5,5,5	0.90	0	5,5,5	1.02	0
4	GOL	L	303	-	5,5,5	0.92	0	5,5,5	0.95	0
4	GOL	C	302	-	5,5,5	0.97	0	5,5,5	1.01	0
4	GOL	D	301	-	5,5,5	0.82	0	5,5,5	1.20	1 (20%)
4	GOL	A	301	-	5,5,5	0.72	0	5,5,5	1.20	0
4	GOL	H	302	-	5,5,5	1.03	0	5,5,5	0.98	0
4	GOL	L	301	-	5,5,5	0.92	0	5,5,5	0.98	0
4	GOL	L	302	-	5,5,5	0.97	0	5,5,5	0.93	0
4	GOL	B	301	-	5,5,5	1.00	0	5,5,5	0.89	0
4	GOL	L	305	-	5,5,5	1.01	0	5,5,5	0.85	0
4	GOL	O	101	-	5,5,5	0.88	0	5,5,5	0.84	0
4	GOL	H	301	-	5,5,5	1.05	0	5,5,5	0.95	0
4	GOL	C	301	-	5,5,5	0.80	0	5,5,5	1.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	L	304	-	-	0/4/4/4	-
4	GOL	L	303	-	-	4/4/4/4	-
4	GOL	C	302	-	-	2/4/4/4	-
4	GOL	D	301	-	-	2/4/4/4	-
4	GOL	A	301	-	-	4/4/4/4	-
4	GOL	H	302	-	-	2/4/4/4	-
4	GOL	L	301	-	-	1/4/4/4	-
4	GOL	L	302	-	-	2/4/4/4	-
4	GOL	B	301	-	-	0/4/4/4	-
4	GOL	L	305	-	-	2/4/4/4	-
4	GOL	O	101	-	-	1/4/4/4	-
4	GOL	H	301	-	-	2/4/4/4	-
4	GOL	C	301	-	-	0/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	301	GOL	C3-C2-C1	-2.21	103.11	111.70

There are no chirality outliers.

5 of 22 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	L	303	GOL	O1-C1-C2-O2
4	L	303	GOL	O1-C1-C2-C3
4	H	301	GOL	C1-C2-C3-O3
4	H	302	GOL	O1-C1-C2-C3
4	A	301	GOL	O1-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

#### 4.7 Other polymers [i](#)

There are no such residues in this entry.

#### 4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 5 Fit of model and data

### 5.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

### 5.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

### 5.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

### 5.4 Ligands

EDS failed to run properly - this section is therefore empty.

### 5.5 Other polymers

EDS failed to run properly - this section is therefore empty.