



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 17, 2023 – 11:07 AM EDT

PDB ID : 1O5X
Title : Plasmodium falciparum TIM complexed to 2-phosphoglycerate
Authors : Parthasarathy, S.; Eaazhisai, K.; Balaram, H.; Balaram, P.; Murthy, M.R.
Deposited on : 2003-10-06
Resolution : 1.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1619 (1.14-1.06)
Clashscore	141614	1671 (1.14-1.06)
Ramachandran outliers	138981	1615 (1.14-1.06)
Sidechain outliers	138945	1613 (1.14-1.06)
RSRZ outliers	127900	1588 (1.14-1.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	248	 2% 90% 8% ..
1	B	248	 4% 78% 19% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	3PY	A	3300	-	X	-	-
4	2PG	B	4400[A]	-	-	X	-
4	2PG	B	4400[B]	X	X	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4784 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Triosephosphate isomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	246	1988	1259	336	388	5	0	11	0
1	B	246	2101	1329	353	414	5	0	34	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	163	VAL	ALA	engineered mutation	UNP Q07412
B	163	VAL	ALA	engineered mutation	UNP Q07412

- Molecule 2 is PHOSPHITE ION (three-letter code: PO3) (formula: O₃P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	4	3	1	0	0

- Molecule 3 is 3-HYDROXYPYRUVIC ACID (three-letter code: 3PY) (formula: $C_3H_4O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	3	4		

- Molecule 4 is 2-PHOSPHOGLYCERIC ACID (three-letter code: 2PG) (formula: $C_3H_7O_7P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	O	P	0	1
			22	6	14	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	337	Total 337	O 337	0	0
5	B	325	Total 325	O 325	0	2



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Triosephosphate isomerase



- Molecule 1: Triosephosphate isomerase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	53.49Å 50.90Å 89.01Å 90.00° 91.90° 90.00°	Depositor
Resolution (Å)	30.00 – 1.10 23.37 – 1.10	Depositor EDS
% Data completeness (in resolution range)	87.9 (30.00-1.10) 88.8 (23.37-1.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.65 (at 1.10Å)	Xtrriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.133 , 0.171 0.131 , 0.130	Depositor DCC
R_{free} test set	2107 reflections (1.22%)	wwPDB-VP
Wilson B-factor (Å ²)	9.2	Xtrriage
Anisotropy	0.356	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 64.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.022 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	4784	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: 2PG, PO3, 3PY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.74	0/2065	1.28	17/2784 (0.6%)
1	B	1.21	4/2193 (0.2%)	1.37	29/2961 (1.0%)
All	All	1.01	4/4258 (0.1%)	1.33	46/5745 (0.8%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	218[B]	SER	CA-CB	30.45	1.98	1.52
1	B	218[A]	SER	CA-CB	30.45	1.98	1.52
1	B	248	MET	C-OXT	7.03	1.36	1.23
1	B	70	GLY	N-CA	5.20	1.53	1.46

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	134	ARG	CD-NE-CZ	17.99	148.79	123.60
1	A	134	ARG	NE-CZ-NH1	-12.26	114.17	120.30
1	A	134	ARG	NE-CZ-NH2	9.95	125.28	120.30
1	B	218[B]	SER	N-CA-CB	-9.81	95.78	110.50
1	B	218[A]	SER	N-CA-CB	-9.81	95.78	110.50
1	A	3	ARG	NE-CZ-NH1	-9.05	115.78	120.30
1	B	229	PHE	CB-CG-CD2	-8.83	114.62	120.80
1	A	98	ARG	NE-CZ-NH2	-8.39	116.11	120.30
1	A	205	ARG	NE-CZ-NH1	8.21	124.41	120.30
1	B	98	ARG	NE-CZ-NH2	-7.97	116.31	120.30
1	B	218[B]	SER	CA-CB-OG	-7.87	89.96	111.20
1	B	218[A]	SER	CA-CB-OG	-7.87	89.96	111.20
1	A	98	ARG	NE-CZ-NH1	7.37	123.98	120.30
1	A	48	TYR	CB-CG-CD1	-7.26	116.64	121.00
1	A	205	ARG	NE-CZ-NH2	-7.14	116.73	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	134	ARG	NE-CZ-NH2	-7.00	116.80	120.30
1	B	165[A]	GLU	OE1-CD-OE2	-6.78	115.16	123.30
1	B	165[B]	GLU	OE1-CD-OE2	-6.78	115.16	123.30
1	B	49[A]	ASP	CB-CG-OD1	-6.71	112.26	118.30
1	B	49[B]	ASP	CB-CG-OD1	-6.71	112.26	118.30
1	B	98	ARG	NE-CZ-NH1	6.49	123.55	120.30
1	B	27	PHE	CB-CG-CD1	6.38	125.27	120.80
1	B	218[B]	SER	CB-CA-C	6.34	122.14	110.10
1	B	218[A]	SER	CB-CA-C	6.34	122.14	110.10
1	A	99	ARG	NE-CZ-NH2	-6.31	117.15	120.30
1	A	50[A]	HIS	CA-CB-CG	6.04	123.87	113.60
1	A	50[B]	HIS	CA-CB-CG	6.04	123.87	113.60
1	B	69	PHE	C-N-CA	-6.01	109.68	122.30
1	A	222	GLN	CA-CB-CG	5.92	126.42	113.40
1	B	110	ARG	NE-CZ-NH2	-5.91	117.34	120.30
1	B	48	TYR	CB-CG-CD2	5.89	124.53	121.00
1	A	158	ASP	CB-CG-OD1	-5.83	113.06	118.30
1	B	101	TYR	CG-CD1-CE1	5.65	125.82	121.30
1	B	27	PHE	CB-CG-CD2	-5.58	116.89	120.80
1	B	31	ASP	CB-CG-OD2	-5.57	113.29	118.30
1	B	229	PHE	CB-CG-CD1	5.55	124.69	120.80
1	B	134	ARG	NH1-CZ-NH2	5.51	125.46	119.40
1	A	3	ARG	NH1-CZ-NH2	5.35	125.28	119.40
1	B	5	TYR	CB-CG-CD2	-5.35	117.79	121.00
1	B	49[A]	ASP	CB-CG-OD2	5.30	123.08	118.30
1	B	49[B]	ASP	CB-CG-OD2	5.30	123.08	118.30
1	A	158	ASP	CB-CG-OD2	5.26	123.03	118.30
1	B	106	ASP	CB-CG-OD1	5.23	123.00	118.30
1	A	48	TYR	CB-CG-CD2	5.12	124.07	121.00
1	B	134	ARG	NE-CZ-NH1	-5.12	117.74	120.30
1	B	110	ARG	NE-CZ-NH1	5.00	122.80	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1988	0	1965	13	0
1	B	2101	0	2062	44	0
2	A	4	0	0	0	0
3	A	7	0	3	3	0
4	B	22	0	7	12	0
5	A	337	0	0	6	0
5	B	325	0	0	6	0
All	All	4784	0	4037	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:53:LYS:HG3	5:B:4549:HOH:O	1.84	0.78
1:A:17:LEU:HD21	1:B:86:LEU:HD21	1.68	0.74
1:B:171[B]:GLY:HA3	5:B:4554:HOH:O	1.91	0.69
1:B:175[B]:THR:HB	1:B:210[B]:GLY:O	1.93	0.68
1:A:237:LYS:HG3	5:A:6203:HOH:O	1.94	0.67
1:B:168[A]:TRP:HE3	1:B:174[A]:LYS:HD3	1.59	0.67
1:A:236:LEU:HB2	5:A:6203:HOH:O	1.95	0.67
1:B:232:GLY:H	4:B:4400[A]:2PG:H32	1.59	0.66
3:A:3300:3PY:H31	5:A:6157:HOH:O	1.95	0.65
1:B:65:ASN:HD21	1:B:112:LYS:NZ	1.97	0.63
1:B:156:ASN:HD22	1:B:158:ASP:H	1.46	0.63
1:B:232:GLY:N	4:B:4400[A]:2PG:H32	2.16	0.60
4:B:4400[A]:2PG:H31	4:B:4400[A]:2PG:O4P	2.00	0.58
1:B:168[A]:TRP:CE3	1:B:174[A]:LYS:HD3	2.39	0.57
1:B:50:HIS:HD2	5:B:4564:HOH:O	1.87	0.56
1:B:131:LEU:HD13	1:B:174[A]:LYS:HD2	1.87	0.55
1:B:165[B]:GLU:OE1	4:B:4400[B]:2PG:H31	2.07	0.54
1:A:65:ASN:HA	5:A:6269:HOH:O	2.08	0.54
1:B:166[A]:PRO:O	1:B:170[A]:ILE:HD13	2.09	0.52
1:B:232:GLY:H	4:B:4400[A]:2PG:C3	2.23	0.52
1:B:12:LYS:NZ	4:B:4400[B]:2PG:H32	2.25	0.52
1:B:156:ASN:ND2	1:B:158:ASP:H	2.07	0.52
1:A:12:LYS:HZ1	3:A:3300:3PY:C3	2.24	0.51
1:B:164:TYR:CE2	1:B:166[B]:PRO:HG3	2.45	0.51
1:B:12:LYS:HZ1	4:B:4400[B]:2PG:C2	2.24	0.51
1:B:65:ASN:HD21	1:B:112:LYS:HZ3	1.59	0.50
1:B:96[B]:PHE:CZ	1:B:100:LYS:HD2	2.45	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:157:PHE:CE1	1:B:200[B]:GLN:HG2	2.47	0.49
1:B:49[B]:ASP:OD2	1:B:53:LYS:HE3	2.12	0.49
1:B:12:LYS:HZ1	4:B:4400[B]:2PG:C3	2.25	0.48
1:B:232:GLY:CA	4:B:4400[A]:2PG:H32	2.43	0.48
1:B:104[B]:GLU:HG2	5:B:4709:HOH:O	2.14	0.48
1:B:96[B]:PHE:HB2	1:B:167[B]:LEU:HD22	1.96	0.48
1:A:222:GLN:HG2	5:A:6174:HOH:O	2.14	0.47
1:B:96[B]:PHE:CD1	1:B:167[B]:LEU:HD22	2.50	0.47
4:B:4400[A]:2PG:H31	4:B:4400[A]:2PG:O2P	2.08	0.47
1:B:232:GLY:HA3	4:B:4400[A]:2PG:H32	1.97	0.46
1:A:193:LYS:HG3	1:A:198:GLU:HA	1.97	0.46
1:B:165[B]:GLU:HG2	1:B:170[B]:ILE:HD11	1.96	0.46
1:B:134:ARG:HH11	1:B:134:ARG:HD2	1.60	0.46
1:B:233:ASN:ND2	5:B:4516:HOH:O	2.50	0.45
1:A:136:GLN:HB3	1:A:138:LYS:HE3	1.98	0.45
1:B:134:ARG:HD3	1:B:168[A]:TRP:CD2	2.52	0.45
1:B:233:ASN:ND2	4:B:4400[A]:2PG:O4P	2.49	0.45
1:B:175[B]:THR:HG21	1:B:216[B]:ASN:ND2	2.32	0.45
1:A:100:LYS:NZ	5:A:6331:HOH:O	2.49	0.44
1:B:176[B]:ALA:HB3	1:B:181:ALA:HB2	1.99	0.44
1:A:17:LEU:HD11	1:B:86:LEU:HD23	1.99	0.44
1:A:230:LEU:O	3:A:3300:3PY:O2	2.36	0.44
1:A:48:TYR:CZ	1:A:88:ILE:HG12	2.53	0.43
1:B:131:LEU:HA	1:B:168[A]:TRP:CB	2.48	0.43
1:B:118:LYS:NZ	5:B:4599:HOH:O	2.48	0.43
1:B:49[B]:ASP:CG	1:B:53:LYS:HE3	2.40	0.42
1:B:169[A]:ALA:C	1:B:170[A]:ILE:HD12	2.39	0.42
1:B:17:LEU:O	1:B:50:HIS:HE1	2.03	0.42
1:B:175[B]:THR:HG21	1:B:211[B]:SER:O	2.20	0.41
1:B:6:PHE:O	1:B:228:GLY:HA3	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	255/248 (103%)	250 (98%)	5 (2%)	0	100	100
1	B	272/248 (110%)	264 (97%)	6 (2%)	2 (1%)	22	4
All	All	527/496 (106%)	514 (98%)	11 (2%)	2 (0%)	47	10

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	211[A]	SER
1	B	211[B]	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/220 (103%)	226 (100%)	1 (0%)	91	74
1	B	240/220 (109%)	235 (98%)	5 (2%)	53	14
All	All	467/440 (106%)	461 (99%)	6 (1%)	67	32

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	229	PHE
1	B	98	ARG
1	B	132	GLU
1	B	156	ASN
1	B	190	LYS
1	B	199	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	120	ASN
1	B	29	ASN
1	B	50	HIS
1	B	56	GLN
1	B	65	ASN
1	B	71	ASN
1	B	120	ASN
1	B	156	ASN
1	B	185	HIS
1	B	233	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	2PG	B	4400[A]	-	9,10,10	4.01	3 (33%)	11,14,14	3.36	3 (27%)
3	3PY	A	3300	-	5,6,6	2.41	3 (60%)	6,7,7	7.09	4 (66%)
2	PO3	A	6000	-	0,3,3	-	-	0,3,3	-	-
4	2PG	B	4400[B]	1	9,10,10	2.56	4 (44%)	11,14,14	4.87	7 (63%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	2PG	B	4400[A]	-	-	4/11/11/11	-
4	2PG	B	4400[B]	1	1/1/3/3	4/11/11/11	-
3	3PY	A	3300	-	-	1/5/6/6	-

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	4400[A]	2PG	C2-C1	8.79	1.60	1.52
4	B	4400[A]	2PG	P-O1P	6.76	1.72	1.59
4	B	4400[B]	2PG	P-O1P	5.54	1.69	1.59
3	A	3300	3PY	O1-C1	3.96	1.33	1.22
4	B	4400[A]	2PG	O1-C1	3.34	1.32	1.22
4	B	4400[B]	2PG	C2-C1	2.80	1.54	1.52
4	B	4400[B]	2PG	O1-C1	2.75	1.30	1.22
3	A	3300	3PY	O3-C2	2.41	1.28	1.23
4	B	4400[B]	2PG	P-O2P	2.39	1.58	1.50
3	A	3300	3PY	C2-C1	2.36	1.56	1.53

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	3300	3PY	O3-C2-C1	14.61	140.60	119.43
4	B	4400[B]	2PG	O1-C1-C2	10.50	147.36	122.57
4	B	4400[B]	2PG	O2-C1-O1	-7.87	106.23	124.09
4	B	4400[A]	2PG	O2-C1-O1	-7.56	106.94	124.09
4	B	4400[A]	2PG	O2-C1-C2	-7.30	92.20	113.03
3	A	3300	3PY	O4-C3-C2	-6.33	95.96	112.48
4	B	4400[B]	2PG	O2-C1-C2	-6.08	95.69	113.03
3	A	3300	3PY	O3-C2-C3	-5.93	111.33	120.50
4	B	4400[B]	2PG	O3-C3-C2	-4.58	100.10	111.73
4	B	4400[B]	2PG	P-O1P-C2	4.06	132.35	123.04
3	A	3300	3PY	O2-C1-O1	-3.10	116.51	123.61
4	B	4400[A]	2PG	P-O1P-C2	-2.19	118.02	123.04
4	B	4400[B]	2PG	O1P-P-O2P	-2.12	101.21	109.39
4	B	4400[B]	2PG	O4P-P-O1P	2.02	115.05	105.99

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	B	4400[B]	2PG	C2

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	4400[A]	2PG	C1-C2-C3-O3
4	B	4400[A]	2PG	O1P-C2-C3-O3
4	B	4400[A]	2PG	C3-C2-O1P-P
4	B	4400[B]	2PG	O2-C1-C2-O1P
4	B	4400[B]	2PG	O1P-C2-C3-O3
4	B	4400[B]	2PG	C3-C2-O1P-P
4	B	4400[B]	2PG	O2-C1-C2-C3
4	B	4400[A]	2PG	O1-C1-C2-C3
3	A	3300	3PY	O1-C1-C2-C3

There are no ring outliers.

3 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	4400[A]	2PG	8	0
3	A	3300	3PY	3	0
4	B	4400[B]	2PG	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	246/248 (99%)	-0.22	6 (2%) 59 55	7, 12, 25, 46	0
1	B	246/248 (99%)	-0.10	9 (3%) 41 38	7, 12, 31, 51	0
All	All	492/496 (99%)	-0.16	15 (3%) 50 47	7, 12, 30, 51	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	175[A]	THR	5.3
1	B	170[A]	ILE	4.9
1	A	172	THR	4.6
1	A	171	GLY	4.3
1	B	174[A]	LYS	4.3
1	A	175	THR	4.1
1	B	169[A]	ALA	3.1
1	A	170	ILE	2.9
1	A	173	GLY	2.9
1	A	174	LYS	2.9
1	B	199	LYS	2.6
1	B	172[A]	THR	2.4
1	B	171[A]	GLY	2.3
1	B	198	GLU	2.2
1	B	197	GLY	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO3	A	6000	4/4	0.84	0.19	23,34,43,55	0
4	2PG	B	4400[A]	11/11	0.84	0.23	11,26,35,38	11
4	2PG	B	4400[B]	11/11	0.84	0.23	19,22,26,30	11
3	3PY	A	3300	7/7	0.93	0.20	13,28,46,52	0

6.5 Other polymers [i](#)

There are no such residues in this entry.