

wwPDB X-ray Structure Validation Summary Report (i)

Jan 7, 2024 – 05:00 pm GMT

PDB ID : 508Q

Title: Crystal structure of R. ruber ADH-A, mutant Y294F, W295A

Authors: Dobritzsch, D.; Maurer, D.; Hamnevik, E.; Enugala, T.R.; Widersten, M.

Deposited on : 2017-06-14

Resolution : 2.22 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS: 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{A})}) \end{array}$
R_{free}	130704	5912 (2.24-2.20)
Clashscore	141614	6646 (2.24-2.20)
Ramachandran outliers	138981	6543 (2.24-2.20)
Sidechain outliers	138945	6544 (2.24-2.20)
RSRZ outliers	127900	5797 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	352	95%	
1	В	352	95%	
1	С	352	96%	
1	D	352	95%	
1	Е	352	94%	



Continued from previous page...

	J		\cdot \cdot \cdot
\mathbf{Mol}	Chain	ain Length	Quality of chain
1	F	352	95%
-			7%
1	G	352	95%
1	TT	1 250	9%
Ţ	H	H 352	94%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 20864 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Alcohol dehydrogenase.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	345	Total	С	N	О	S	0	0	0
1	A	340	2463	1553	432	466	12	U	0	
1	В	345	Total	С	N	О	S	0	0	0
1	Ъ	340	2463	1553	432	466	12	U		
1	С	345	Total	С	N	О	S	0	1	0
1		040	2471	1558	435	466	12	U	1	
1	D	345	Total	С	N	O	S	0	0	0
1	D	040	2463	1553	432	466	12	U		
1	E	345	Total	С	N	O	S	0	0	0
1	ш	949	2463	1553	432	466	12	U		
1	F	345	Total	С	N	O	S	0	0	0
1	I.	040	2463	1553	432	466	12	U	U	U
1	G	345	Total	С	N	Ο	\mathbf{S}	0	0	0
1	G	040	2463	1553	432	466	12	U	U	U
1	Н	345	Total	С	N	О	S	0	4	0
1	1 H	345	2495	1573	440	469	13	U	4	

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	294	PHE	TYR	engineered mutation	UNP A0A1Q8I6M1
A	295	ALA	TRP	engineered mutation	UNP A0A1Q8I6M1
A	346	THR	-	expression tag	UNP A0A1Q8I6M1
A	347	SER	-	expression tag	UNP A0A1Q8I6M1
A	348	HIS	-	expression tag	UNP A0A1Q8I6M1
A	349	HIS	-	expression tag	UNP A0A1Q8I6M1
A	350	HIS	-	expression tag	UNP A0A1Q8I6M1
A	351	HIS	-	expression tag	UNP A0A1Q8I6M1
A	352	HIS	-	expression tag	UNP A0A1Q8I6M1
В	294	PHE	TYR	engineered mutation	UNP A0A1Q8I6M1
В	295	ALA	TRP	engineered mutation	UNP A0A1Q8I6M1
В	346	THR	-	expression tag	UNP A0A1Q8I6M1
В	347	SER	-	expression tag	UNP A0A1Q8I6M1



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	348	HIS	_	expression tag	UNP A0A1Q8I6M1
В	349	HIS	-	expression tag	UNP A0A1Q8I6M1
В	350	HIS	-	expression tag	UNP A0A1Q8I6M1
В	351	HIS	_	expression tag	UNP A0A1Q8I6M1
В	352	HIS	_	expression tag	UNP A0A1Q8I6M1
С	294	PHE	TYR	engineered mutation	UNP A0A1Q8I6M1
С	295	ALA	TRP	engineered mutation	UNP A0A1Q8I6M1
С	346	THR	-	expression tag	UNP A0A1Q8I6M1
С	347	SER	-	expression tag	UNP A0A1Q8I6M1
С	348	HIS	-	expression tag	UNP A0A1Q8I6M1
С	349	HIS	-	expression tag	UNP A0A1Q8I6M1
С	350	HIS	-	expression tag	UNP A0A1Q8I6M1
С	351	HIS	-	expression tag	UNP A0A1Q8I6M1
С	352	HIS	-	expression tag	UNP A0A1Q8I6M1
D	294	PHE	TYR	engineered mutation	UNP A0A1Q8I6M1
D	295	ALA	TRP	engineered mutation	UNP A0A1Q8I6M1
D	346	THR	-	expression tag	UNP A0A1Q8I6M1
D	347	SER	-	expression tag	UNP A0A1Q8I6M1
D	348	HIS	-	expression tag	UNP A0A1Q8I6M1
D	349	HIS	-	expression tag	UNP A0A1Q8I6M1
D	350	HIS	-	expression tag	UNP A0A1Q8I6M1
D	351	HIS	-	expression tag	UNP A0A1Q8I6M1
D	352	HIS	-	expression tag	UNP A0A1Q8I6M1
E	294	PHE	TYR	engineered mutation	UNP A0A1Q8I6M1
E	295	ALA	TRP	engineered mutation	UNP A0A1Q8I6M1
E	346	THR	-	expression tag	UNP A0A1Q8I6M1
E	347	SER	-	expression tag	UNP A0A1Q8I6M1
E	348	HIS	-	expression tag	UNP A0A1Q8I6M1
E	349	HIS	_	expression tag	UNP A0A1Q8I6M1
E	350	HIS	_	expression tag	UNP A0A1Q8I6M1
Е	351	HIS	_	expression tag	UNP A0A1Q8I6M1
E	352	HIS	_	expression tag	UNP A0A1Q8I6M1
F	294	PHE	TYR	engineered mutation	UNP A0A1Q8I6M1
F	295	ALA	TRP	engineered mutation	UNP A0A1Q8I6M1
F	346	THR	-	expression tag	UNP A0A1Q8I6M1
F	347	SER	-	expression tag	UNP A0A1Q8I6M1
F	348	HIS	-	expression tag	UNP A0A1Q8I6M1
F	349	HIS	-	expression tag	UNP A0A1Q8I6M1
F	350	HIS	-	expression tag	UNP A0A1Q8I6M1
F	351	HIS	-	expression tag	UNP A0A1Q8I6M1
F	352	HIS		expression tag	UNP A0A1Q8I6M1
G	294	PHE	TYR	engineered mutation	UNP A0A1Q8I6M1



Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
G	295	ALA	TRP	engineered mutation	UNP A0A1Q8I6M1
G	346	THR	-	expression tag	UNP A0A1Q8I6M1
G	347	SER	-	expression tag	UNP A0A1Q8I6M1
G	348	HIS	-	expression tag	UNP A0A1Q8I6M1
G	349	HIS	-	expression tag	UNP A0A1Q8I6M1
G	350	HIS	-	expression tag	UNP A0A1Q8I6M1
G	351	HIS	-	expression tag	UNP A0A1Q8I6M1
G	352	HIS	-	expression tag	UNP A0A1Q8I6M1
Н	294	PHE	TYR	engineered mutation	UNP A0A1Q8I6M1
Н	295	ALA	TRP	engineered mutation	UNP A0A1Q8I6M1
Н	346	THR	-	expression tag	UNP A0A1Q8I6M1
Н	347	SER	-	expression tag	UNP A0A1Q8I6M1
Н	348	HIS	-	expression tag	UNP A0A1Q8I6M1
Н	349	HIS	-	expression tag	UNP A0A1Q8I6M1
Н	350	HIS	-	expression tag	UNP A0A1Q8I6M1
Н	351	HIS	-	expression tag	UNP A0A1Q8I6M1
Н	352	HIS	-	expression tag	UNP A0A1Q8I6M1

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Zn 2 2	0	0
2	В	2	Total Zn 2 2	0	0
2	С	2	Total Zn 2 2	0	0
2	D	2	Total Zn 2 2	0	0
2	Е	2	Total Zn 2 2	0	0
2	F	2	Total Zn 2 2	0	0
2	G	2	Total Zn 2 2	0	0
2	Н	2	Total Zn 2 2	0	0

• Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).





Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf				
3	A	1	Total	С	N	О	Р	0	0				
3	A	1	44	21	7	14	2	0	0				
3	В	1	Total	С	N	О	Р	0	0				
3	Б	1	44	21	7	14	2	U	0				
3	С	1	Total	С	N	О	Р	0	0				
3		1	44	21	7	14	2	U	U				
3	D	1	Total	С	N	О	Р	0	0				
3	ע	1	44	21	7	14	2	0	U				
3	Е	1	Total	С	N	О	Р	0	0				
9	l Li	1	44	21	7	14	2	0					
3	F	1	Total	С	N	О	Р	0	0				
3	I.	1	44	21	7	14	2	U	U				
3	С	С	C	С	G	1	Total	С	N	О	Р	0	0
	G	G 1	44	21	7	14	2	U					
3	Н	1	Total	С	N	О	Р	0	0				
3 H	11	1	44	21	7	14	2	0	Ü				

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	93	Total O 93 93	0	0
4	В	108	Total O 108 108	0	0
4	С	89	Total O 89 89	0	0
4	D	120	Total O 120 120	0	0



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	Е	90	Total O 90 90	0	0
4	F	112	Total O 112 112	0	0
4	G	61	Total O 61 61	0	0
4	Н	79	Total O 79 79	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Alcohol dehydrogenase




• Molecule 1: Alcohol dehydrogenase



• Molecule 1: Alcohol dehydrogenase



• Molecule 1: Alcohol dehydrogenase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	103.69Å 109.46Å 256.65Å	Donositon
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.64 - 2.22	Depositor
Resolution (A)	29.64 - 2.22	EDS
% Data completeness	99.3 (29.64-2.22)	Depositor
(in resolution range)	99.3 (29.64-2.22)	EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.74 (at 2.22Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
D D.	0.194 , 0.216	Depositor
R, R_{free}	0.199 , 0.219	DCC
R_{free} test set	7309 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	30.3	Xtriage
Anisotropy	0.551	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.31, 30.9	EDS
L-test for twinning ²	$ < L > = 0.46, < L^2> = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	20864	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 8.47% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAD, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Во	nd lengths	В	ond angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.53	$1/2512 \ (0.0\%)$	0.73	$2/3429 \ (0.1\%)$
1	В	0.48	0/2512	0.69	1/3429~(0.0%)
1	С	0.48	0/2523	0.70	0/3443
1	D	0.49	0/2512	0.71	0/3429
1	Е	0.49	0/2512	0.77	5/3429~(0.1%)
1	F	0.49	0/2512	0.70	1/3429~(0.0%)
1	G	0.47	0/2512	0.72	2/3429~(0.1%)
1	Н	0.52	0/2548	0.73	3/3477~(0.1%)
All	All	0.49	1/20143 (0.0%)	0.72	14/27494 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$\operatorname{Ideal}(\text{\AA})$
1	A	232	ARG	CZ-NH1	6.11	1.41	1.33

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	Е	47	MET	CA-CB-CG	11.47	132.80	113.30
1	Е	206	ASP	CB-CG-OD1	-10.95	108.44	118.30
1	G	222	LYS	CD-CE-NZ	10.18	135.10	111.70
1	Е	125	MET	CA-CB-CG	-8.51	98.83	113.30
1	Е	125	MET	CG-SD-CE	-7.75	87.80	100.20

There are no chirality outliers.

There are no planarity outliers.



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2463	0	2460	3	1
1	В	2463	0	2461	2	1
1	С	2471	0	2474	2	0
1	D	2463	0	2460	3	1
1	Ε	2463	0	2460	5	0
1	F	2463	0	2460	3	0
1	G	2463	0	2461	5	0
1	Н	2495	0	2495	6	1
2	A	2	0	0	0	0
2	В	2	0	0	0	0
2	С	2	0	0	0	0
2	D	2	0	0	0	0
2	Ε	2	0	0	0	0
2	F	2	0	0	0	0
2	G	2	0	0	0	0
2	Н	2	0	0	0	0
3	A	44	0	26	0	0
3	В	44	0	26	0	0
3	С	44	0	26	0	0
3	D	44	0	26	0	0
3	Ε	44	0	26	0	0
3	F	44	0	26	0	0
3	G	44	0	26	0	0
3	Н	44	0	26	0	0
4	A	93	0	0	1	0
4	В	108	0	0	0	0
4	С	89	0	0	1	0
4	D	120	0	0	0	0
4	Ε	90	0	0	0	0
4	F	112	0	0	0	0
4	G	61	0	0	0	0
4	Н	79	0	0	1	0
All	All	20864	0	19939	29	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 29 close contacts within the same asymmetric unit are listed below, sorted by their



clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:C:213[B]:ARG:NH1	4:C:601:HOH:O	2.24	0.69
1:H:208:ARG:NH2	4:H:602:HOH:O	2.39	0.55
1:A:147:ALA:HB3	4:A:620:HOH:O	2.09	0.52
1:E:47:MET:HG3	1:E:52:TYR:HD2	1.77	0.50
1:H:165:VAL:HG11	1:H:265:HIS:CG	2.48	0.49

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{aligned} & ext{Interatomic} \ & ext{distance} \ & ext{(Å)} \end{aligned}$	Clash overlap (Å)
1:D:207:ASP:OD1	1:H:213[A]:ARG:NH1[2_485]	2.01	0.19
1:A:229:ASP:OD2	1:B:20:THR:N[3_545]	2.13	0.07

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	343/352 (97%)	338 (98%)	5 (2%)	0	100 100
1	В	343/352~(97%)	338 (98%)	5 (2%)	0	100 100
1	С	344/352 (98%)	340 (99%)	4 (1%)	0	100 100
1	D	343/352 (97%)	339 (99%)	4 (1%)	0	100 100
1	E	343/352 (97%)	339 (99%)	4 (1%)	0	100 100
1	F	343/352 (97%)	339 (99%)	4 (1%)	0	100 100
1	G	343/352 (97%)	337 (98%)	6 (2%)	0	100 100
1	Н	347/352 (99%)	342 (99%)	5 (1%)	0	100 100
All	All	2749/2816 (98%)	2712 (99%)	37 (1%)	0	100 100

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	247/254 (97%)	244 (99%)	3 (1%)	71 82
1	В	247/254 (97%)	244 (99%)	3 (1%)	71 82
1	C	248/254 (98%)	244 (98%)	4 (2%)	62 75
1	D	247/254 (97%)	245 (99%)	2 (1%)	81 89
1	E	247/254 (97%)	243 (98%)	4 (2%)	62 75
1	F	247/254 (97%)	245 (99%)	2 (1%)	81 89
1	G	247/254 (97%)	245 (99%)	2 (1%)	81 89
1	Н	251/254 (99%)	249 (99%)	2 (1%)	81 89
All	All	1981/2032 (98%)	1959 (99%)	22 (1%)	73 84

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Ε	272	HIS
1	F	340	ARG
1	F	272	HIS
1	G	272	HIS
1	С	62	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	HIS
1	Н	5	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 24 ligands modelled in this entry, 16 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type Chain Re		Res	Link	Во	Bond lengths			Bond angles		
MIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
3	NAD	Н	503	-	42,48,48	0.84	1 (2%)	50,73,73	1.32	7 (14%)	
3	NAD	В	503	-	42,48,48	0.87	2 (4%)	50,73,73	1.33	8 (16%)	
3	NAD	С	503	-	42,48,48	0.86	2 (4%)	50,73,73	1.28	6 (12%)	
3	NAD	D	503	-	42,48,48	0.89	3 (7%)	50,73,73	1.40	9 (18%)	
3	NAD	Е	503	-	42,48,48	0.81	1 (2%)	50,73,73	1.28	6 (12%)	
3	NAD	G	503	-	42,48,48	0.87	1 (2%)	50,73,73	1.32	6 (12%)	
3	NAD	F	503	-	42,48,48	0.84	2 (4%)	50,73,73	1.35	8 (16%)	
3	NAD	A	503	-	42,48,48	0.86	2 (4%)	50,73,73	1.31	6 (12%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	\mathbf{Type}	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	Н	503	1	-	7/26/62/62	0/5/5/5
3	NAD	В	503	-	-	7/26/62/62	0/5/5/5
3	NAD	С	503	-	-	8/26/62/62	0/5/5/5
3	NAD	D	503	-	-	7/26/62/62	0/5/5/5



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	E	503	-	-	8/26/62/62	0/5/5/5
3	NAD	G	503	-	-	7/26/62/62	0/5/5/5
3	NAD	F	503	-	-	8/26/62/62	0/5/5/5
3	NAD	A	503	-	-	8/26/62/62	0/5/5/5

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(A)
3	G	503	NAD	C5A-C4A	2.78	1.48	1.40
3	Н	503	NAD	C5A-C4A	2.50	1.47	1.40
3	С	503	NAD	C5A-C4A	2.42	1.47	1.40
3	Е	503	NAD	C5A-C4A	2.37	1.47	1.40
3	В	503	NAD	O4D-C1D	2.36	1.44	1.41

The worst 5 of 56 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
3	Е	503	NAD	N3A-C2A-N1A	-4.20	122.11	128.68
3	A	503	NAD	N3A-C2A-N1A	-4.03	122.38	128.68
3	F	503	NAD	N3A-C2A-N1A	-4.02	122.40	128.68
3	С	503	NAD	N3A-C2A-N1A	-4.01	122.42	128.68
3	D	503	NAD	N3A-C2A-N1A	-3.96	122.48	128.68

There are no chirality outliers.

5 of 60 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	503	NAD	O4D-C1D-N1N-C2N
3	A	503	NAD	O4D-C1D-N1N-C6N
3	A	503	NAD	C2D-C1D-N1N-C2N
3	A	503	NAD	C2D-C1D-N1N-C6N
3	В	503	NAD	O4D-C1D-N1N-C2N


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.




Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q < 0.9
1	A	345/352~(98%)	-0.06	15 (4%) 35 33	19, 34, 65, 98	0
1	В	345/352~(98%)	-0.12	10 (2%) 51 49	17, 33, 61, 82	0
1	С	345/352 (98%)	0.02	15 (4%) 35 33	19, 36, 76, 109	0
1	D	345/352 (98%)	-0.18	8 (2%) 60 58	19, 32, 60, 98	0
1	E	345/352 (98%)	-0.10	15 (4%) 35 33	18, 35, 65, 93	0
1	F	345/352~(98%)	-0.13	13 (3%) 40 38	18, 30, 63, 95	0
1	G	345/352 (98%)	0.22	26 (7%) 14 12	22, 45, 87, 120	0
1	Н	345/352 (98%)	0.42	32 (9%) 8 7	19, 43, 91, 118	0
All	All	2760/2816 (98%)	0.01	134 (4%) 29 28	17, 35, 76, 120	0

The worst 5 of 134 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Н	43	PHE	6.5
1	Н	38[A]	CYS	5.9
1	Н	49	ALA	5.9
1	Н	50	ALA	5.5
1	G	51	GLN	4.9

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathrm{A}}^2)$	Q<0.9
2	ZN	D	501	1/1	0.70	0.11	75,75,75,75	0
2	ZN	A	501	1/1	0.77	0.09	74,74,74,74	0
2	ZN	Ε	501	1/1	0.84	0.10	71,71,71,71	0
2	ZN	F	501	1/1	0.84	0.06	72,72,72,72	0
2	ZN	В	501	1/1	0.85	0.07	84,84,84,84	0
2	ZN	Н	501	1/1	0.86	0.60	61,61,61,61	1
2	ZN	G	501	1/1	0.90	0.07	86,86,86,86	0
2	ZN	С	501	1/1	0.94	0.06	86,86,86,86	0
3	NAD	G	503	44/44	0.95	0.11	24,31,33,35	0
3	NAD	С	503	44/44	0.96	0.11	22,26,27,27	0
3	NAD	A	503	44/44	0.96	0.11	23,28,30,32	0
3	NAD	Η	503	44/44	0.96	0.10	26,31,33,34	0
3	NAD	D	503	44/44	0.97	0.11	24,25,28,28	0
3	NAD	Ε	503	44/44	0.97	0.09	23,25,27,28	0
3	NAD	В	503	44/44	0.98	0.11	23,26,27,28	0
3	NAD	F	503	44/44	0.98	0.08	22,23,24,26	0
2	ZN	Н	502	1/1	0.99	0.04	25,25,25,25	0
2	ZN	A	502	1/1	0.99	0.04	27,27,27,27	0
2	ZN	F	502	1/1	0.99	0.03	25,25,25,25	0
2	ZN	Ε	502	1/1	1.00	0.05	22,22,22,22	0
2	ZN	В	502	1/1	1.00	0.03	28,28,28,28	0
2	ZN	D	502	1/1	1.00	0.02	24,24,24,24	0
2	ZN	С	502	1/1	1.00	0.05	27,27,27,27	0
2	ZN	G	502	1/1	1.00	0.03	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.










Electron density around NAD A 503:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NAD H 503:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around NAD D 503: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive) Electron density around NAD E 503:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around NAD B 503: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.

