

wwPDB NMR Structure Validation Summary Report (i)

Jun 6, 2023 – 01:47 pm BST

PDB ID : 5O9B BMRB ID : 34149

Title : Solution NMR structure of human GATA2 C-terminal zinc finger domain

Authors: Nurmohamed, S.S.; Broadhurst, R.W.; May, G.; Enver, T.

Deposited on : 2017-06-16

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

 $\begin{array}{ccc} wwPDB\text{-ShiftChecker} &:& v1.2\\ BMRB \ Restraints \ Analysis &:& v1.2 \end{array}$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 80%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$ m NMR~archive \ (\#Entries)$		
Clashscore	158937	12864		
Ramachandran outliers	154571	11451		
Sidechain outliers	154315	11428		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mo	Chain	Length	Quality of chain			
1	A	68	21%	32%	•	44%



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 2 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues					
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model		
1	A:5-A:42 (38)	0.26	2		

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 2 single-model clusters were found.

Cluster number	Models		
1	3, 6, 7, 11, 12, 14, 17, 18		
2	1, 4, 9, 10, 15, 20		
3	2, 5, 13, 19		
Single-model clusters	8; 16		



3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 1056 atoms, of which 519 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called Endothelial transcription factor GATA-2.

Mol	Chain	Residues	Atoms				Trace		
1	Λ	60	Total	С	Н	N	О	S	0
1		68	1055	326	519	109	94	7	0

There are 19 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MET	-	initiating methionine	UNP P23769
A	-17	ALA	-	expression tag	UNP P23769
A	-16	HIS	-	expression tag	UNP P23769
A	-15	HIS	-	expression tag	UNP P23769
A	-14	HIS	-	expression tag	UNP P23769
A	-13	HIS	-	expression tag	UNP P23769
A	-12	HIS	-	expression tag	UNP P23769
A	-11	HIS	-	expression tag	UNP P23769
A	-10	SER	-	expression tag	UNP P23769
A	-9	SER	-	expression tag	UNP P23769
A	-8	GLY	-	expression tag	UNP P23769
A	-7	LEU	-	expression tag	UNP P23769
A	-6	GLU	-	expression tag	UNP P23769
A	-5	VAL	-	expression tag	UNP P23769
A	-4	LEU	-	expression tag	UNP P23769
A	-3	PHE		expression tag	UNP P23769
A	-2	GLN	-	expression tag	UNP P23769
A	-1	GLY	-	expression tag	UNP P23769
A	0	PRO	-	expression tag	UNP P23769

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms
2	A	1	Total Zn
	11		1 1




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


• Molecule 1: Endothelial transcription factor GATA-2



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 2. Colouring as in section 4.1 above.

• Molecule 1: Endothelial transcription factor GATA-2





Refinement protocol and experimental data overview (i) 5



The models were refined using the following method: simulated annealing.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: structures with the lowest energy.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
ARIA	refinement	2.3
ARIA	structure calculation	2.3

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	594
Number of shifts mapped to atoms	594
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	80%



6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain	В	ond lengths	Bond angles		
MIOI	Chain	RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.56 ± 0.08	$0\pm1/301~(~0.1\pm~0.2\%)$	0.62 ± 0.03	$0\pm0/413~(~0.0\pm~0.1\%)$
All	All	0.57	4/6020 (0.1%)	0.62	2/8260 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0 ± 0.0	0.3 ± 0.5
All	All	0	7

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain Res Type Atoms Z Observed(Å)		$oxed{Atoms} oxed{f Z} oxed{f Observed(\AA)} oxed{f Ideal(\AA)}$		Models				
WIOI	Chain	nes	Type	Atoms		Observed(A)	ideal(A)	Worst	Total
1	A	34	TYR	CE2-CZ	-8.23	1.27	1.38	8	1
1	A	34	TYR	CE1-CZ	7.77	1.48	1.38	8	3

All unique angle outliers are listed below.

Mal	Chain	Dec	Type	$f Z = f Atoms = f Z = f Observed(^o)$		Atoms Z		$\operatorname{Ideal}({}^o)$	Mod	dels
MIOI	$\operatorname{Iol} \mid \operatorname{Chain} \mid \operatorname{Res} $	nes	es Type Atoms	Atoms		Observed()	ideai()	Worst	Total	
1	A	19	ARG	NE-CZ-NH1	5.66	123.13	120.30	12	2	

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.



Mol	Chain	Res	Type	Group	Models (Total)
1	A	34	TYR	Sidechain	4
1	A	35	TYR	Sidechain	3

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	295	278	277	10±3
All	All	5920	5560	5540	199

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

5 of 49 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	$Distance(\mathring{A})$	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:17:LEU:HG	1:A:19:ARG:NH1	0.70	2.02	18	5
1:A:33:LEU:O	1:A:37:LEU:HG	0.70	1.86	6	17
1:A:32:GLY:O	1:A:36:LYS:HD2	0.69	1.88	16	3
1:A:36:LYS:N	1:A:36:LYS:HE2	0.67	2.05	14	1
1:A:32:GLY:O	1:A:36:LYS:HD3	0.67	1.89	3	15

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Per	centiles
1	A	38/68 (56%)	33±1 (87±3%)	3±1 (7±3%)	2±1 (5±1%)	3	23
All	All	760/1360 (56%)	664 (87%)	56 (7%)	40 (5%)	3	23

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.



Mol	Chain	Res	Type	Models (Total)
1	A	8	ALA	19
1	A	11	GLN	19
1	A	9	ASN	2

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perc	entiles
1	A	33/58 (57%)	26±1 (79±4%)	7±1 (21±4%)	3	31
All	All	660/1160 (57%)	519 (79%)	141 (21%)	3	31

5 of 15 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	19	ARG	20
1	A	27	VAL	20
1	A	29	ASN	17
1	A	25	ASP	16
1	A	40	VAL	16

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.



6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 80% for the well-defined parts and 65% for the entire structure.

7.1 Chemical shift list 1

File name: working cs.cif

Chemical shift list name: CF_170605_csdep.txt

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	594
Number of shifts mapped to atoms	594
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	7

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\mathrm{C}_{\alpha}$	61	-0.20 ± 0.14	None needed ($< 0.5 \text{ ppm}$)
$^{13}C_{\beta}$	55	0.02 ± 0.11	None needed (< 0.5 ppm)
¹³ C′	0	_	None (insufficient data)
^{15}N	55	-0.04 ± 0.45	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 80%, i.e. 393 atoms were assigned a chemical shift out of a possible 494. 0 out of 5 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathbf{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	150/190 (79%)	76/77 (99%)	38/76~(50%)	36/37 (97%)
Sidechain	211/266 (79%)	142/171 (83%)	$62/78 \ (79\%)$	7/17 (41%)

Continued on next page...



Continued from previous page...

	Total	$^{1}\mathbf{H}$	$^{13}\mathbf{C}$	$^{15}\mathbf{N}$
Aromatic	32/38 (84%)	16/18 (89%)	15/17 (88%)	1/3 (33%)
Overall	393/494 (80%)	234/266 (88%)	115/171 (67%)	44/57 (77%)

7.1.4 Statistically unusual chemical shifts (i)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	15	THR	HG21	-0.83	0.08 - 2.19	-9.3
1	A	15	THR	HG22	-0.83	0.08 - 2.19	-9.3
1	A	15	THR	HG23	-0.83	0.08 - 2.19	-9.3
1	A	5	THR	HG21	-0.10	0.08 - 2.19	-5.8
1	A	5	THR	HG22	-0.10	0.08 - 2.19	-5.8
1	A	5	THR	HG23	-0.10	0.08 - 2.19	-5.8
1	A	15	THR	CG2	15.27	16.06 - 27.03	-5.7

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:





8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	1161
Intra-residue ($ i-j =0$)	438
Sequential ($ i-j =1$)	328
Medium range ($ i-j >1$ and $ i-j <5$)	162
Long range (i-j ≥5)	233
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	17.1
Number of long range restraints per residue ¹	3.4

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	72.9	0.2
0.2-0.5 (Medium)	90.2	0.5
>0.5 (Large)	66.1	3.1



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations



9 Distance violation analysis (i)

9.1 Summary of distance violations (i)


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Donatus into topo o	Count	% ¹	Vi	olated	3	Consis	tently	$\mathbf{Violated}^4$
Restraints type	Count	%0°	Count	$\%^2$	$\%^{1}$	Count	$\%^2$	$\%^1$
Intra-residue (i-j =0)	438	37.7	128	29.2	11.0	45	10.3	3.9
Backbone-Backbone	75	6.5	13	17.3	1.1	8	10.7	0.7
Backbone-Sidechain	235	20.2	67	28.5	5.8	20	8.5	1.7
Sidechain-Sidechain	128	11.0	48	37.5	4.1	17	13.3	1.5
Sequential (i-j =1)	328	28.3	93	28.4	8.0	27	8.2	2.3
Backbone-Backbone	121	10.4	26	21.5	2.2	7	5.8	0.6
Backbone-Sidechain	169	14.6	48	28.4	4.1	16	9.5	1.4
Sidechain-Sidechain	38	3.3	19	50.0	1.6	4	10.5	0.3
Medium range ($ i-j >1 \& i-j <5$)	162	14.0	67	41.4	5.8	20	12.3	1.7
Backbone-Backbone	48	4.1	9	18.8	0.8	2	4.2	0.2
Backbone-Sidechain	71	6.1	35	49.3	3.0	13	18.3	1.1
Sidechain-Sidechain	43	3.7	23	53.5	2.0	5	11.6	0.4
Long range ($ i-j \ge 5$)	233	20.1	97	41.6	8.4	32	13.7	2.8
Backbone-Backbone	22	1.9	3	13.6	0.3	0	0.0	0.0
Backbone-Sidechain	99	8.5	40	40.4	3.4	13	13.1	1.1
Sidechain-Sidechain	112	9.6	54	48.2	4.7	19	17.0	1.6
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	1161	100.0	385	33.2	33.2	124	10.7	10.7
Backbone-Backbone	266	22.9	51	19.2	4.4	17	6.4	1.5
Backbone-Sidechain	574	49.4	190	33.1	16.4	62	10.8	5.3
Sidechain-Sidechain	321	27.6	144	44.9	12.4	45	14.0	3.9

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models



9.1.1 Bar chart: Distribution of distance restraints and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Madal ID		Nun	nber o	f viola	ations	5	Mean (Å)	M (Å)	${ m SD}^6$ (Å)	Madian (8)
Model ID	IR^1	SQ^2	$ m MR^3$	LR^4	IC^5	Total	Mean (A)	Max (Å)	$SD^*(A)$	Median (Å)
1	77	52	38	65	0	232	0.51	2.88	0.53	0.3
2	77	51	37	65	0	230	0.49	2.9	0.5	0.3
3	78	49	37	66	0	230	0.51	3.02	0.52	0.31
4	75	50	36	65	0	226	0.48	2.9	0.5	0.29
5	71	52	38	61	0	222	0.49	2.9	0.49	0.29
6	77	46	39	65	0	227	0.51	2.9	0.51	0.32
7	75	49	36	64	0	224	0.51	2.97	0.52	0.3
8	71	54	41	60	0	226	0.5	2.68	0.48	0.3
9	79	52	42	62	0	235	0.49	2.91	0.51	0.29
10	70	47	34	57	0	208	0.5	2.93	0.51	0.32
11	75	57	41	67	0	240	0.5	3.02	0.52	0.3

Continued on next page...




Continued from previous page...

Model ID		Nun	nber o	f viola	ations	3	Mean (Å)	Max (Å)	\mathbf{SD}^6 (Å)	Median (Å)
Model 1D	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (A)	Max (A)	$SD^*(A)$	Median (A)
12	83	52	40	65	0	240	0.52	3.0	0.52	0.32
13	74	56	42	62	0	234	0.5	2.88	0.5	0.29
14	77	52	33	65	0	227	0.53	3.02	0.52	0.32
15	79	54	37	64	0	234	0.51	2.95	0.51	0.3
16	79	45	38	57	0	219	0.51	2.97	0.51	0.31
17	78	49	42	65	0	234	0.51	2.86	0.53	0.31
18	79	54	44	62	0	239	0.5	2.99	0.51	0.3
19	75	54	36	64	0	229	0.49	3.1	0.51	0.3
20	73	52	40	62	0	227	0.48	3.01	0.49	0.29

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation

9.2.1 Bar graph: Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right



9.3 Distance violation statistics for the ensemble (i)


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 776(IR:310, SQ:235, MR:95, LR:136, IC:0) restraints are not violated in the ensemble.

Nu	Number of violated restraints						n of the ensemble
IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Count ⁶	%
13	9	9	8	0	39	1	5.0
11	6	4	5	0	26	2	10.0
10	11	5	4	0	30	3	15.0
4	6	4	5	0	19	4	20.0
2	5	1	5	0	13	5	25.0
2	1	2	0	0	5	6	30.0
5	4	1	1	0	11	7	35.0
0	3	2	1	0	6	8	40.0
8	1	3	3	0	15	9	45.0
6	2	0	3	0	11	10	50.0
1	1	2	4	0	8	11	55.0
1	1	1	2	0	5	12	60.0
2	0	1	4	0	7	13	65.0
1	1	0	1	0	3	14	70.0
3	2	1	1	0	7	15	75.0
4	3	2	1	0	10	16	80.0
3	1	4	4	0	12	17	85.0
4	6	1	4	0	15	18	90.0
3	3	4	9	0	19	19	95.0
45	27	20	32	0	124	20	100.0

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations



9.3.1 Bar graph: Distance violation statistics for the ensemble (i)




9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram : Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)	Median (Å)
(2,396)	1:A:27:VAL:HG11	1:A:19:ARG:HB3	20	2.91	0.08	2.9
(2,396)	1:A:27:VAL:HG12	1:A:19:ARG:HB3	20	2.91	0.08	2.9
(2,396)	1:A:27:VAL:HG13	1:A:19:ARG:HB3	20	2.91	0.08	2.9
(2,162)	1:A:36:LYS:HB3	1:A:19:ARG:HD2	20	2.62	0.13	2.64
(2,20)	1:A:35:TYR:HB2	1:A:36:LYS:HB3	20	2.52	0.04	2.53
(1,102)	1:A:45:THR:H	1:A:43:PRO:HB2	20	2.51	0.31	2.34
(1,208)	1:A:9:ASN:H	1:A:27:VAL:HG21	20	2.1	0.17	2.14
(1,208)	1:A:9:ASN:H	1:A:27:VAL:HG22	20	2.1	0.17	2.14
(1,208)	1:A:9:ASN:H	1:A:27:VAL:HG23	20	2.1	0.17	2.14
(2,123)	1:A:27:VAL:HG21	1:A:9:ASN:H	20	2.0	0.17	2.04
(2,123)	1:A:27:VAL:HG22	1:A:9:ASN:H	20	2.0	0.17	2.04
(2,123)	1:A:27:VAL:HG23	1:A:9:ASN:H	20	2.0	0.17	2.04
(1,270)	1:A:29:ASN:HD21	1:A:17:LEU:HD11	20	1.9	0.56	1.58
(1,270)	1:A:29:ASN:HD21	1:A:17:LEU:HD12	20	1.9	0.56	1.58
(1,270)	1:A:29:ASN:HD21	1:A:17:LEU:HD13	20	1.9	0.56	1.58
(1,338)	1:A:29:ASN:HD22	1:A:17:LEU:HD11	20	1.89	0.28	1.81

Continued on next page...



Continued from previous page...


Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)	Median (Å)
(1,338)	1:A:29:ASN:HD22	1:A:17:LEU:HD12	20	1.89	0.28	1.81
(1,338)	1:A:29:ASN:HD22	1:A:17:LEU:HD13	20	1.89	0.28	1.81
(2,543)	1:A:32:GLY:HA3	1:A:35:TYR:HB2	20	1.59	0.12	1.58
(2,461)	1:A:43:PRO:HD2	1:A:40:VAL:HG11	20	1.57	0.08	1.61

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints (i)

9.5.1 Histogram: Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,102)	1:A:45:THR:H	1:A:43:PRO:HB2	19	3.1

Continued on next page...



Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(2,396)	1:A:27:VAL:HG11	1:A:19:ARG:HB3	3	3.02
(2,396)	1:A:27:VAL:HG12	1:A:19:ARG:HB3	3	3.02
(2,396)	1:A:27:VAL:HG13	1:A:19:ARG:HB3	3	3.02
(2,396)	1:A:27:VAL:HG11	1:A:19:ARG:HB3	11	3.02
(2,396)	1:A:27:VAL:HG12	1:A:19:ARG:HB3	11	3.02
(2,396)	1:A:27:VAL:HG13	1:A:19:ARG:HB3	11	3.02
(2,396)	1:A:27:VAL:HG11	1:A:19:ARG:HB3	14	3.02
(2,396)	1:A:27:VAL:HG12	1:A:19:ARG:HB3	14	3.02
(2,396)	1:A:27:VAL:HG13	1:A:19:ARG:HB3	14	3.02
(1,102)	1:A:45:THR:H	1:A:43:PRO:HB2	20	3.01
(2,396)	1:A:27:VAL:HG11	1:A:19:ARG:HB3	12	3.0
(2,396)	1:A:27:VAL:HG12	1:A:19:ARG:HB3	12	3.0
(2,396)	1:A:27:VAL:HG13	1:A:19:ARG:HB3	12	3.0
(2,396)	1:A:27:VAL:HG11	1:A:19:ARG:HB3	18	2.99
(2,396)	1:A:27:VAL:HG12	1:A:19:ARG:HB3	18	2.99
(2,396)	1:A:27:VAL:HG13	1:A:19:ARG:HB3	18	2.99
(2,396)	1:A:27:VAL:HG11	1:A:19:ARG:HB3	7	2.97
(2,396)	1:A:27:VAL:HG12	1:A:19:ARG:HB3	7	2.97
(2,396)	1:A:27:VAL:HG13	1:A:19:ARG:HB3	7	2.97
(2,396)	1:A:27:VAL:HG11	1:A:19:ARG:HB3	16	2.97
(2,396)	1:A:27:VAL:HG12	1:A:19:ARG:HB3	16	2.97
(2,396)	1:A:27:VAL:HG13	1:A:19:ARG:HB3	16	2.97
(1,102)	1:A:45:THR:H	1:A:43:PRO:HB2	15	2.95



10 Dihedral-angle violation analysis (i)

No dihedral-angle restraints found

