

# wwPDB X-ray Structure Validation Summary Report (i)

### Jun 16, 2020 - 12:03 am BST

PDB ID	:	50AN
$\operatorname{Title}$	:	Crystal structure of mutant AChBP in complex with glycine (T53F, Q74R,
		Y110A, I135S, G162E, S206CCP_KGTG)
Authors	:	Dawson, A.; Hunter, W.N.; de Souza, J.O.; Trumper, P.
Deposited on	:	2017-06-23
$\operatorname{Resolution}$	:	2.60  Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
$\operatorname{Refmac}$	:	5.8.0158
$\rm CCP4$	:	$7.0.044 (\mathrm{Gargrove})$
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries},{ m resolution\ range}({ m \AA}))$
R <sub>free</sub>	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455(2.60-2.60)
Sidechain outliers	138945	3455(2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
-	٨	240	2%		
	A	249	79%	•	17%
		240	2%		
	В	249	78%	6%	15%
-	~	2.4.2	2%		
1	С	249	79%	5% •	15%
	_		2%		
1	D	249	79%	••	16%
			3%		
1	Ε	249	78%	•	17%



# 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 8744 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Λ	206	Total	С	Ν	Ο	S	0	2	0
	A	200	1660	1049	276	328	7	0	Δ	0
1	В	211	Total	С	Ν	Ο	S	0	3	0
	D	211	1700	1077	283	333	7	0	J	0
1	С	212	Total	С	Ν	Ο	S	0	4	0
			1726	1091	293	334	8		1	0
1	р	208	Total	С	Ν	Ο	S	0	4	0
	D	200	1693	1075	279	331	8	0	4	0
1	Е	206	Total	С	Ν	Ο	S	0	4	0
		200	1670	1056	277	329	8	0	4	U

• Molecule 1 is a protein called Soluble acetylcholine receptor.

There are 120 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	53	PHE	THR	engineered mutation	UNP Q8WSF8
А	60	VAL	ALA	engineered mutation	UNP Q8WSF8
A	74	ARG	GLN	engineered mutation	UNP Q8WSF8
А	110	ALA	TYR	engineered mutation	UNP Q8WSF8
А	135	SER	ILE	engineered mutation	UNP Q8WSF8
А	155	VAL	ALA	engineered mutation	UNP Q8WSF8
А	162	GLU	GLY	engineered mutation	UNP Q8WSF8
А	206	LYS	SER	engineered mutation	UNP Q8WSF8
А	207	GLY	CYS	engineered mutation	UNP Q8WSF8
A	208	THR	CYS	engineered mutation	UNP Q8WSF8
А	209	GLY	PRO	engineered mutation	UNP Q8WSF8
А	237	GLU	-	expression tag	UNP Q8WSF8
A	238	ASN	-	expression tag	UNP Q8WSF8
А	239	LEU	-	expression tag	UNP Q8WSF8
А	240	TYR	-	expression tag	UNP Q8WSF8
А	241	PHE	-	expression tag	UNP Q8WSF8
А	242	GLN	-	expression tag	UNP Q8WSF8
А	243	GLY	-	expression tag	UNP Q8WSF8
А	244	HIS	-	expression tag	UNP Q8WSF8



5O	AN

Chain	Residue	Modelled	Actual	Comment	Reference
А	245	HIS	-	expression tag	UNP Q8WSF8
А	246	HIS	-	expression tag	UNP Q8WSF8
А	247	HIS	-	expression tag	UNP Q8WSF8
А	248	HIS	-	expression tag	UNP Q8WSF8
А	249	HIS	-	expression tag	UNP Q8WSF8
В	53	PHE	THR	engineered mutation	UNP Q8WSF8
В	60	VAL	ALA	engineered mutation	UNP Q8WSF8
В	74	ARG	GLN	engineered mutation	UNP Q8WSF8
В	110	ALA	TYR	engineered mutation	UNP Q8WSF8
В	135	SER	ILE	engineered mutation	UNP Q8WSF8
В	155	VAL	ALA	engineered mutation	UNP Q8WSF8
В	162	GLU	GLY	engineered mutation	UNP Q8WSF8
В	206	LYS	SER	engineered mutation	UNP Q8WSF8
В	207	GLY	CYS	engineered mutation	UNP Q8WSF8
В	208	THR	CYS	engineered mutation	UNP Q8WSF8
В	209	GLY	PRO	engineered mutation	UNP Q8WSF8
В	237	GLU	-	expression tag	UNP Q8WSF8
В	238	ASN	-	expression tag	UNP Q8WSF8
В	239	LEU	-	expression tag	UNP Q8WSF8
В	240	TYR	-	expression tag	UNP Q8WSF8
В	241	PHE	-	expression tag	UNP Q8WSF8
В	242	GLN	-	expression tag	UNP Q8WSF8
В	243	GLY	_	expression tag	UNP Q8WSF8
В	244	HIS	-	expression tag	UNP Q8WSF8
В	245	HIS	-	expression tag	UNP Q8WSF8
В	246	HIS	_	expression tag	UNP Q8WSF8
В	247	HIS	_	expression tag	UNP Q8WSF8
В	248	HIS	_	expression tag	UNP Q8WSF8
В	249	HIS	_	expression tag	UNP Q8WSF8
С	53	PHE	THR	engineered mutation	UNP Q8WSF8
С	60	VAL	ALA	engineered mutation	UNP Q8WSF8
С	74	ARG	GLN	engineered mutation	UNP Q8WSF8
С	110	ALA	TYR	engineered mutation	UNP Q8WSF8
С	135	SER	ILE	engineered mutation	UNP Q8WSF8
С	155	VAL	ALA	engineered mutation	UNP Q8WSF8
C	162	GLU	GLY	engineered mutation	UNP Q8WSF8
С	206	LYS	SER	engineered mutation	UNP Q8WSF8
C	207	GLY	CYS	engineered mutation	UNP Q8WSF8
C	208	THR	CYS	engineered mutation	UNP Q8WSF8
C	209	GLY	PRO	engineered mutation	UNP Q8WSF8
C	238	GLU	_	expression tag	UNP Q8WSF8
C	239	ASN	_	expression tag	UNP Q8WSF8



51	$\cap$	A	M
0	U.		N L

Chain	Residue	Modelled	Actual	Comment	Reference
С	240	LEU	-	expression tag	UNP Q8WSF8
С	241	TYR	-	expression tag	UNP Q8WSF8
С	242	PHE	-	expression tag	UNP Q8WSF8
С	243	GLN	-	expression tag	UNP Q8WSF8
С	244	GLY	-	expression tag	UNP Q8WSF8
С	245	HIS	-	expression tag	UNP Q8WSF8
С	246	HIS	-	expression tag	UNP Q8WSF8
С	247	HIS	-	expression tag	UNP Q8WSF8
С	248	HIS	-	expression tag	UNP Q8WSF8
С	249	HIS	-	expression tag	UNP Q8WSF8
С	250	HIS	-	expression tag	UNP Q8WSF8
D	53	PHE	THR	engineered mutation	UNP Q8WSF8
D	60	VAL	ALA	engineered mutation	UNP Q8WSF8
D	74	ARG	GLN	engineered mutation	UNP Q8WSF8
D	110	ALA	TYR	engineered mutation	UNP Q8WSF8
D	135	SER	ILE	engineered mutation	UNP Q8WSF8
D	155	VAL	ALA	engineered mutation	UNP Q8WSF8
D	162	GLU	GLY	engineered mutation	UNP Q8WSF8
D	206	LYS	SER	engineered mutation	UNP Q8WSF8
D	207	GLY	CYS	engineered mutation	UNP Q8WSF8
D	208	THR	CYS	engineered mutation	UNP Q8WSF8
D	209	GLY	PRO	engineered mutation	UNP Q8WSF8
D	237	GLU	-	expression tag	UNP Q8WSF8
D	238	ASN	-	expression tag	UNP Q8WSF8
D	239	LEU	-	expression tag	UNP Q8WSF8
D	240	TYR	-	expression tag	UNP Q8WSF8
D	241	PHE	-	expression tag	UNP Q8WSF8
D	242	GLN	-	expression tag	UNP Q8WSF8
D	243	GLY	-	expression tag	UNP Q8WSF8
D	244	HIS	-	expression tag	UNP Q8WSF8
D	245	HIS	_	expression tag	UNP Q8WSF8
D	246	HIS	-	expression tag	UNP Q8WSF8
D	247	HIS	-	expression tag	UNP Q8WSF8
D	248	HIS	_	expression tag	UNP Q8WSF8
D	249	HIS	_	expression tag	UNP Q8WSF8
Е	53	PHE	THR	engineered mutation	UNP Q8WSF8
Е	60	VAL	ALA	engineered mutation	UNP Q8WSF8
Е	74	ARG	GLN	engineered mutation	UNP Q8WSF8
Е	110	ALA	TYR	engineered mutation	UNP Q8WSF8
Е	135	SER	ILE	engineered mutation	UNP Q8WSF8
Е	155	VAL	ALA	engineered mutation	UNP Q8WSF8
Е	162	GLU	GLY	engineered mutation	UNP Q8WSF8




Chain	Residue	Modelled	Actual	Comment	Reference
Е	206	LYS	SER	engineered mutation	UNP Q8WSF8
Е	207	GLY	CYS	engineered mutation	UNP Q8WSF8
E	208	THR	CYS	engineered mutation	UNP Q8WSF8
Е	209	GLY	PRO	engineered mutation	UNP Q8WSF8
E	237	GLU	-	expression tag	UNP Q8WSF8
E	238	ASN	-	expression tag	UNP Q8WSF8
Е	239	LEU	-	expression tag	UNP Q8WSF8
E	240	TYR	-	expression tag	UNP Q8WSF8
E	241	PHE	-	expression tag	UNP Q8WSF8
E	242	GLN	-	expression tag	UNP Q8WSF8
E	243	GLY	-	expression tag	UNP Q8WSF8
E	244	HIS	-	expression tag	UNP Q8WSF8
E	245	HIS	-	expression tag	UNP Q8WSF8
Е	246	HIS	-	expression tag	UNP Q8WSF8
Е	247	HIS	-	expression tag	UNP Q8WSF8
Е	248	HIS	-	expression tag	UNP Q8WSF8
Е	249	HIS	_	expression tag	UNP Q8WSF8

• Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	1	$\begin{array}{cc} {\rm Total} & {\rm Cl} \\ 1 & 1 \end{array}$	0	0
2	А	1	Total Cl 1 1	0	0
2	D	1	Total Cl 1 1	0	0
2	С	1	Total Cl 1 1	0	0
2	Е	1	Total Cl 1 1	0	0


• Molecule 3 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
3	В	1	TotalCO422	0	0
3	С	1	TotalCO422	0	0
3	D	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 4  2  2 \end{array}$	0	0
3	Е	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0

• Molecule 4 is GLYCINE (three-letter code: GLY) (formula:  $C_2H_5NO_2$ ).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	В	1	$\begin{array}{cccc} \text{Total}  \text{C}  \text{N}  \text{O} \\ 5  2  1  2 \end{array}$	0	0
4	С	1	$\begin{array}{ccccc} \text{Total} & \text{C} & \text{N} & \text{O} \\ 5 & 2 & 1 & 2 \end{array}$	0	0
4	D	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{N} & \text{O} \\ 5 & 2 & 1 & 2 \end{array}$	0	0
4	Е	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{N} & \text{O} \\ 5 & 2 & 1 & 2 \end{array}$	0	0


• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	44	Total O 44 44	0	0
5	В	55	Total O 55 55	0	0
5	С	66	Total O 66 66	0	0
5	D	43	Total         O           43         43	0	0
5	Е	42	Total         O           42         42	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.




• Molecule 1: Soluble acetylcholine receptor



#### GLN GLN GLY HIS HIS HIS HIS HIS HIS HIS

• Molecule 1: Soluble acetylcholine receptor



#### ASN LEU TYR PHE GLN GLN HIS HIS HIS HIS HIS



# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants	89.92Å $100.27$ Å $165.24$ Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	32.90 - 2.60	Depositor
Resolution (A)	32.90 - 2.60	EDS
% Data completeness	99.9 (32.90-2.60)	Depositor
(in resolution range)	$100.0 \ (32.90-2.60)$	EDS
R <sub>merge</sub>	0.15	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$3.05 (at 2.61 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
D D .	0.202 , $0.235$	Depositor
$R, R_{free}$	0.206 , $0.239$	DCC
$R_{free}$ test set	2381 reflections $(5.10%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	26.9	Xtriage
Anisotropy	0.188	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.34, 28.9	EDS
L-test for twinning <sup>2</sup>	$ \langle L  \rangle = 0.46, \langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	8744	wwPDB-VP
Average B, all atoms $(Å^2)$	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.34% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles		
		RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.51	0/1705	0.72	1/2320~(0.0%)	
1	В	0.54	0/1751	0.76	2/2381~(0.1%)	
1	С	0.52	0/1782	0.74	2/2423~(0.1%)	
1	D	0.50	0/1746	0.72	1/2374~(0.0%)	
1	Е	0.50	0/1722	0.73	2/2344~(0.1%)	
All	All	0.52	0/8706	0.73	8/11842~(0.1%)	

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	С	25	ARG	NE-CZ-NH1	6.34	123.47	120.30
1	Е	25	ARG	NE-CZ-NH1	5.84	123.22	120.30
1	D	25	ARG	NE-CZ-NH1	5.69	123.14	120.30
1	С	192	GLU	OE1-CD-OE2	-5.27	116.97	123.30
1	В	33	ARG	CD-NE-CZ	5.19	130.87	123.60

There are no chirality outliers.

There are no planarity outliers.

# 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1660	0	1600	9	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	В	1700	0	1639	10	1
1	С	1726	0	1659	8	1
1	D	1693	0	1635	9	0
1	Ε	1670	0	1616	11	0
2	А	1	0	0	0	0
2	В	1	0	0	0	0
2	С	1	0	0	0	0
2	D	1	0	0	0	0
2	Е	1	0	0	0	0
3	А	4	0	3	1	0
3	В	4	0	3	0	0
3	С	4	0	3	1	0
3	D	4	0	3	0	0
3	Ε	4	0	3	1	0
4	В	5	0	2	0	0
4	С	5	0	2	0	0
4	D	5	0	2	0	0
4	Ε	5	0	2	0	0
5	А	44	0	0	0	0
5	В	55	0	0	0	0
5	С	66	0	0	0	1
5	D	43	0	0	0	0
5	Ε	42	0	0	2	0
All	All	8744	0	8172	40	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 40 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)	
1:B:114[B]:ARG:HH11	1:B:114[B]:ARG:HG3	1.15	1.12	
1:A:114[B]:ARG:HG3	1:A:114[B]:ARG:HH11	0.98	1.07	
1:A:114[B]:ARG:CG	1:A:114[B]:ARG:HH11	1.79	0.94	
1:E:114[B]:ARG:HH11	1:E:114[B]:ARG:HG3	1.29	0.94	
1:A:114[B]:ARG:HG3	1:A:114[B]:ARG:NH1	1.78	0.88	

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)	
5:C:718:HOH:O	5:C:718:HOH:O[2_575]	1.50	0.70	
1:B:179:GLN:NE2	1:C:201:GLN:OE1[3_745]	2.18	0.02	

### 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percer	ntiles
1	А	205/249~(82%)	200~(98%)	5(2%)	0	100	100
1	В	210/249~(84%)	206~(98%)	4 (2%)	0	100	100
1	С	212/249~(85%)	207~(98%)	4 (2%)	1 (0%)	29	52
1	D	208/249~(84%)	204~(98%)	4 (2%)	0	100	100
1	Ε	208/249~(84%)	204~(98%)	4 (2%)	0	100	100
All	All	1043/1245~(84%)	1021~(98%)	21 (2%)	1 (0%)	51	75

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	С	248	HIS

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	189/222~(85%)	186~(98%)	3(2%)	62 82
1	В	192/222~(86%)	188 (98%)	4 (2%)	53 77



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	С	196/222~(88%)	192~(98%)	4 (2%)	55 78	
1	D	193/222~(87%)	189~(98%)	4 (2%)	53 77	,
1	Ε	191/222~(86%)	188~(98%)	3~(2%)	62 82	
All	All	961/1110 (87%)	943~(98%)	18 (2%)	57 79	

Continued from previous page...

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	$\mathbf{Res}$	Type
1	С	206	LYS
1	С	208	THR
1	D	225	ARG
1	В	133	MET
1	С	20	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	$\mathbf{Res}$	Type
1	D	20	GLN
1	Е	55	GLN
1	D	91	ASN
1	В	201	GLN
1	D	87	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.



# 5.6 Ligand geometry (i)

Of 14 ligands modelled in this entry, 5 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	ol Type Chain Res Link		Bond lengths			Bond angles				
	Type	Cham	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2
3	ACT	В	603	-	1,3,3	0.32	0	$_{0,3,3}$	0.00	-
3	ACT	С	603	-	1,3,3	0.77	0	$_{0,3,3}$	0.00	-
3	ACT	D	603	-	1,3,3	0.79	0	$_{0,3,3}$	0.00	-
3	ACT	А	302	-	1,3,3	0.31	0	$_{0,3,3}$	0.00	-
3	ACT	Е	301	_	1,3,3	0.13	0	$_{0,3,3}$	0.00	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	С	603	ACT	1	0
3	А	302	ACT	1	0
3	Е	301	ACT	1	0

# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	А	1



All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	224:ARG	С	225:ARG	Ν	3.29



# 6 Fit of model and data (i)

# 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ $>$ 2	$OWAB(Å^2)$	Q<0.9
1	А	206/249~(82%)	-0.42	4 (1%) 66 62	14,  25,  56,  80	0
1	В	211/249~(84%)	-0.30	4 (1%) 66 62	12, 23, 59, 88	0
1	С	212/249~(85%)	-0.45	4 (1%) 66 62	12, 22, 62, 95	0
1	D	208/249~(83%)	-0.40	5 (2%) 59 53	13, 26, 59, 105	0
1	Ε	206/249~(82%)	-0.32	7 (3%) 45 38	14, 28, 65, 119	0
All	All	1043/1245~(83%)	-0.38	24 (2%) 60 54	12, 25, 61, 119	0

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	227	GLY	4.6
1	Е	206	LYS	4.1
1	D	208	THR	4.1
1	Е	205	TYR	3.7
1	D	207	GLY	3.6

# 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

# 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GLY	Е	302	5/5	0.78	0.29	$57,\!60,\!64,\!65$	0
4	GLY	С	601	5/5	0.84	0.29	$54,\!56,\!57,\!61$	0
4	GLY	D	601	5/5	0.85	0.24	$45,\!50,\!52,\!55$	0
4	GLY	В	601	5/5	0.93	0.17	42,47,49,50	0
3	ACT	D	603	4/4	0.93	0.22	$23,\!24,\!24,\!24$	0
2	CL	D	602	1/1	0.95	0.06	$32,\!32,\!32,\!32$	0
2	CL	А	301	1/1	0.96	0.06	$38,\!38,\!38,\!38$	0
2	CL	В	602	1/1	0.96	0.07	$26,\!26,\!26,\!26$	0
3	ACT	С	603	4/4	0.96	0.22	$31,\!31,\!32,\!32$	0
3	ACT	А	302	4/4	0.96	0.19	$28,\!29,\!29,\!29$	0
3	ACT	Е	301	4/4	0.97	0.16	$26,\!28,\!28,\!30$	0
3	ACT	В	603	4/4	0.97	0.17	$27,\!29,\!29,\!30$	0
2	CL	Е	303	1/1	0.98	0.06	32,32,32,32	0
2	CL	С	602	1/1	0.99	0.08	27,27,27,27	0

median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

# 6.5 Other polymers (i)

There are no such residues in this entry.

