



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2022 – 01:03 am BST

PDB ID : 7OAF  
Title : conserved hypothetical protein residues 311-335 from Candidatus Magnetomorphum sp. HK-1 fused to GCN4 adaptors, mutant beta1/A, crystal form III  
Authors : Adlakha, J.; Albrecht, R.; Hartmann, M.D.  
Deposited on : 2021-04-19  
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.28  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.28


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1156 (1.46-1.46)
Clashscore	141614	1202 (1.46-1.46)
Ramachandran outliers	138981	1178 (1.46-1.46)
Sidechain outliers	138945	1178 (1.46-1.46)
RSRZ outliers	127900	1139 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	86	 6% 86% 8% 6%
1	B	86	 8% 88% 6% 6%
1	C	86	 3% 84% 10% 6%

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4618 atoms, of which 2305 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called General control transcription factor GCN4, conserved hypothetical protein residues 311-335 from Candidatus Magnetomorum sp. HK-1 fused to GCN4 adaptors, mutant beta1/A, General control transcription factor GCN4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	A	81	Total	C	H	N	O	S	10	4	0
			1454	452	755	115	128	4			
1	B	81	Total	C	H	N	O	S	10	4	0
			1465	454	764	118	126	3			
1	C	81	Total	C	H	N	O	S	10	4	0
			1463	454	764	116	126	3			

There are 69 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	278	GLY	-	expression tag	UNP P03069
A	279	GLY	-	expression tag	UNP P03069
A	280	GLY	-	expression tag	UNP P03069
A	281	SER	-	expression tag	UNP P03069
A	282	GLY	-	expression tag	UNP P03069
A	286	ILE	LEU	engineered mutation	UNP P03069
A	288	MET	ASP	engineered mutation	UNP P03069
A	290	ILE	VAL	engineered mutation	UNP P03069
A	293	ILE	LEU	engineered mutation	UNP P03069
A	297	ILE	ASN	engineered mutation	UNP P03069
A	300	ILE	LEU	engineered mutation	UNP P03069
A	304	ILE	VAL	engineered mutation	UNP P03069
A	307	ILE	LEU	engineered mutation	UNP P03069
A	320	ALA	VAL	engineered mutation	UNP A0A0N0D484
A	326	ALA	LEU	engineered mutation	UNP A0A0N0D484
A	339	ILE	LEU	engineered mutation	UNP P03069
A	341	TRP	ASP	engineered mutation	UNP P03069
A	343	ILE	VAL	engineered mutation	UNP P03069
A	346	ILE	LEU	engineered mutation	UNP P03069
A	350	ILE	ASN	engineered mutation	UNP P03069
A	353	ILE	LEU	engineered mutation	UNP P03069

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
A	357	ILE	VAL	engineered mutation	UNP P03069
A	360	ILE	LEU	engineered mutation	UNP P03069
B	278	GLY	-	expression tag	UNP P03069
B	279	GLY	-	expression tag	UNP P03069
B	280	GLY	-	expression tag	UNP P03069
B	281	SER	-	expression tag	UNP P03069
B	282	GLY	-	expression tag	UNP P03069
B	286	ILE	LEU	engineered mutation	UNP P03069
B	288	MET	ASP	engineered mutation	UNP P03069
B	290	ILE	VAL	engineered mutation	UNP P03069
B	293	ILE	LEU	engineered mutation	UNP P03069
B	297	ILE	ASN	engineered mutation	UNP P03069
B	300	ILE	LEU	engineered mutation	UNP P03069
B	304	ILE	VAL	engineered mutation	UNP P03069
B	307	ILE	LEU	engineered mutation	UNP P03069
B	320	ALA	VAL	engineered mutation	UNP A0A0N0D484
B	326	ALA	LEU	engineered mutation	UNP A0A0N0D484
B	339	ILE	LEU	engineered mutation	UNP P03069
B	341	TRP	ASP	engineered mutation	UNP P03069
B	343	ILE	VAL	engineered mutation	UNP P03069
B	346	ILE	LEU	engineered mutation	UNP P03069
B	350	ILE	ASN	engineered mutation	UNP P03069
B	353	ILE	LEU	engineered mutation	UNP P03069
B	357	ILE	VAL	engineered mutation	UNP P03069
B	360	ILE	LEU	engineered mutation	UNP P03069
C	278	GLY	-	expression tag	UNP P03069
C	279	GLY	-	expression tag	UNP P03069
C	280	GLY	-	expression tag	UNP P03069
C	281	SER	-	expression tag	UNP P03069
C	282	GLY	-	expression tag	UNP P03069
C	286	ILE	LEU	engineered mutation	UNP P03069
C	288	MET	ASP	engineered mutation	UNP P03069
C	290	ILE	VAL	engineered mutation	UNP P03069
C	293	ILE	LEU	engineered mutation	UNP P03069
C	297	ILE	ASN	engineered mutation	UNP P03069
C	300	ILE	LEU	engineered mutation	UNP P03069
C	304	ILE	VAL	engineered mutation	UNP P03069
C	307	ILE	LEU	engineered mutation	UNP P03069
C	320	ALA	VAL	engineered mutation	UNP A0A0N0D484
C	326	ALA	LEU	engineered mutation	UNP A0A0N0D484
C	339	ILE	LEU	engineered mutation	UNP P03069
C	341	TRP	ASP	engineered mutation	UNP P03069

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	343	ILE	VAL	engineered mutation	UNP P03069
C	346	ILE	LEU	engineered mutation	UNP P03069
C	350	ILE	ASN	engineered mutation	UNP P03069
C	353	ILE	LEU	engineered mutation	UNP P03069
C	357	ILE	VAL	engineered mutation	UNP P03069
C	360	ILE	LEU	engineered mutation	UNP P03069

- Molecule 2 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula:  $C_{10}H_{22}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	H			O
2	A	1	38	10	22	6	1	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	67	Total	O	0	0
			67	67		
3	B	61	Total	O	0	0
			61	61		
3	C	70	Total	O	0	0
			70	70		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: General control transcription factor GCN4, conserved hypothetical protein residues 311-335 from *Candidatus Magnetomorum* sp. HK-1 fused to GCN4 adaptors, mutant beta1/A, General control transcription factor GCN4



- Molecule 1: General control transcription factor GCN4, conserved hypothetical protein residues 311-335 from *Candidatus Magnetomorum* sp. HK-1 fused to GCN4 adaptors, mutant beta1/A, General control transcription factor GCN4



- Molecule 1: General control transcription factor GCN4, conserved hypothetical protein residues 311-335 from *Candidatus Magnetomorum* sp. HK-1 fused to GCN4 adaptors, mutant beta1/A, General control transcription factor GCN4



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	26.01Å 38.85Å 128.81Å 90.00° 96.18° 90.00°	Depositor
Resolution (Å)	33.21 – 1.45 33.21 – 1.45	Depositor EDS
% Data completeness (in resolution range)	99.7 (33.21-1.45) 99.7 (33.21-1.45)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.28 (at 1.45Å)	Xtrriage
Refinement program	REFMAC 5.8.0049 2013/06/30	Depositor
R, $R_{free}$	0.198 , 0.215 0.200 , 0.217	Depositor DCC
$R_{free}$ test set	2279 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.9	Xtrriage
Anisotropy	0.067	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.026 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4618	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
1PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.90	0/718	0.82	1/956 (0.1%)
1	B	0.98	0/720	0.82	1/959 (0.1%)
1	C	0.89	0/718	0.77	0/956
All	All	0.92	0/2156	0.80	2/2871 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	359	ARG	NE-CZ-NH1	-5.60	117.50	120.30
1	A	340	GLU	CA-CB-CG	5.04	124.48	113.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	699	755	756	12	0
1	B	701	764	765	9	0
1	C	699	764	765	10	0
2	A	16	22	22	0	0
3	A	67	0	0	2	0
3	B	61	0	0	0	0

*Continued on next page...*



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	70	0	0	1	0
All	All	2313	2305	2308	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:345[B]:GLU:HG2	3:A:556:HOH:O	1.71	0.90
1:B:290[A]:ILE:HD11	1:C:290[A]:ILE:HD11	1.56	0.87
1:A:290[A]:ILE:HD11	1:B:290[A]:ILE:HD11	1.62	0.82
1:A:290[B]:ILE:HD12	1:C:286:ILE:HD12	1.67	0.75
1:C:356:GLU:OE2	1:C:359:ARG:NH1	2.23	0.71
1:B:290[A]:ILE:CD1	1:C:290[A]:ILE:HD11	2.20	0.71
1:A:290[A]:ILE:HD11	1:C:290[A]:ILE:HD11	1.77	0.66
1:B:314:LYS:O	1:C:317:LYS:NZ	2.33	0.61
1:A:343[B]:ILE:HD11	1:B:343[B]:ILE:HD11	1.85	0.58
1:A:343[B]:ILE:CD1	1:B:343[B]:ILE:HD11	2.34	0.58
1:A:290[A]:ILE:CD1	1:B:290[A]:ILE:HD11	2.36	0.53
1:A:286:ILE:HG23	1:B:290[B]:ILE:HD11	1.92	0.51
1:A:290[A]:ILE:HD11	1:C:290[A]:ILE:CD1	2.42	0.49
1:A:306:ARG:HG3	3:A:532:HOH:O	2.13	0.47
1:A:302:ASN:HB3	1:A:306:ARG:NH2	2.33	0.44
1:C:337:LYS:HE2	3:C:452:HOH:O	2.17	0.43
1:A:290[B]:ILE:CD1	1:C:286:ILE:HD12	2.43	0.43
1:C:283:MET:CG	1:C:284:LYS:N	2.82	0.42
1:B:283:MET:HE2	1:B:283:MET:HB3	1.86	0.42

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	83/86 (96%)	83 (100%)	0	0	100	100
1	B	83/86 (96%)	83 (100%)	0	0	100	100
1	C	83/86 (96%)	83 (100%)	0	0	100	100
All	All	249/258 (96%)	249 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	78/76 (103%)	78 (100%)	0	100	100
1	B	78/76 (103%)	78 (100%)	0	100	100
1	C	78/76 (103%)	77 (99%)	1 (1%)	69	40
All	All	234/228 (103%)	233 (100%)	1 (0%)	88	80

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	362	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	1PE	A	401	-	15,15,15	0.58	0	14,14,14	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1PE	A	401	-	-	2/13/13/13	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	1PE	OH6-C15-C25-OH5
2	A	401	1PE	C15-C25-OH5-C14

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	81/86 (94%)	0.28	5 (6%) 20 22	17, 25, 43, 65	0
1	B	81/86 (94%)	0.31	7 (8%) 10 12	18, 27, 51, 66	0
1	C	81/86 (94%)	0.21	3 (3%) 41 44	18, 26, 42, 63	0
All	All	243/258 (94%)	0.27	15 (6%) 20 22	17, 26, 48, 66	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	282	GLY	6.8
1	B	283	MET	4.6
1	C	362	LYS	4.3
1	B	284	LYS	4.1
1	A	283[A]	MET	3.5
1	B	362	LYS	3.4
1	C	283	MET	3.3
1	B	288	MET	3.3
1	A	341	TRP	3.1
1	A	361	LYS	2.8
1	C	360	ILE	2.6
1	B	285	GLN	2.6
1	A	362	LYS	2.3
1	B	287	GLU	2.3
1	A	284	LYS	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	1PE	A	401	16/16	0.94	0.10	31,38,54,54	1

### 6.5 Other polymers [i](#)

There are no such residues in this entry.