

wwPDB X-ray Structure Validation Summary Report (i)

Jun 21, 2021 – 11:10 PM BST

PDB ID : 5OBH

Title : Crystal structure of glycine binding protein in complex with bicuculline

Authors: Dawson, A.; Hunter, W.N.; Jones, M.

Deposited on : 2017-06-27

Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.20

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.20

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	249	80%		17%
			3%	•	1770
1	В	249	78% 4%	••	18%
1	С	249	81%	•	17%
1	D	249	81%		18%
			4%		1070
1	Е	249	80%	••	18%



Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	3	100%



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 8813 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Soluble acetylcholine receptor.

Mol	Chain	Residues		Atoms				ZeroOcc	AltConf	Trace
1	A	207	Total	С	N	О	S	0	0	0
1	Λ	207	1649	1042	273	327	7	0	U	
1	В	203	Total	С	N	О	S	0	1	0
1	Ъ	203	1629	1031	269	322	7		1	
1	С	206	Total	С	N	О	S	0	1	0
1		200	1646	1040	273	326	7	0		
1	D	204	Total	С	N	О	S	0	0	0
1	ע	204	1633	1033	270	323	7	U	U	
1	Е	205	Total	С	N	О	S	0	0	0
1	<u> 1</u> 2	200	1637	1035	271	324	7		U	

There are 120 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	53	PHE	THR	engineered mutation	UNP Q8WSF8
A	60	VAL	ALA	conflict	UNP Q8WSF8
A	74	ARG	GLN	engineered mutation	UNP Q8WSF8
A	110	ALA	TYR	engineered mutation	UNP Q8WSF8
A	135	SER	ILE	engineered mutation	UNP Q8WSF8
A	155	VAL	ALA	conflict	UNP Q8WSF8
A	162	GLU	GLY	engineered mutation	UNP Q8WSF8
A	206	LYS	SER	engineered mutation	UNP Q8WSF8
A	207	GLY	CYS	engineered mutation	UNP Q8WSF8
A	208	THR	CYS	engineered mutation	UNP Q8WSF8
A	209	GLY	PRO	engineered mutation	UNP Q8WSF8
A	237	GLU	_	expression tag	UNP Q8WSF8
A	238	ASN	_	expression tag	UNP Q8WSF8
A	239	LEU	_	expression tag	UNP Q8WSF8
A	240	TYR	-	expression tag	UNP Q8WSF8
A	241	PHE	-	expression tag	UNP Q8WSF8
A	242	GLN	-	expression tag	UNP Q8WSF8
A	243	GLY	-	expression tag	UNP Q8WSF8
A	244	HIS	-	expression tag	UNP Q8WSF8



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	245	HIS	_	expression tag	UNP Q8WSF8
A	246	HIS	-	expression tag	UNP Q8WSF8
A	247	HIS	-	expression tag	UNP Q8WSF8
A	248	HIS	-	expression tag	UNP Q8WSF8
A	249	HIS	-	expression tag	UNP Q8WSF8
В	53	PHE	THR	engineered mutation	UNP Q8WSF8
В	60	VAL	ALA	conflict	UNP Q8WSF8
В	74	ARG	GLN	engineered mutation	UNP Q8WSF8
В	110	ALA	TYR	engineered mutation	UNP Q8WSF8
В	135	SER	ILE	engineered mutation	UNP Q8WSF8
В	155	VAL	ALA	conflict	UNP Q8WSF8
В	162	GLU	GLY	engineered mutation	UNP Q8WSF8
В	206	LYS	SER	engineered mutation	UNP Q8WSF8
В	207	GLY	CYS	engineered mutation	UNP Q8WSF8
В	208	THR	CYS	engineered mutation	UNP Q8WSF8
В	209	GLY	PRO	engineered mutation	UNP Q8WSF8
В	237	GLU	=	expression tag	UNP Q8WSF8
В	238	ASN	=	expression tag	UNP Q8WSF8
В	239	LEU	=	expression tag	UNP Q8WSF8
В	240	TYR	=	expression tag	UNP Q8WSF8
В	241	PHE	-	expression tag	UNP Q8WSF8
В	242	GLN	-	expression tag	UNP Q8WSF8
В	243	GLY	ı	expression tag	UNP Q8WSF8
В	244	HIS	-	expression tag	UNP Q8WSF8
В	245	HIS	ı	expression tag	UNP Q8WSF8
В	246	HIS	ı	expression tag	UNP Q8WSF8
В	247	HIS	-	expression tag	UNP Q8WSF8
В	248	HIS	-	expression tag	UNP Q8WSF8
В	249	HIS	1	expression tag	UNP Q8WSF8
С	53	PHE	THR	engineered mutation	UNP Q8WSF8
С	60	VAL	ALA	conflict	UNP Q8WSF8
С	74	ARG	GLN	engineered mutation	UNP Q8WSF8
С	110	ALA	TYR	engineered mutation	UNP Q8WSF8
С	135	SER	ILE	engineered mutation	UNP Q8WSF8
С	155	VAL	ALA	conflict	UNP Q8WSF8
С	162	GLU	GLY	engineered mutation	UNP Q8WSF8
С	206	LYS	SER	engineered mutation	UNP Q8WSF8
С	207	GLY	CYS	engineered mutation	UNP Q8WSF8
С	208	THR	CYS	engineered mutation	UNP Q8WSF8
С	209	GLY	PRO	engineered mutation	UNP Q8WSF8
С	237	GLU		expression tag	UNP Q8WSF8
С	238	ASN	_	expression tag	UNP Q8WSF8



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
С	239	LEU	_	expression tag	UNP Q8WSF8
С	240	TYR	_	expression tag	UNP Q8WSF8
С	241	PHE	_	expression tag	UNP Q8WSF8
С	242	GLN	_	expression tag	UNP Q8WSF8
С	243	GLY	_	expression tag	UNP Q8WSF8
С	244	HIS	_	expression tag	UNP Q8WSF8
С	245	HIS	_	expression tag	UNP Q8WSF8
С	246	HIS	-	expression tag	UNP Q8WSF8
С	247	HIS	-	expression tag	UNP Q8WSF8
С	248	HIS	-	expression tag	UNP Q8WSF8
С	249	HIS	_	expression tag	UNP Q8WSF8
D	53	PHE	THR	engineered mutation	UNP Q8WSF8
D	60	VAL	ALA	conflict	UNP Q8WSF8
D	74	ARG	GLN	engineered mutation	UNP Q8WSF8
D	110	ALA	TYR	engineered mutation	UNP Q8WSF8
D	135	SER	ILE	engineered mutation	UNP Q8WSF8
D	155	VAL	ALA	conflict	UNP Q8WSF8
D	162	GLU	GLY	engineered mutation	UNP Q8WSF8
D	206	LYS	SER	engineered mutation	UNP Q8WSF8
D	207	GLY	CYS	engineered mutation	UNP Q8WSF8
D	208	THR	CYS	engineered mutation	UNP Q8WSF8
D	209	GLY	PRO	engineered mutation	UNP Q8WSF8
D	237	GLU	-	expression tag	UNP Q8WSF8
D	238	ASN	-	expression tag	UNP Q8WSF8
D	239	LEU	_	expression tag	UNP Q8WSF8
D	240	TYR	-	expression tag	UNP Q8WSF8
D	241	PHE	_	expression tag	UNP Q8WSF8
D	242	GLN	_	expression tag	UNP Q8WSF8
D	243	GLY	_	expression tag	UNP Q8WSF8
D	244	HIS	-	expression tag	UNP Q8WSF8
D	245	HIS	-	expression tag	UNP Q8WSF8
D	246	HIS		expression tag	UNP Q8WSF8
D	247	HIS	-	expression tag	UNP Q8WSF8
D	248	HIS	-	expression tag	UNP Q8WSF8
D	249	HIS	-	expression tag	UNP Q8WSF8
Е	53	PHE	THR	engineered mutation	UNP Q8WSF8
Е	60	VAL	ALA	conflict	UNP Q8WSF8
Е	74	ARG	GLN	engineered mutation	UNP Q8WSF8
Е	110	ALA	TYR	engineered mutation	UNP Q8WSF8
Е	135	SER	ILE	engineered mutation	UNP Q8WSF8
Е	155	VAL	ALA	conflict	UNP Q8WSF8
Е	162	GLU	GLY	engineered mutation	UNP Q8WSF8



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
Е	206	LYS	SER	engineered mutation	UNP Q8WSF8
Е	207	GLY	CYS	engineered mutation	UNP Q8WSF8
E	208	THR	CYS	engineered mutation	UNP Q8WSF8
Е	209	GLY	PRO	engineered mutation	UNP Q8WSF8
Е	237	GLU	-	expression tag	UNP Q8WSF8
E	238	ASN	ı	expression tag	UNP Q8WSF8
Е	239	LEU	-	expression tag	UNP Q8WSF8
E	240	TYR	-	expression tag	UNP Q8WSF8
Е	241	PHE	-	expression tag	UNP Q8WSF8
Е	242	GLN	-	expression tag	UNP Q8WSF8
E	243	GLY	ı	expression tag	UNP Q8WSF8
Е	244	HIS	-	expression tag	UNP Q8WSF8
Е	245	HIS	-	expression tag	UNP Q8WSF8
Е	246	HIS		expression tag	UNP Q8WSF8
Е	247	HIS	-	expression tag	UNP Q8WSF8
Е	248	HIS	_	expression tag	UNP Q8WSF8
Е	249	HIS	-	expression tag	UNP Q8WSF8


• Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	A	tom	ıs		ZeroOcc	AltConf	Trace
2	F	3	Total 42	C 24	N 3	O 15	0	0	0

• Molecule 3 is (5S)-6,6-dimethyl-5-[(6R)-8-oxo-6,8-dihydrofuro[3,4-e][1,3]benzodioxol-6-y l]-5,6,7,8-tetrahydro[1,3]dioxolo[4,5-g]isoquinolin-6-ium (three-letter code: J94) (formula: $C_{21}H_{20}NO_6$) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
3	A	1	Total C N O	0	0	
J	Λ	1	28 21 1 6	U	U	
3	В	1	Total C N O	0	0	
	Ъ	1	28 21 1 6	U	0	
3	C	1	Total C N O	0	0	
		1	28 21 1 6	0		
3	D	1	Total C N O	0	0	
	D	1	28 21 1 6	0	U	
3	E	1	Total C N O	0	0	
'		1	28 21 1 6		U	

• Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Cl 1 1	0	0
4	В	1	Total Cl 1 1	0	0
4	С	1	Total Cl 1 1	0	0
4	D	1	Total Cl 1 1	0	0
4	Е	1	Total Cl 1 1	0	0

 \bullet Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $\rm C_8H_{15}NO_6).$





Mol	Chain	Residues	${f Atoms}$				ZeroOcc	AltConf
5	D	1	Total				0	0
			14					
5	E	1	Total	C	Ν	O	0	0
0	ь	_	14	8	1	5		

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	86	Total O 86 86	0	0
6	В	100	Total O 100 100	0	0
6	С	73	Total O 73 73	0	0
6	D	62	Total O 62 62	0	0
6	Е	83	Total O 83 83	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Soluble acetylcholine receptor



• Molecule 1: Soluble acetylcholine receptor



GLY HIS HIS HIS HIS HIS

 $\bullet \ \, \text{Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2$

01 ' D	
Chain F:	100%

NAG1 NAG2 NAG3



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	71.24Å 132.24Å 132.63Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.82 - 2.40	Depositor
Resolution (A)	41.83 - 2.40	EDS
% Data completeness	99.9 (46.82-2.40)	Depositor
(in resolution range)	$100.0 \ (41.83-2.40)$	EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.99~({ m at}~2.39{ m \AA})$	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
D D.	0.230 , 0.249	Depositor
R, R_{free}	0.234 , 0.255	DCC
R_{free} test set	2447 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	39.9	Xtriage
Anisotropy	0.260	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.29 , 23.5	EDS
L-test for twinning ²	$< L > = 0.54, < L^2> = 0.38$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8813	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.19% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, CL, J94

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Chain		Bond lengths		ond angles
MIOI	Chain	RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/1688	0.75	1/2300~(0.0%)
1	В	0.51	0/1672	0.79	3/2276~(0.1%)
1	С	0.51	0/1689	0.80	5/2299~(0.2%)
1	D	0.47	0/1671	0.75	$2/2275 \ (0.1\%)$
1	E	0.51	0/1675	0.88	$4/2280 \ (0.2\%)$
All	All	0.50	0/8395	0.79	$15/11430 \ (0.1\%)$

There are no bond length outliers.

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^o)$
1	E	224	ARG	NE-CZ-NH2	17.40	129.00	120.30
1	E	224	ARG	NE-CZ-NH1	-10.65	114.97	120.30
1	С	76	ARG	NE-CZ-NH2	7.67	124.14	120.30
1	D	74	ARG	NE-CZ-NH1	-6.91	116.84	120.30
1	С	76	ARG	NE-CZ-NH1	-6.85	116.88	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mo	l Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1649	0	1582	4	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	В	1629	0	1567	5	0
1	С	1646	0	1582	2	0
1	D	1633	0	1565	2	0
1	E	1637	0	1571	1	0
2	F	42	0	37	0	0
3	A	28	0	20	1	0
3	В	28	0	20	0	0
3	С	28	0	20	4	0
3	D	28	0	20	0	0
3	E	28	0	20	0	0
4	A	1	0	0	0	0
4	В	1	0	0	0	0
4	С	1	0	0	0	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
5	D	14	0	13	0	0
5	E	14	0	13	0	0
6	A	86	0	0	1	0
6	В	100	0	0	0	0
6	С	73	0	0	2	0
6	D	62	0	0	0	0
6	Ε	83	0	0	0	0
All	All	8813	0	8030	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 14 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{aligned}$	$egin{aligned} ext{Clash} \ ext{overlap} & (ext{Å}) \end{aligned}$
1:B:69:LEU:HD13	1:B:142:PHE:HE1	1.43	0.83
1:B:69:LEU:HD13	1:B:142:PHE:CE1	2.24	0.73
1:C:162:GLU:OE2	3:C:601:J94:H6	2.07	0.54
1:B:69:LEU:CD1	1:B:142:PHE:HE1	2.19	0.53
3:C:601:J94:O12	1:D:74:ARG:HD2	2.10	0.51

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	$205/249 \ (82\%)$	202 (98%)	3 (2%)	0	100	100
1	В	200/249 (80%)	197 (98%)	3 (2%)	0	100	100
1	С	203/249 (82%)	199 (98%)	4 (2%)	0	100	100
1	D	200/249 (80%)	197 (98%)	3 (2%)	0	100	100
1	Е	201/249 (81%)	198 (98%)	3 (2%)	0	100	100
All	All	1009/1245 (81%)	993 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	$186/222 \; (84\%)$	183 (98%)	3 (2%)	62	79
1	В	$186/222 \; (84\%)$	181 (97%)	5 (3%)	44	65
1	С	187/222 (84%)	187 (100%)	0	100	100
1	D	185/222 (83%)	183 (99%)	2 (1%)	73	87
1	${ m E}$	$185/222 \ (83\%)$	183 (99%)	2 (1%)	73	87
All	All	929/1110 (84%)	917 (99%)	12 (1%)	69	84

5 of 12 residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	В	174	LYS
1	D	74	ARG
1	E	74	ARG
1	D	205	TYR
1	В	33	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	203	GLN
1	В	20	GLN
1	В	128	HIS
1	D	203	GLN
1	E	203	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	F	1	1,2	14,14,15	0.47	0	17,19,21	1.46	3 (17%)
2	NAG	F	2	2	14,14,15	0.72	0	17,19,21	1.47	2 (11%)
2	NAG	F	3	2	14,14,15	0.88	1 (7%)	17,19,21	1.54	4 (23%)



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	F	1	1,2	-	4/6/23/26	0/1/1/1
2	NAG	F	2	2	-	2/6/23/26	0/1/1/1
2	NAG	F	3	2	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(\mathbf{\mathring{A}})$	$\operatorname{Ideal}(ext{\AA})$
2	F	3	NAG	C1-C2	2.63	1.56	1.52

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
2	F	3	NAG	C2-N2-C7	3.25	127.53	122.90
2	F	2	NAG	C2-N2-C7	3.20	127.47	122.90
2	F	1	NAG	O5-C5-C6	3.12	112.10	107.20
2	F	2	NAG	C8-C7-N2	3.08	121.32	116.10
2	F	3	NAG	C1-C2-N2	2.85	115.36	110.49

There are no chirality outliers.

5 of 8 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	F	3	NAG	C1-C2-N2-C7
2	F	1	NAG	O5-C5-C6-O6
2	F	1	NAG	C4-C5-C6-O6
2	F	1	NAG	C8-C7-N2-C2
2	F	1	NAG	O7-C7-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 5 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trans	Chain	Dog	Link	Bond lengths			В	ond ang	les
MIOI	Type	Chain	m Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	J94	E	601	-	33,33,33	0.29	0	48,52,52	0.58	0
3	J94	D	601	_	33,33,33	0.38	0	48,52,52	0.53	0
5	NAG	E	602	1	14,14,15	0.68	0	17,19,21	1.72	5 (29%)
3	J94	В	601	-	33,33,33	0.32	0	48,52,52	0.55	0
3	J94	A	601	-	33,33,33	0.43	0	48,52,52	0.61	0
3	J94	С	601	-	33,33,33	0.57	1 (3%)	48,52,52	0.55	0



Mol	Type	Chain	Res	Link	Bo	ond leng	$ ag{ths}$	В	ond ang	les
MIOI	туре	Chain	res	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	D	602	1	14,14,15	0.55	0	17,19,21	1.42	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	J94	Е	601	-	-	3/4/44/44	0/6/6/6
3	J94	D	601	-	-	0/4/44/44	0/6/6/6
5	NAG	Ε	602	1	-	3/6/23/26	0/1/1/1
3	J94	В	601	=	-	2/4/44/44	0/6/6/6
3	J94	A	601	-	-	2/4/44/44	0/6/6/6
3	J94	С	601	-	-	0/4/44/44	0/6/6/6
5	NAG	D	602	1	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$\mathbf{Ideal}(\mathbf{\mathring{A}})$
3	С	601	J94	C07-C15	-2.74	1.48	1.52

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
5	D	602	NAG	C1-O5-C5	4.31	118.04	112.19
5	E	602	NAG	C2-N2-C7	3.81	128.33	122.90
5	E	602	NAG	O5-C1-C2	3.00	116.02	111.29
5	E	602	NAG	C1-C2-N2	-2.51	106.20	110.49
5	D	602	NAG	O5-C1-C2	2.39	115.07	111.29

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	601	J94	N02-C15-C16-O17
3	Е	601	J94	N02-C15-C16-C28
3	E	601	J94	C07-C15-C16-O17
5	E	602	NAG	C3-C2-N2-C7
5	E	602	NAG	C4-C5-C6-O6




There are no ring outliers.

2 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	601	J94	1	0
3	С	601	J94	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed			$OWAB(A^2)$	Q < 0.9
1	A	207/249 (83%)	0.06	6 (2%) 51 50	24, 38, 77, 114	0
1	В	203/249 (81%)	0.07	7 (3%) 45 44	24, 36, 69, 115	0
1	С	$206/249 \ (82\%)$	0.26	10 (4%) 29 28	25, 39, 82, 123	0
1	D	204/249 (81%)	0.17	9 (4%) 34 33	28, 46, 88, 132	0
1	E	$205/249 \ (82\%)$	0.06	10 (4%) 29 28	26, 40, 78, 135	0
All	All	1025/1245 (82%)	0.12	42 (4%) 37 36	24, 39, 83, 135	0

The worst 5 of 42 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	205	TYR	6.3
1	В	209	GLY	5.8
1	E	33	ARG	5.3
1	С	227	GLY	4.9
1	Е	205	TYR	4.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	\mathbf{Res}	Atoms	RSCC	RSR	${f B\text{-factors}}({f A}^2)$	Q<0.9
2	NAG	F	3	14/15	0.75	0.26	75,81,88,89	0



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B\text{-factors}}({f \AA}^2)$	Q < 0.9
2	NAG	F	2	14/15	0.76	0.26	73,84,86,89	0
2	NAG	F	1	14/15	0.76	0.21	71,76,80,81	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B-factors(\AA^2)}$	Q < 0.9
5	NAG	E	602	14/15	0.52	0.30	107,115,119,119	0
3	J94	E	601	28/28	0.54	0.30	97,119,126,128	0
3	J94	D	601	28/28	0.54	0.36	97,110,126,129	0
3	J94	С	601	28/28	0.55	0.31	88,101,109,112	0
3	J94	A	601	28/28	0.68	0.28	74,88,115,118	0
3	J94	В	601	28/28	0.72	0.26	55,66,78,85	0




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$ extbf{B-factors}(extbf{A}^2)$	Q < 0.9
5	NAG	D	602	14/15	0.73	0.29	94,102,108,113	0
4	CL	D	603	1/1	0.89	0.14	56, 56, 56, 56	0
4	CL	В	605	1/1	0.96	0.10	37,37,37,37	0
4	CL	С	602	1/1	0.97	0.09	49,49,49,49	0
4	CL	Ε	603	1/1	0.97	0.09	58,58,58,58	0
4	CL	A	602	1/1	0.99	0.08	37,37,37,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









Electron density around J94 D 601: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)














6.5 Other polymers (i)

There are no such residues in this entry.

