



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 3, 2023 – 02:15 pm GMT

PDB ID : 1OCW
Title : Free conformation Ab2 of the IgE SPE-7
Authors : James, L.C.; Roversi, P.; Tawfik, D.
Deposited on : 2003-02-11
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	H	121	 12% 51% 27% 8% .
2	L	120	 42% 45% 5% 8%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1835 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IMMUNOGLOBULIN E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	120	954	611	156	182	5	0	0	0

- Molecule 2 is a protein called IMMUNOGLOBULIN E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	110	813	510	138	163	2	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	H	43	Total	O	0	0
			43	43		
3	L	25	Total	O	0	0
			25	25		

4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	79.55Å 79.55Å 67.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.17 – 2.00 28.12 – 2.02	Depositor EDS
% Data completeness (in resolution range)	94.4 (28.17-2.00) 98.1 (28.12-2.02)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.24 (at 2.01Å)	Xtrriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.267 , 0.290 0.410 , 0.409	Depositor DCC
R_{free} test set	695 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	20.1	Xtrriage
Anisotropy	0.084	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 36.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.487 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.64	EDS
Total number of atoms	1835	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	H	0.35	0/982	1.11	10/1331 (0.8%)
2	L	0.28	0/829	0.80	1/1133 (0.1%)
All	All	0.32	0/1811	0.98	11/2464 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	H	0	1

There are no bond length outliers.

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	43	ARG	NE-CZ-NH2	11.60	126.10	120.30
1	H	101	TYR	C-N-CA	10.74	148.54	121.70
1	H	100	TRP	CA-CB-CG	6.97	126.95	113.70
1	H	105	TYR	C-N-CA	6.21	137.24	121.70
1	H	100	TRP	C-N-CA	5.83	136.27	121.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	H	41	PRO	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	954	0	913	205	75
2	L	813	0	794	118	36
3	H	43	0	0	24	0
3	L	25	0	0	9	0
All	All	1835	0	1707	238	75

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 69.

The worst 5 of 238 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:47:TRP:CZ3	2:L:100:PHE:HB2	1.18	1.64
1:H:110:TRP:CD1	2:L:46:PHE:HB2	1.37	1.58
1:H:110:TRP:CG	2:L:46:PHE:HB2	1.38	1.55
1:H:47:TRP:HZ3	2:L:100:PHE:CB	1.29	1.44
1:H:110:TRP:CD2	2:L:46:PHE:HB3	1.56	1.40

The worst 5 of 75 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:112:GLN:NE2	2:L:8:SER:OG[3_655]	0.50	1.70
1:H:112:GLN:OE1	2:L:8:SER:C[3_655]	0.52	1.68
1:H:3:GLN:NE2	2:L:10:LEU:CG[3_655]	0.95	1.25
1:H:112:GLN:OE1	2:L:9:ALA:N[3_655]	0.97	1.23
1:H:3:GLN:NE2	2:L:10:LEU:CD1[3_655]	1.08	1.12

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	118/121 (98%)	88 (75%)	16 (14%)	14 (12%)	0	0
2	L	108/120 (90%)	99 (92%)	9 (8%)	0	100	100
All	All	226/241 (94%)	187 (83%)	25 (11%)	14 (6%)	1	0

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	32	TYR
1	H	33	TRP
1	H	42	GLY
1	H	77	SER
1	H	102	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	101/101 (100%)	60 (59%)	41 (41%)	0	0
2	L	87/97 (90%)	77 (88%)	10 (12%)	5	3
All	All	188/198 (95%)	137 (73%)	51 (27%)	0	0

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	96	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	H	106	TYR
2	L	105	LYS
1	H	98	ARG
1	H	101	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	39	GLN
1	H	112	GLN
2	L	81	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.