



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 22, 2024 – 07:22 PM EDT

PDB ID : 6OCZ
Title : Crystal Structure of Mycobacterium tuberculosis Proteasome in Complex with Phenylimidazole-based Inhibitor A86
Authors : Hsu, H.C.; Li, H.
Deposited on : 2019-03-25
Resolution : 2.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1332 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	240	 2% 5% 90% 10% 9%
1	B	240	 5% 5% 89% 10% 5%
1	C	240	 5% 5% 90% 10% 5%
1	D	240	 2% 5% 89% 10% 2%
1	E	240	 3% 5% 90% 10% 3%
1	F	240	 4% 5% 90% 10% 4%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	240	90% 10%
1	O	240	90% 10%
1	P	240	91% 9%
1	Q	240	89% 10%
1	R	240	88% 10%
1	S	240	89% 9%
1	T	240	90% 10%
1	U	240	90% 10%
2	H	234	94% 5%
2	I	234	95% 5%
2	J	234	94% 5%
2	K	234	95% 5%
2	L	234	95% 5%
2	M	234	95% 5%
2	N	234	95% 5%
2	V	234	95% 5%
2	W	234	95% 5%
2	X	234	94% 5%
2	Y	234	95% 5%
2	Z	234	94% 5%
2	a	234	95% 5%
2	b	234	95% 5%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 48235 atoms, of which 210 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	218	1677	1050	306	317	4	0	0	0
1	B	215	1660	1041	303	312	4	0	0	0
1	C	217	1672	1047	305	316	4	0	0	0
1	D	215	1655	1035	303	313	4	0	0	0
1	E	217	1670	1046	305	315	4	0	0	0
1	F	216	1663	1041	304	314	4	0	0	0
1	G	216	1661	1039	304	314	4	0	0	0
1	O	217	1670	1045	305	316	4	0	0	0
1	P	219	1685	1054	307	320	4	0	0	0
1	Q	216	1668	1045	304	315	4	0	0	0
1	R	215	1657	1038	303	312	4	0	0	0
1	S	218	1678	1050	306	318	4	0	0	0
1	T	217	1671	1047	305	315	4	0	0	0
1	U	216	1664	1043	304	313	4	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	MET	-	initiating methionine	UNP P9WHU1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	9	MET	-	initiating methionine	UNP P9WHU1
C	9	MET	-	initiating methionine	UNP P9WHU1
D	9	MET	-	initiating methionine	UNP P9WHU1
E	9	MET	-	initiating methionine	UNP P9WHU1
F	9	MET	-	initiating methionine	UNP P9WHU1
G	9	MET	-	initiating methionine	UNP P9WHU1
O	9	MET	-	initiating methionine	UNP P9WHU1
P	9	MET	-	initiating methionine	UNP P9WHU1
Q	9	MET	-	initiating methionine	UNP P9WHU1
R	9	MET	-	initiating methionine	UNP P9WHU1
S	9	MET	-	initiating methionine	UNP P9WHU1
T	9	MET	-	initiating methionine	UNP P9WHU1
U	9	MET	-	initiating methionine	UNP P9WHU1

- Molecule 2 is a protein called Proteasome subunit beta.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	222	1638	1027	282	324	5	0	0	0
2	I	222	1638	1027	282	324	5	0	0	0
2	J	222	1638	1027	282	324	5	0	0	0
2	K	223	1642	1029	283	325	5	0	0	0
2	L	223	1642	1029	283	325	5	0	0	0
2	M	222	1638	1027	282	324	5	0	0	0
2	N	223	1642	1029	283	325	5	0	0	0
2	V	223	1642	1029	283	325	5	0	0	0
2	W	223	1642	1029	283	325	5	0	0	0
2	X	222	1638	1027	282	324	5	0	0	0
2	Y	223	1642	1029	283	325	5	0	0	0
2	Z	222	1638	1027	282	324	5	0	0	0
2	a	223	1642	1029	283	325	5	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	b	223	1642	1029	283	325	5	0	0	0

- Molecule 3 is DIMETHYLFORMAMIDE (three-letter code: DMF) (formula: C₃H₇NO).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
3	A	1	Total 12	C 3	H 7	N 1	O 1	0	0
3	B	1	Total 12	C 3	H 7	N 1	O 1	0	0
3	C	1	Total 12	C 3	H 7	N 1	O 1	0	0
3	E	1	Total 12	C 3	H 7	N 1	O 1	0	0
3	F	1	Total 12	C 3	H 7	N 1	O 1	0	0
3	F	1	Total 12	C 3	H 7	N 1	O 1	0	0
3	G	1	Total 12	C 3	H 7	N 1	O 1	0	0
3	G	1	Total 12	C 3	H 7	N 1	O 1	0	0
3	I	1	Total 12	C 3	H 7	N 1	O 1	0	0
3	J	1	Total 12	C 3	H 7	N 1	O 1	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	J	1	Total	C	H	N	O	0	0
			12	3	7	1	1		
3	O	1	Total	C	H	N	O	0	0
			12	3	7	1	1		
3	P	1	Total	C	H	N	O	0	0
			12	3	7	1	1		
3	Q	1	Total	C	H	N	O	0	0
			12	3	7	1	1		
3	R	1	Total	C	H	N	O	0	0
			12	3	7	1	1		
3	R	1	Total	C	H	N	O	0	0
			12	3	7	1	1		
3	S	1	Total	C	H	N	O	0	0
			12	3	7	1	1		
3	S	1	Total	C	H	N	O	0	0
			12	3	7	1	1		
3	T	1	Total	C	H	N	O	0	0
			12	3	7	1	1		
3	U	1	Total	C	H	N	O	0	0
			12	3	7	1	1		

- Molecule 4 is N-{(2S)-1-((2S)-1-[(2,4-difluorobenzyl)amino]-1-oxopropan-2-yl)amino)-1,4-dioxo-4-[(2R)-2-phenylpyrrolidin-1-yl]butan-2-yl}-5-methyl-1,2-oxazole-3-carboxamide (non-preferred name) (three-letter code: M6Y) (formula: C₂₉H₃₁F₂N₅O₅).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	H	1	Total 41	C 29	F 2	N 5	O 5	0	0
4	I	1	Total 41	C 29	F 2	N 5	O 5	0	0
4	J	1	Total 41	C 29	F 2	N 5	O 5	0	0
4	K	1	Total 41	C 29	F 2	N 5	O 5	0	0
4	L	1	Total 41	C 29	F 2	N 5	O 5	0	0
4	M	1	Total 41	C 29	F 2	N 5	O 5	0	0
4	N	1	Total 41	C 29	F 2	N 5	O 5	0	0
4	V	1	Total 41	C 29	F 2	N 5	O 5	0	0
4	W	1	Total 41	C 29	F 2	N 5	O 5	0	0
4	X	1	Total 41	C 29	F 2	N 5	O 5	0	0
4	Y	1	Total 41	C 29	F 2	N 5	O 5	0	0
4	Z	1	Total 41	C 29	F 2	N 5	O 5	0	0
4	a	1	Total 41	C 29	F 2	N 5	O 5	0	0
4	b	1	Total 41	C 29	F 2	N 5	O 5	0	0

- Molecule 5 is CITRIC ACID (three-letter code: CIT) (formula: C₆H₈O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	H			O
5	H	1	Total 18	C 6	H 5	O 7	0	0
5	I	1	Total 18	C 6	H 5	O 7	0	0
5	J	1	Total 18	C 6	H 5	O 7	0	0
5	K	1	Total 18	C 6	H 5	O 7	0	0
5	L	1	Total 18	C 6	H 5	O 7	0	0
5	M	1	Total 18	C 6	H 5	O 7	0	0
5	N	1	Total 18	C 6	H 5	O 7	0	0
5	V	1	Total 18	C 6	H 5	O 7	0	0
5	W	1	Total 18	C 6	H 5	O 7	0	0
5	X	1	Total 18	C 6	H 5	O 7	0	0
5	Y	1	Total 18	C 6	H 5	O 7	0	0
5	Z	1	Total 18	C 6	H 5	O 7	0	0
5	a	1	Total 18	C 6	H 5	O 7	0	0
5	b	1	Total 18	C 6	H 5	O 7	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	27	Total O 27 27	0	0
6	B	20	Total O 20 20	0	0
6	C	29	Total O 29 29	0	0
6	D	20	Total O 20 20	0	0
6	E	24	Total O 24 24	0	0
6	F	16	Total O 16 16	0	0
6	G	21	Total O 21 21	0	0
6	H	36	Total O 36 36	0	0
6	I	49	Total O 49 49	0	0
6	J	36	Total O 36 36	0	0
6	K	41	Total O 41 41	0	0
6	L	42	Total O 42 42	0	0
6	M	35	Total O 35 35	0	0
6	N	34	Total O 34 34	0	0
6	O	17	Total O 17 17	0	0
6	P	26	Total O 26 26	0	0
6	Q	31	Total O 31 31	0	0
6	R	28	Total O 28 28	0	0
6	S	28	Total O 28 28	0	0
6	T	13	Total O 13 13	0	0
6	U	27	Total O 27 27	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	V	42	Total O 42 42	0	0
6	W	39	Total O 39 39	0	0
6	X	35	Total O 35 35	0	0
6	Y	40	Total O 40 40	0	0
6	Z	37	Total O 37 37	0	0
6	a	29	Total O 29 29	0	0
6	b	32	Total O 32 32	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




• Molecule 1: Proteasome subunit alpha




• Molecule 1: Proteasome subunit alpha




• Molecule 1: Proteasome subunit alpha




• Molecule 1: Proteasome subunit alpha




• Molecule 1: Proteasome subunit alpha




• Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 2: Proteasome subunit beta




- Molecule 2: Proteasome subunit beta




- Molecule 2: Proteasome subunit beta



- Molecule 2: Proteasome subunit beta




- Molecule 2: Proteasome subunit beta




- Molecule 2: Proteasome subunit beta

Chain M:  95% 5%




- Molecule 2: Proteasome subunit beta

Chain N:  95% 5%




- Molecule 2: Proteasome subunit beta

Chain V:  95% 5%




- Molecule 2: Proteasome subunit beta

Chain W:  95% 5%




- Molecule 2: Proteasome subunit beta

Chain X:  94% 5%




- Molecule 2: Proteasome subunit beta

Chain Y:  95% 5%




- Molecule 2: Proteasome subunit beta

Chain Z:  94% 5%




- Molecule 2: Proteasome subunit beta

Chain a:  95% 5%



- Molecule 2: Proteasome subunit beta

Chain b:  95% 5%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	120.77Å 198.06Å 167.92Å 90.00° 103.72° 90.00°	Depositor
Resolution (Å)	42.92 – 2.65 42.92 – 2.65	Depositor EDS
% Data completeness (in resolution range)	93.5 (42.92-2.65) 93.5 (42.92-2.65)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.98 (at 2.65Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.184 , 0.235 0.187 , 0.237	Depositor DCC
R_{free} test set	10226 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	35.4	Xtrriage
Anisotropy	0.442	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	48235	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: M6Y, DMF, CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/1701	0.47	0/2297
1	B	0.26	0/1684	0.44	0/2274
1	C	0.26	0/1696	0.45	0/2290
1	D	0.27	0/1679	0.46	0/2266
1	E	0.26	0/1694	0.45	0/2287
1	F	0.33	0/1687	0.47	0/2277
1	G	0.27	0/1684	0.46	0/2272
1	O	0.33	0/1694	0.50	2/2287 (0.1%)
1	P	0.26	0/1709	0.45	0/2308
1	Q	0.27	0/1692	0.46	0/2285
1	R	0.27	0/1681	0.46	0/2269
1	S	0.27	0/1702	0.46	0/2298
1	T	0.26	0/1695	0.45	0/2289
1	U	0.31	0/1688	0.48	0/2279
2	H	0.27	0/1662	0.48	0/2254
2	I	0.27	0/1662	0.47	0/2254
2	J	0.27	0/1662	0.48	0/2254
2	K	0.28	0/1666	0.49	0/2259
2	L	0.27	0/1666	0.48	0/2259
2	M	0.26	0/1662	0.47	0/2254
2	N	0.27	0/1666	0.48	0/2259
2	V	0.31	0/1666	0.48	0/2259
2	W	0.27	0/1666	0.48	0/2259
2	X	0.27	0/1662	0.47	0/2254
2	Y	0.34	0/1666	0.50	0/2259
2	Z	0.27	0/1662	0.48	0/2254
2	a	0.29	0/1666	0.46	0/2259
2	b	0.27	0/1666	0.47	0/2259
All	All	0.28	0/46986	0.47	2/63574 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	O	85	ARG	CG-CD-NE	-5.90	99.41	111.80
1	O	85	ARG	NE-CZ-NH2	-5.78	117.41	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/240 (89%)	210 (98%)	4 (2%)	0	100	100
1	B	211/240 (88%)	201 (95%)	10 (5%)	0	100	100
1	C	213/240 (89%)	203 (95%)	10 (5%)	0	100	100
1	D	211/240 (88%)	203 (96%)	8 (4%)	0	100	100
1	E	213/240 (89%)	207 (97%)	6 (3%)	0	100	100
1	F	212/240 (88%)	205 (97%)	7 (3%)	0	100	100
1	G	211/240 (88%)	206 (98%)	5 (2%)	0	100	100
1	O	213/240 (89%)	207 (97%)	6 (3%)	0	100	100
1	P	215/240 (90%)	209 (97%)	6 (3%)	0	100	100
1	Q	212/240 (88%)	204 (96%)	8 (4%)	0	100	100
1	R	211/240 (88%)	205 (97%)	5 (2%)	1 (0%)	29	43
1	S	214/240 (89%)	206 (96%)	7 (3%)	1 (0%)	29	43
1	T	213/240 (89%)	205 (96%)	8 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	U	212/240 (88%)	203 (96%)	9 (4%)	0	100	100
2	H	220/234 (94%)	215 (98%)	5 (2%)	0	100	100
2	I	220/234 (94%)	210 (96%)	10 (4%)	0	100	100
2	J	220/234 (94%)	213 (97%)	7 (3%)	0	100	100
2	K	221/234 (94%)	216 (98%)	5 (2%)	0	100	100
2	L	221/234 (94%)	215 (97%)	6 (3%)	0	100	100
2	M	220/234 (94%)	216 (98%)	4 (2%)	0	100	100
2	N	221/234 (94%)	212 (96%)	9 (4%)	0	100	100
2	V	221/234 (94%)	218 (99%)	3 (1%)	0	100	100
2	W	221/234 (94%)	217 (98%)	4 (2%)	0	100	100
2	X	220/234 (94%)	217 (99%)	3 (1%)	0	100	100
2	Y	221/234 (94%)	217 (98%)	4 (2%)	0	100	100
2	Z	220/234 (94%)	215 (98%)	5 (2%)	0	100	100
2	a	221/234 (94%)	215 (97%)	6 (3%)	0	100	100
2	b	221/234 (94%)	218 (99%)	3 (1%)	0	100	100
All	All	6063/6636 (91%)	5888 (97%)	173 (3%)	2 (0%)	100	100

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	S	151	PRO
1	R	151	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	167/184 (91%)	165 (99%)	2 (1%)	71	84
1	B	165/184 (90%)	164 (99%)	1 (1%)	86	92
1	C	166/184 (90%)	164 (99%)	2 (1%)	71	84

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	164/184 (89%)	162 (99%)	2 (1%)	71	84
1	E	166/184 (90%)	165 (99%)	1 (1%)	86	92
1	F	165/184 (90%)	165 (100%)	0	100	100
1	G	165/184 (90%)	165 (100%)	0	100	100
1	O	166/184 (90%)	165 (99%)	1 (1%)	86	92
1	P	168/184 (91%)	168 (100%)	0	100	100
1	Q	166/184 (90%)	163 (98%)	3 (2%)	59	75
1	R	164/184 (89%)	161 (98%)	3 (2%)	59	75
1	S	167/184 (91%)	163 (98%)	4 (2%)	49	67
1	T	166/184 (90%)	166 (100%)	0	100	100
1	U	165/184 (90%)	165 (100%)	0	100	100
2	H	165/172 (96%)	162 (98%)	3 (2%)	59	75
2	I	165/172 (96%)	165 (100%)	0	100	100
2	J	165/172 (96%)	163 (99%)	2 (1%)	71	84
2	K	165/172 (96%)	165 (100%)	0	100	100
2	L	165/172 (96%)	165 (100%)	0	100	100
2	M	165/172 (96%)	165 (100%)	0	100	100
2	N	165/172 (96%)	165 (100%)	0	100	100
2	V	165/172 (96%)	164 (99%)	1 (1%)	86	92
2	W	165/172 (96%)	164 (99%)	1 (1%)	86	92
2	X	165/172 (96%)	163 (99%)	2 (1%)	71	84
2	Y	165/172 (96%)	164 (99%)	1 (1%)	86	92
2	Z	165/172 (96%)	163 (99%)	2 (1%)	71	84
2	a	165/172 (96%)	165 (100%)	0	100	100
2	b	165/172 (96%)	164 (99%)	1 (1%)	86	92
All	All	4630/4984 (93%)	4598 (99%)	32 (1%)	84	91

5 of 32 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	Y	82	ARG
2	Z	31	VAL
2	J	219	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	J	208	SER
2	Z	144	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	U	101	ASN
2	W	110	HIS
1	O	98	GLN
1	Q	101	ASN
1	R	98	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

48 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	DMF	C	301	-	4,4,4	0.40	0	4,4,4	0.46	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	M6Y	L	301	-	42,44,44	2.86	12 (28%)	53,61,61	1.55	10 (18%)
3	DMF	S	302	-	4,4,4	0.43	0	4,4,4	0.39	0
4	M6Y	Z	301	-	42,44,44	2.83	10 (23%)	53,61,61	1.53	12 (22%)
5	CIT	a	302	-	12,12,12	1.07	0	17,17,17	1.43	1 (5%)
5	CIT	N	302	-	12,12,12	1.06	0	17,17,17	1.49	2 (11%)
3	DMF	S	301	-	4,4,4	0.42	0	4,4,4	0.46	0
4	M6Y	N	301	-	42,44,44	2.84	11 (26%)	53,61,61	1.52	11 (20%)
4	M6Y	I	301	-	42,44,44	2.86	10 (23%)	53,61,61	1.58	9 (16%)
4	M6Y	X	301	-	42,44,44	2.83	12 (28%)	53,61,61	1.61	11 (20%)
5	CIT	X	302	-	12,12,12	1.08	0	17,17,17	1.42	1 (5%)
5	CIT	V	302	-	12,12,12	1.11	0	17,17,17	1.38	1 (5%)
3	DMF	R	301	-	4,4,4	0.37	0	4,4,4	0.44	0
3	DMF	T	301	-	4,4,4	0.39	0	4,4,4	0.53	0
4	M6Y	V	301	-	42,44,44	2.87	11 (26%)	53,61,61	1.52	9 (16%)
4	M6Y	b	301	-	42,44,44	3.53	25 (59%)	53,61,61	1.86	17 (32%)
3	DMF	U	301	-	4,4,4	0.36	0	4,4,4	0.43	0
3	DMF	J	303	-	4,4,4	0.39	0	4,4,4	0.47	0
3	DMF	O	301	-	4,4,4	0.34	0	4,4,4	0.33	0
4	M6Y	a	301	-	42,44,44	2.84	10 (23%)	53,61,61	1.64	10 (18%)
5	CIT	H	302	-	12,12,12	1.05	0	17,17,17	1.47	2 (11%)
3	DMF	G	301	-	4,4,4	0.36	0	4,4,4	0.35	0
3	DMF	F	302	-	4,4,4	0.39	0	4,4,4	0.40	0
3	DMF	J	304	-	4,4,4	0.46	0	4,4,4	0.47	0
4	M6Y	J	301	-	42,44,44	2.84	11 (26%)	53,61,61	1.66	11 (20%)
3	DMF	F	301	-	4,4,4	0.36	0	4,4,4	0.43	0
5	CIT	M	302	-	12,12,12	1.03	0	17,17,17	1.70	5 (29%)
5	CIT	W	302	-	12,12,12	1.08	0	17,17,17	1.46	2 (11%)
5	CIT	b	302	-	12,12,12	1.07	0	17,17,17	1.57	3 (17%)
4	M6Y	Y	301	-	42,44,44	2.86	11 (26%)	53,61,61	1.57	12 (22%)
4	M6Y	H	301	-	42,44,44	2.85	10 (23%)	53,61,61	1.56	10 (18%)
3	DMF	I	303	-	4,4,4	0.43	0	4,4,4	0.38	0
5	CIT	K	302	-	12,12,12	1.01	0	17,17,17	1.64	3 (17%)
3	DMF	A	301	-	4,4,4	0.38	0	4,4,4	0.40	0
5	CIT	I	302	-	12,12,12	1.11	0	17,17,17	1.45	2 (11%)
4	M6Y	M	301	-	42,44,44	2.86	10 (23%)	53,61,61	1.53	9 (16%)
3	DMF	R	302	-	4,4,4	0.37	0	4,4,4	0.46	0
3	DMF	B	301	-	4,4,4	0.40	0	4,4,4	0.41	0
4	M6Y	W	301	-	42,44,44	2.82	10 (23%)	53,61,61	1.64	13 (24%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	M6Y	K	301	-	42,44,44	2.85	10 (23%)	53,61,61	1.60	11 (20%)
3	DMF	Q	301	-	4,4,4	0.37	0	4,4,4	0.34	0
3	DMF	G	302	-	4,4,4	0.40	0	4,4,4	0.45	0
3	DMF	E	301	-	4,4,4	0.39	0	4,4,4	0.38	0
5	CIT	Y	302	-	12,12,12	1.07	0	17,17,17	1.48	1 (5%)
5	CIT	J	302	-	12,12,12	1.09	0	17,17,17	1.50	2 (11%)
3	DMF	P	301	-	4,4,4	0.37	0	4,4,4	0.33	0
5	CIT	L	302	-	12,12,12	1.04	0	17,17,17	1.59	3 (17%)
5	CIT	Z	302	-	12,12,12	1.02	0	17,17,17	1.54	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DMF	C	301	-	-	2/2/2/2	-
4	M6Y	L	301	-	-	2/34/47/47	0/4/4/4
3	DMF	S	302	-	-	0/2/2/2	-
4	M6Y	Z	301	-	-	4/34/47/47	0/4/4/4
5	CIT	a	302	-	-	7/16/16/16	-
5	CIT	N	302	-	-	9/16/16/16	-
3	DMF	S	301	-	-	0/2/2/2	-
4	M6Y	N	301	-	-	1/34/47/47	0/4/4/4
4	M6Y	I	301	-	-	1/34/47/47	0/4/4/4
4	M6Y	X	301	-	-	1/34/47/47	0/4/4/4
5	CIT	X	302	-	-	1/16/16/16	-
5	CIT	V	302	-	-	1/16/16/16	-
3	DMF	R	301	-	-	2/2/2/2	-
3	DMF	T	301	-	-	2/2/2/2	-
4	M6Y	V	301	-	-	3/34/47/47	0/4/4/4
4	M6Y	b	301	-	-	3/34/47/47	0/4/4/4
3	DMF	U	301	-	-	2/2/2/2	-
3	DMF	J	303	-	-	2/2/2/2	-
3	DMF	O	301	-	-	2/2/2/2	-
4	M6Y	a	301	-	-	1/34/47/47	0/4/4/4
5	CIT	H	302	-	-	4/16/16/16	-
3	DMF	G	301	-	-	0/2/2/2	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DMF	F	302	-	-	0/2/2/2	-
3	DMF	J	304	-	-	0/2/2/2	-
4	M6Y	J	301	-	-	1/34/47/47	0/4/4/4
3	DMF	F	301	-	-	0/2/2/2	-
5	CIT	M	302	-	-	8/16/16/16	-
5	CIT	W	302	-	-	3/16/16/16	-
5	CIT	b	302	-	-	3/16/16/16	-
4	M6Y	Y	301	-	-	2/34/47/47	0/4/4/4
4	M6Y	H	301	-	-	0/34/47/47	0/4/4/4
3	DMF	I	303	-	-	0/2/2/2	-
5	CIT	K	302	-	-	4/16/16/16	-
3	DMF	A	301	-	-	0/2/2/2	-
5	CIT	I	302	-	-	2/16/16/16	-
4	M6Y	M	301	-	-	0/34/47/47	0/4/4/4
3	DMF	R	302	-	-	2/2/2/2	-
3	DMF	B	301	-	-	0/2/2/2	-
4	M6Y	W	301	-	-	3/34/47/47	0/4/4/4
4	M6Y	K	301	-	-	1/34/47/47	0/4/4/4
3	DMF	Q	301	-	-	0/2/2/2	-
3	DMF	G	302	-	-	2/2/2/2	-
3	DMF	E	301	-	-	0/2/2/2	-
5	CIT	Y	302	-	-	5/16/16/16	-
5	CIT	J	302	-	-	1/16/16/16	-
3	DMF	P	301	-	-	0/2/2/2	-
5	CIT	L	302	-	-	9/16/16/16	-
5	CIT	Z	302	-	-	7/16/16/16	-

The worst 5 of 163 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	b	301	M6Y	C11-N10	-10.13	1.28	1.47
4	V	301	M6Y	C14-N10	9.89	1.63	1.47
4	Y	301	M6Y	C14-N10	9.79	1.63	1.47
4	H	301	M6Y	C14-N10	9.78	1.63	1.47
4	K	301	M6Y	C14-N10	9.77	1.63	1.47

The worst 5 of 184 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	J	301	M6Y	C13-C14-N10	6.25	108.89	101.93
4	I	301	M6Y	C13-C14-N10	6.19	108.82	101.93
4	X	301	M6Y	C13-C14-N10	6.13	108.75	101.93
4	L	301	M6Y	C13-C14-N10	5.99	108.60	101.93
4	K	301	M6Y	C13-C14-N10	5.95	108.55	101.93

There are no chirality outliers.



5 of 103 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
4	I	301	M6Y	O23-C22-C24-C25
4	J	301	M6Y	O23-C22-C24-C25
4	K	301	M6Y	O23-C22-C24-C25
4	L	301	M6Y	O23-C22-C24-C25
4	W	301	M6Y	O23-C22-C24-C25



There are no ring outliers.



No monomer is involved in short contacts.



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	218/240 (90%)	-0.35	3 (1%) 75 73	19, 33, 55, 83	0
1	B	215/240 (89%)	-0.05	11 (5%) 28 25	23, 44, 73, 89	0
1	C	217/240 (90%)	-0.06	11 (5%) 28 25	21, 42, 73, 86	0
1	D	215/240 (89%)	-0.07	6 (2%) 53 49	20, 40, 66, 80	0
1	E	217/240 (90%)	-0.06	8 (3%) 41 38	22, 41, 69, 83	0
1	F	216/240 (90%)	-0.04	10 (4%) 32 29	21, 45, 70, 87	0
1	G	216/240 (90%)	-0.32	2 (0%) 84 83	21, 35, 62, 71	0
1	O	217/240 (90%)	-0.07	10 (4%) 32 29	23, 42, 74, 89	0
1	P	219/240 (91%)	-0.24	3 (1%) 75 73	20, 37, 64, 93	0
1	Q	216/240 (90%)	-0.31	2 (0%) 84 83	19, 35, 60, 99	0
1	R	215/240 (89%)	-0.20	3 (1%) 75 73	19, 34, 59, 68	0
1	S	218/240 (90%)	-0.35	5 (2%) 60 56	19, 32, 60, 87	0
1	T	217/240 (90%)	-0.17	5 (2%) 60 56	22, 43, 65, 93	0
1	U	216/240 (90%)	-0.16	9 (4%) 36 33	20, 36, 62, 79	0
2	H	222/234 (94%)	-0.54	0 100 100	18, 26, 44, 64	0
2	I	222/234 (94%)	-0.63	0 100 100	17, 25, 42, 58	0
2	J	222/234 (94%)	-0.58	1 (0%) 91 91	17, 25, 42, 80	0
2	K	223/234 (95%)	-0.66	0 100 100	14, 25, 42, 53	0
2	L	223/234 (95%)	-0.62	1 (0%) 92 93	17, 26, 42, 63	0
2	M	222/234 (94%)	-0.60	0 100 100	15, 27, 47, 69	0
2	N	223/234 (95%)	-0.54	2 (0%) 84 83	18, 29, 51, 68	0
2	V	223/234 (95%)	-0.66	0 100 100	15, 24, 41, 53	0
2	W	223/234 (95%)	-0.70	1 (0%) 92 93	17, 25, 41, 58	0
2	X	222/234 (94%)	-0.57	1 (0%) 91 91	18, 25, 40, 62	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
2	Y	223/234 (95%)	-0.66	0 100 100	16, 24, 41, 68	0
2	Z	222/234 (94%)	-0.61	1 (0%) 91 91	17, 26, 44, 61	0
2	a	223/234 (95%)	-0.60	0 100 100	19, 28, 48, 66	0
2	b	223/234 (95%)	-0.52	1 (0%) 92 93	17, 27, 44, 67	0
All	All	6148/6636 (92%)	-0.39	96 (1%) 72 69	14, 30, 63, 99	0

The worst 5 of 96 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	236	ASP	6.6
2	J	115	GLN	5.5
1	F	191	GLY	4.3
1	T	202	THR	4.3
1	B	235	VAL	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	DMF	S	302	5/5	0.90	0.27	34,47,56,57	0
5	CIT	N	302	13/13	0.90	0.19	31,47,61,62	0
5	CIT	Z	302	13/13	0.90	0.21	32,42,51,67	0
5	CIT	L	302	13/13	0.91	0.17	31,40,52,55	0
5	CIT	a	302	13/13	0.91	0.15	29,35,48,52	0
5	CIT	b	302	13/13	0.91	0.17	31,40,56,58	0
3	DMF	J	303	5/5	0.92	0.23	30,37,47,50	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	DMF	B	301	5/5	0.92	0.25	27,39,47,47	0
3	DMF	F	302	5/5	0.92	0.23	27,34,56,56	0
5	CIT	M	302	13/13	0.92	0.17	30,47,57,63	0
5	CIT	H	302	13/13	0.93	0.15	26,39,50,56	0
5	CIT	V	302	13/13	0.93	0.16	28,35,54,57	0
5	CIT	K	302	13/13	0.93	0.13	27,36,42,46	0
3	DMF	Q	301	5/5	0.93	0.21	21,33,40,46	0
3	DMF	F	301	5/5	0.93	0.21	25,34,44,46	0
5	CIT	W	302	13/13	0.94	0.16	26,38,47,52	0
5	CIT	J	302	13/13	0.94	0.17	31,39,47,47	0
3	DMF	I	303	5/5	0.94	0.24	24,35,42,44	0
3	DMF	S	301	5/5	0.94	0.20	22,28,35,36	0
3	DMF	J	304	5/5	0.95	0.26	26,41,50,54	0
5	CIT	I	302	13/13	0.95	0.13	33,38,49,52	0
3	DMF	G	302	5/5	0.95	0.20	34,41,51,51	0
5	CIT	X	302	13/13	0.95	0.21	27,36,44,55	0
5	CIT	Y	302	13/13	0.95	0.18	29,38,45,53	0
3	DMF	A	301	5/5	0.95	0.22	35,42,47,47	0
3	DMF	E	301	5/5	0.95	0.15	26,34,45,45	0
3	DMF	U	301	5/5	0.95	0.22	33,39,50,50	0
4	M6Y	H	301	41/41	0.96	0.14	17,24,38,47	0
4	M6Y	I	301	41/41	0.96	0.15	17,27,38,46	0
4	M6Y	M	301	41/41	0.96	0.14	20,26,36,48	0
4	M6Y	N	301	41/41	0.96	0.14	19,26,37,44	0
4	M6Y	V	301	41/41	0.96	0.15	16,25,33,39	0
4	M6Y	Y	301	41/41	0.96	0.14	18,24,35,47	0
4	M6Y	Z	301	41/41	0.96	0.14	14,24,33,43	0
4	M6Y	a	301	41/41	0.96	0.14	17,24,35,39	0
3	DMF	O	301	5/5	0.96	0.24	27,33,39,42	0
3	DMF	G	301	5/5	0.96	0.15	22,30,37,37	0
3	DMF	R	302	5/5	0.96	0.18	36,43,52,52	0
3	DMF	C	301	5/5	0.97	0.16	21,37,41,48	0
4	M6Y	b	301	41/41	0.97	0.12	20,27,37,47	0
3	DMF	R	301	5/5	0.97	0.19	26,31,38,38	0
4	M6Y	J	301	41/41	0.97	0.14	17,23,31,31	0
4	M6Y	W	301	41/41	0.97	0.14	19,25,34,42	0
4	M6Y	X	301	41/41	0.97	0.15	20,26,35,47	0
4	M6Y	K	301	41/41	0.97	0.13	17,24,36,41	0
4	M6Y	L	301	41/41	0.97	0.15	15,25,34,42	0
3	DMF	T	301	5/5	0.98	0.23	27,37,44,45	0
3	DMF	P	301	5/5	0.99	0.13	30,36,41,41	0




The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around M6Y I 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around M6Y M 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around M6Y N 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around M6Y V 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around M6Y Y 301:**




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)








Electron density around M6Y a 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around M6Y b 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around M6Y J 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around M6Y W 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around M6Y X 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around M6Y K 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.