



wwPDB X-ray Structure Validation Summary Report ⓘ

May 21, 2020 – 12:55 am BST

PDB ID : 1OGK
Title : The crystal structure of Trypanosoma cruzi dUTPase in complex with dUDP
Authors : Harkiolaki, M.; Dodson, E.J.; Bernier-Villamor, V.; Turkenburg, J.P.;
Gonzalez-Pacanowska, D.; Wilson, K.S.
Deposited on : 2003-05-07
Resolution : 2.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3168 (2.90-2.82)
Clashscore	141614	3438 (2.90-2.82)
Ramachandran outliers	138981	3348 (2.90-2.82)
Sidechain outliers	138945	3351 (2.90-2.82)
RSRZ outliers	127900	3103 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	283	 % 51% 24% 6% 19%
1	B	283	 49% 28% 7% • 15%
1	D	283	 6% 45% 31% 8% • 16%
1	E	283	 2% 48% 30% 5% • 14%

2 Entry composition [i](#)


There are 2 unique types of molecules in this entry. The entry contains 7784 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DEOXYURIDINE TRIPHOSPHATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	230	Total 1865	C 1200	N 311	O 345	S 9	0	0	1
1	B	240	Total 1949	C 1256	N 327	O 357	S 9	0	0	1
1	D	238	Total 1934	C 1247	N 325	O 353	S 9	0	0	1
1	E	242	Total 1964	C 1267	N 329	O 359	S 9	0	0	1

- Molecule 2 is DEOXYURIDINE-5'-DIPHOSPHATE (three-letter code: DUD) (formula: $C_9H_{14}N_2O_{11}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	B	1	Total 24	C 9	N 2	O 11	P 2	0	0
2	D	1	Total 24	C 9	N 2	O 11	P 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	E	1	24	9	2	11	2	0	0



• Molecule 1: DEOXYURIDINE TRIPHOSPHATASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	106.39Å 50.31Å 111.39Å 90.00° 101.35° 90.00°	Depositor
Resolution (Å)	111.80 – 2.85 20.03 – 2.85	Depositor EDS
% Data completeness (in resolution range)	99.1 (111.80-2.85) 99.5 (20.03-2.85)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 2.83Å)	Xtriage
Refinement program	REFMAC 5.1.13	Depositor
R, R_{free}	0.205 , 0.273 0.209 , 0.267	Depositor DCC
R_{free} test set	1362 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	59.5	Xtriage
Anisotropy	0.017	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 36.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.020 for l,-k,h	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7784	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DUD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.98	0/1906	1.07	7/2584 (0.3%)
1	B	0.98	2/1993 (0.1%)	1.10	15/2698 (0.6%)
1	D	0.75	0/1978	0.96	6/2677 (0.2%)
1	E	0.90	1/2008 (0.0%)	1.07	12/2720 (0.4%)
All	All	0.91	3/7885 (0.0%)	1.05	40/10679 (0.4%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	77	GLU	CG-CD	6.28	1.61	1.51
1	E	223	ASP	CB-CG	-5.87	1.39	1.51
1	B	146	MET	CG-SD	-5.19	1.67	1.81

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	238	ASP	CB-CG-OD2	11.55	128.69	118.30
1	A	139	ASP	CB-CG-OD2	9.23	126.61	118.30
1	B	77	GLU	OE1-CD-OE2	-8.33	113.30	123.30
1	E	52	GLU	OE1-CD-OE2	-7.88	113.85	123.30
1	A	140	ASP	CB-CG-OD2	7.60	125.14	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1865	0	1826	54	0
1	B	1949	0	1922	60	0
1	D	1934	0	1909	71	0
1	E	1964	0	1941	75	0
2	B	24	0	11	0	0
2	D	24	0	11	3	0
2	E	24	0	11	2	0
All	All	7784	0	7631	245	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 245 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:65:VAL:CG1	1:E:66:LYS:H	1.63	1.11
1:E:65:VAL:HG12	1:E:66:LYS:H	1.06	1.10
1:E:10:VAL:CG2	1:E:180:LEU:HD22	1.81	1.08
1:E:10:VAL:HG21	1:E:180:LEU:CD2	1.95	0.95
1:E:65:VAL:HG12	1:E:66:LYS:N	1.86	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	222/283 (78%)	200 (90%)	18 (8%)	4 (2%)	8 25
1	B	234/283 (83%)	211 (90%)	20 (8%)	3 (1%)	12 33
1	D	232/283 (82%)	197 (85%)	29 (12%)	6 (3%)	5 17

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	236/283 (83%)	209 (89%)	19 (8%)	8 (3%)	3	12
All	All	924/1132 (82%)	817 (88%)	86 (9%)	21 (2%)	6	20

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	92	LYS
1	B	223	ASP
1	E	65	VAL
1	E	92	LYS
1	E	153	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/238 (84%)	167 (84%)	33 (16%)	2	5
1	B	208/238 (87%)	167 (80%)	41 (20%)	1	3
1	D	206/238 (87%)	164 (80%)	42 (20%)	1	3
1	E	210/238 (88%)	166 (79%)	44 (21%)	1	2
All	All	824/952 (87%)	664 (81%)	160 (19%)	1	3

5 of 160 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	275	LYS
1	D	106	LYS
1	E	223	ASP
1	D	15	LEU
1	D	37	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	201	ASN
1	D	26	ASN
1	E	20	HIS
1	B	165	GLN
1	E	165	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	DUD	D	1280	-	20,25,25	0.99	1 (5%)	25,38,38	1.13	1 (4%)
2	DUD	B	1280	-	20,25,25	1.40	4 (20%)	25,38,38	1.41	4 (16%)
2	DUD	E	1281	-	20,25,25	1.21	2 (10%)	25,38,38	1.32	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DUD	D	1280	-	-	1/13/28/28	0/2/2/2
2	DUD	B	1280	-	-	1/13/28/28	0/2/2/2
2	DUD	E	1281	-	-	1/13/28/28	0/2/2/2

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	1281	DUD	PA-O5'	-2.80	1.48	1.59
2	B	1280	DUD	PA-O5'	-2.63	1.48	1.59
2	B	1280	DUD	O4'-C4'	-2.60	1.39	1.45
2	B	1280	DUD	C2-N3	-2.58	1.33	1.38
2	D	1280	DUD	C4-N3	2.53	1.37	1.33

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1280	DUD	O2B-PB-O3A	3.73	117.13	104.64
2	D	1280	DUD	O2B-PB-O3A	3.59	116.68	104.64
2	E	1281	DUD	O2B-PB-O3A	3.26	115.57	104.64
2	E	1281	DUD	O4'-C1'-C2'	-3.23	100.15	106.25
2	E	1281	DUD	C4'-O4'-C1'	2.61	115.77	109.45

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	1280	DUD	PB-O3A-PA-O1A
2	B	1280	DUD	PB-O3A-PA-O1A
2	E	1281	DUD	PB-O3A-PA-O1A


There are no ring outliers.


2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1280	DUD	3	0
2	E	1281	DUD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	230/283 (81%)	-0.35	2 (0%) 84 84	22, 34, 50, 78	0
1	B	240/283 (84%)	-0.45	1 (0%) 92 92	24, 34, 45, 61	0
1	D	238/283 (84%)	0.18	16 (6%) 17 13	23, 36, 47, 84	0
1	E	242/283 (85%)	-0.28	6 (2%) 57 54	27, 34, 45, 64	0
All	All	950/1132 (83%)	-0.22	25 (2%) 56 52	22, 35, 48, 84	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	112	GLY	6.2
1	A	101	ASP	4.7
1	A	139	ASP	4.1
1	E	102	ASP	3.7
1	D	113	PHE	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	DUD	D	1280	24/24	0.96	0.12	56,61,63,64	0
2	DUD	E	1281	24/24	0.96	0.14	55,61,63,63	0
2	DUD	B	1280	24/24	0.97	0.17	55,61,63,63	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around DUD E 1281:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around DUD B 1280:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers

There are no such residues in this entry.