



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 8, 2024 – 02:27 am GMT

PDB ID : 5OHT
Title : A GH31 family sulfoquinovosidase from *E. coli* in complex with aza-sugar inhibitor IFGSQ
Authors : Jin, Y.; Williams, S.J.; Goddard-Borger, E.; Davies, G.J.
Deposited on : 2017-07-18
Resolution : 1.87 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	9470 (1.90-1.86)
Clashscore	141614	10282 (1.90-1.86)
Ramachandran outliers	138981	10152 (1.90-1.86)
Sidechain outliers	138945	10152 (1.90-1.86)
RSRZ outliers	127900	9303 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	686	 2% 94%
1	B	686	 3% 94%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11339 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sulfoquinovosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	670	5368	3439	897	1004	28	0	0	0
1	B	666	5316	3409	884	995	28	0	2	0


There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	679	LEU	-	expression tag	UNP P32138
A	680	GLU	-	expression tag	UNP P32138
A	681	HIS	-	expression tag	UNP P32138
A	682	HIS	-	expression tag	UNP P32138
A	683	HIS	-	expression tag	UNP P32138
A	684	HIS	-	expression tag	UNP P32138
A	685	HIS	-	expression tag	UNP P32138
A	686	HIS	-	expression tag	UNP P32138
B	679	LEU	-	expression tag	UNP P32138
B	680	GLU	-	expression tag	UNP P32138
B	681	HIS	-	expression tag	UNP P32138
B	682	HIS	-	expression tag	UNP P32138
B	683	HIS	-	expression tag	UNP P32138
B	684	HIS	-	expression tag	UNP P32138
B	685	HIS	-	expression tag	UNP P32138
B	686	HIS	-	expression tag	UNP P32138

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		
2	B	1	Total	Ca	0	0
			1	1		

- Molecule 3 is [(3 {S},4 {R},5 {R})-4,5-bis(oxidanyl)piperidin-3-yl]methanesulfonic acid (three-letter code: 9VH) (formula: C₆H₁₃NO₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	A	1	13	6	1	5	1	0	0
3	B	1	13	6	1	5	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	375	378	378	0	3
4	B	243	249	249	0	6


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Sulfoquinovosidase



- Molecule 1: Sulfoquinovosidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	68.47Å 86.27Å 86.81Å 100.76° 113.77° 97.14°	Depositor
Resolution (Å)	44.62 – 1.87 44.62 – 1.87	Depositor EDS
% Data completeness (in resolution range)	97.8 (44.62-1.87) 97.8 (44.62-1.87)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 1.87Å)	Xtrriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.166 , 0.202 0.176 , 0.211	Depositor DCC
R_{free} test set	6991 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	33.8	Xtrriage
Anisotropy	0.451	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 49.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	11339	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: 9VH, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.96	2/5526 (0.0%)	0.89	9/7512 (0.1%)
1	B	0.90	1/5479 (0.0%)	0.88	9/7456 (0.1%)
All	All	0.93	3/11005 (0.0%)	0.89	18/14968 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	611	ARG	CZ-NH1	5.94	1.40	1.33
1	A	642	GLU	CD-OE2	5.61	1.31	1.25
1	B	611	ARG	CZ-NH1	5.44	1.40	1.33

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	481	ASP	CB-CG-OD1	11.30	128.47	118.30
1	A	611	ARG	NE-CZ-NH2	-11.26	114.67	120.30
1	B	611	ARG	NE-CZ-NH2	-10.67	114.97	120.30
1	A	611	ARG	NE-CZ-NH1	8.35	124.48	120.30
1	B	611	ARG	NE-CZ-NH1	8.15	124.38	120.30
1	A	481	ASP	CB-CG-OD1	7.65	125.19	118.30
1	B	176	ASP	CB-CG-OD1	7.26	124.84	118.30
1	A	663	ARG	NE-CZ-NH1	6.91	123.75	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	276	ARG	NE-CZ-NH1	-6.46	117.07	120.30
1	A	176	ASP	CB-CG-OD1	6.19	123.87	118.30
1	B	663	ARG	NE-CZ-NH1	5.88	123.24	120.30
1	B	198	ASP	CB-CG-OD2	-5.88	113.01	118.30
1	B	276	ARG	NE-CZ-NH2	5.60	123.10	120.30
1	A	623	ARG	NE-CZ-NH1	5.37	122.98	120.30
1	A	527	PHE	CB-CG-CD1	5.29	124.50	120.80
1	A	198	ASP	CB-CG-OD2	-5.26	113.57	118.30
1	A	574	GLU	OE1-CD-OE2	-5.05	117.23	123.30
1	B	623	ARG	NE-CZ-NH1	5.01	122.81	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	17	ASN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5368	0	5067	4	0
1	B	5316	0	4983	4	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	13	0	0	0	0
3	B	13	0	0	0	0
4	A	378	0	0	0	0
4	B	249	0	0	0	0
All	All	11339	0	10050	7	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (7) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:14:HIS:O	1:B:15:GLN:HG3	1.99	0.61
1:A:65:ILE:HG12	1:B:65:ILE:HG12	1.92	0.52
1:A:102:ARG:HG2	1:A:227:GLU:CG	2.45	0.47
1:A:170:LYS:HE2	1:A:171:GLU:N	2.30	0.46
1:B:644:PHE:CE2	1:B:649:VAL:HG21	2.51	0.45
1:B:469:TRP:HA	1:B:502:HIS:O	2.17	0.45
1:A:469:TRP:HA	1:A:502:HIS:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	666/686 (97%)	648 (97%)	18 (3%)	0	100	100
1	B	662/686 (96%)	645 (97%)	17 (3%)	0	100	100
All	All	1328/1372 (97%)	1293 (97%)	35 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	558/580 (96%)	548 (98%)	10 (2%)	59	52
1	B	549/580 (95%)	543 (99%)	6 (1%)	73	70

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1107/1160 (95%)	1091 (99%)	16 (1%)	67 62

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	LEU
1	A	10	ASP
1	A	25	GLN
1	A	80	LEU
1	A	170	LYS
1	A	182	PHE
1	A	469	TRP
1	A	488	PRO
1	A	628	LEU
1	A	648	GLU
1	B	9	LEU
1	B	10	ASP
1	B	182	PHE
1	B	469	TRP
1	B	628	LEU
1	B	676	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	288	GLN
1	A	351	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	9VH	A	702	-	11,13,13	1.00	0	14,19,19	1.74	3 (21%)
3	9VH	B	702	-	11,13,13	1.05	1 (9%)	14,19,19	1.97	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	9VH	A	702	-	-	0/5/19/19	1/1/1/1
3	9VH	B	702	-	-	1/5/19/19	1/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	702	9VH	O3-S1	2.65	1.52	1.45

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	702	9VH	O1-S1-O2	5.12	123.78	111.27
3	B	702	9VH	O1-S1-O3	-4.16	101.11	111.27
3	A	702	9VH	O1-S1-O2	3.99	121.01	111.27
3	A	702	9VH	O3-S1-O2	-2.72	104.53	113.95
3	A	702	9VH	O1-S1-C1	2.58	109.85	105.74

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	702	9VH	C2-C1-S1-O2

All (2) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	702	9VH	C2-C3-C4-C5-C6-N1
3	B	702	9VH	C2-C3-C4-C5-C6-N1

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	670/686 (97%)	-0.37	16 (2%) 59 60	24, 35, 61, 104	0
1	B	666/686 (97%)	-0.10	23 (3%) 44 45	31, 45, 69, 99	0
All	All	1336/1372 (97%)	-0.24	39 (2%) 51 53	24, 39, 67, 104	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	679	LEU	4.9
1	A	87	SER	4.6
1	B	359	SER	4.6
1	B	358	ALA	4.3
1	B	350	GLN	4.0
1	B	17	ASN	4.0
1	A	76	PRO	3.4
1	A	13	PHE	3.2
1	B	316	LEU	2.9
1	B	678	ILE	2.9
1	A	86	GLY	2.9
1	B	343	ASP	2.8
1	B	371	TYR	2.8
1	B	87	SER	2.7
1	B	315	GLN	2.6
1	B	11	PHE	2.5
1	A	17	ASN	2.5
1	B	360	GLY	2.5
1	A	35	ASP	2.5
1	A	18	ASP	2.4
1	B	644	PHE	2.4
1	A	469	TRP	2.4
1	B	18	ASP	2.3
1	A	78	GLY	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	340	SER	2.3
1	B	169	CYS	2.3
1	B	552	ALA	2.3
1	B	364	LEU	2.3
1	A	359	SER	2.3
1	B	677	SER	2.3
1	A	470	ALA	2.3
1	A	148	LEU	2.3
1	A	9	LEU	2.2
1	A	79	TRP	2.1
1	B	415	TYR	2.1
1	A	99	ASP	2.1
1	B	651	VAL	2.1
1	B	470	ALA	2.0
1	B	171	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	9VH	B	702	13/13	0.96	0.20	36,39,42,43	13
2	CA	B	701	1/1	0.98	0.09	34,34,34,34	0
3	9VH	A	702	13/13	0.99	0.15	26,28,33,35	0
2	CA	A	701	1/1	1.00	0.09	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.