

wwPDB EM Validation Summary Report (i)

Nov 4, 2024 - 01:10 am GMT

PDB ID	:	8OH2
EMDB ID	:	EMD-16876
Title	:	Structure of the Tau-PAM4 Type 1 amyloid fibril
Authors	:	Wilkinson, M.; Louros, N.; Tsaka, G.; Ramakers, M.; Morelli, C.; Garcia,
		T.; Gallardo, R.U.; D'Haeyer, S.; Goossens, V.; Audenaert, D.; Thal, D.R.;
		Ranson, N.A.; Radford, S.E.; Rousseau, F.; Schymkowitz, J.
Deposited on	:	2023-03-20
Resolution	:	2.60 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev113
Mogul	:	1.8.4, CSD as 541 be (2020)
MolProbity	:	4.02b-467
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 2.60 Å.

Ramachandran outliers

Sidechain outliers

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.

Metric	Percentile Ranks	Value
Ramachandran outliers		0
Sidechain outliers		0
Worse		Better
Percentile relat	ive to all structures	
Percentile relat	ive to all EM structures	
Metric	Whole archive	EM structures
wietric	(# Entries)	$(\# { m Entries})$

207382

206894

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for $\geq=3, 2, 1$ and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq=5\%$ The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

16835

16415

Mol	Chain	Length	Quality of chain
1	А	15	100%
1	В	15	100%
1	С	15	100%
1	D	15	100%
1	Е	15	100%
1	F	15	27%
1	G	15	100%
1	Н	15	100%
1	Ι	15	



Mol	Chain	Length	Quality of chain
1	J	15	100%
1	K	15	100%
1	L	15	73%
1	M	15	100%
1	N	15	7%
			100%
1	О	15	100%
1	Р	15	100%
1	Q	15	100%
1	R	15	100%
1	S	15	100%
1	Т	15	13%
1	U	15	100%
1	V	15	100%
1	W	15	100%
1	Х	15	73%
1	Y	15	100%
	Z		7%
1		15	100%
1	a	15	100%
1	b	15	100%
1	с	15	100%
1	d	15	100%
1	е	15	100%
1	f	15	13%
1	g	15	100%
1	h	15	100%
L	1		Continued on nert nage

Continued from previous page...



Continued from previous page...

Mol	Chain	Length	Quality of chain
1	i	15	100%
1	j	15	67%



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 3702 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	_	Ator	ns		AltConf	Trace
1	٨	15	Total	С	Ν	0	0	1
1	А	15	102	62	19	21	0	1
1	р	15	Total	С	Ν	0	0	1
1	В	15	102	62	19	21	0	1
1	C	15	Total	С	Ν	0	0	1
1	С	15	102	62	19	21	0	1
1	D	15	Total	С	Ν	0	0	1
	D	10	102	62	19	21	0	1
1	Е	15	Total	С	Ν	0	0	1
	E	10	102	62	19	21	0	1
1	F	15	Total	С	Ν	0	0	1
	Г	10	102	62	19	21	0	
1	G	15	Total	С	Ν	0	0	1
	G	10	102	62	19	21	0	
1	Н	15	Total	С	Ν	0	0	1
	п	10	102	62	19	21	0	
1	Ι	15	Total	С	Ν	0	0	1
	1	10	102	62	19	21	U	1
1	J	15	Total	С	Ν	0	0	1
1	J	10	102	62	19	21	0	
1	K	15	Total	С	Ν	0	0	1
	Λ	10	102	62	19	21	0	1
1	L	15	Total	С	Ν	0	0	1
		10	102	62	19	21	0	1
1	М	15	Total	С	Ν	0	0	1
	111	10	102	62	19	21	0	1
1	Ν	15	Total	С	Ν	0	0	1
	1 N	10	102	62	19	21		1
1	О	15	Total	С	Ν	0	0	1
	U	10	102	62	19	21		1
1	Р	15	Total	С	Ν	0	0	1
	1	10	102	62	19	21		1
1	Q	15	Total	С	Ν	0	0	1
	v ع	10	102	62	19	21		1

• Molecule 1 is a protein called Microtubule-associated protein tau.



Mol	Chain	<i>i previous pa</i> Residues	-	Ator	ns		AltConf	Trace
			Total	C	N	0		
1	R	15	102	62	19	21	0	1
	a	1 5	Total	С	Ν	0	0	1
1	S	15	102	62	19	21	0	1
1	т	15	Total	С	Ν	0	0	1
1	Т	15	102	62	19	21	0	1
1	U	15	Total	С	Ν	0	0	1
	U	10	102	62	19	21	0	L
1	V	15	Total	С	Ν	Ο	0	1
	v	10	102	62	19	21	0	L
1	W	15	Total	С	Ν	Ο	0	1
	**	10	102	62	19	21	0	1
1	Х	15	Total	С	Ν	Ο	0	1
		10	102	62	19	21	0	1
1	Y	15	Total	С	Ν	Ο	0	1
	-	10	102	62	19	21		-
1	Z	15	Total	С	Ν	Ο	0	1
			102	62	19	21	Ŭ	-
1	a	15	Total	С	Ν	Ο	0	1
			102	62	19	21		
1	b	15	Total	С	Ν	0	0	1
	-	_	102	62	19	21		
1	с	15	Total	С	N	0	0	1
			102	62	19 	21		
1	d	15	Total	C	N 10	0	0	1
			102	$\frac{62}{C}$	19 N	21		
1	е	15	Total		N 10	0	0	1
			102	$\frac{62}{C}$	19 N	$\frac{21}{0}$		
1	f	15	Total 102	62	19	0 21	0	1
			Total	$\frac{02}{C}$	<u>19</u> N	$\frac{21}{0}$		
1	g	15	10tai 102	62	19	0 21	0	1
			Total	$\frac{02}{C}$	19 N	$\frac{21}{0}$		
1	h	15	10tar 102	62	19	$\frac{0}{21}$	0	1
			Total	02 C	N	$\frac{21}{0}$		
1	i	15	100	62	19	$\frac{0}{21}$	0	1
<u> </u>			Total	02 C	N	0		
1	j	15	102	62	19	21	0	1
L				~ -	_0			

Continued from previous page...

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	349	ACE	ARG	conflict	UNP P10636
	•			α 1 1	1



Continue	Continued from previous page						
Chain	Residue	Modelled	Actual	Comment	Reference		
A	363	NH2	VAL	conflict	UNP P10636		
В	349	ACE	ARG	conflict	UNP P10636		
В	363	NH2	VAL	conflict	UNP P10636		
С	349	ACE	ARG	conflict	UNP P10636		
С	363	NH2	VAL	conflict	UNP P10636		
D	349	ACE	ARG	conflict	UNP P10636		
D	363	NH2	VAL	conflict	UNP P10636		
E	349	ACE	ARG	conflict	UNP P10636		
E	363	NH2	VAL	conflict	UNP P10636		
F	349	ACE	ARG	conflict	UNP P10636		
F	363	NH2	VAL	conflict	UNP P10636		
G	349	ACE	ARG	conflict	UNP P10636		
G	363	NH2	VAL	conflict	UNP P10636		
Н	349	ACE	ARG	conflict	UNP P10636		
Н	363	NH2	VAL	conflict	UNP P10636		
Ι	349	ACE	ARG	conflict	UNP P10636		
Ι	363	NH2	VAL	conflict	UNP P10636		
J	349	ACE	ARG	conflict	UNP P10636		
J	363	NH2	VAL	conflict	UNP P10636		
K	349	ACE	ARG	conflict	UNP P10636		
K	363	NH2	VAL	conflict	UNP P10636		
L	349	ACE	ARG	conflict	UNP P10636		
L	363	NH2	VAL	conflict	UNP P10636		
М	349	ACE	ARG	conflict	UNP P10636		
М	363	NH2	VAL	conflict	UNP P10636		
N	349	ACE	ARG	conflict	UNP P10636		
N	363	NH2	VAL	conflict	UNP P10636		
0	349	ACE	ARG	conflict	UNP P10636		
0	363	NH2	VAL	conflict	UNP P10636		
Р	349	ACE	ARG	conflict	UNP P10636		
Р	363	NH2	VAL	conflict	UNP P10636		
Q	349	ACE	ARG	conflict	UNP P10636		
Q	363	NH2	VAL	conflict	UNP P10636		
R	349	ACE	ARG	conflict	UNP P10636		
R	363	NH2	VAL	conflict	UNP P10636		
S	349	ACE	ARG	conflict	UNP P10636		
S	363	NH2	VAL	conflict	UNP P10636		
Т	349	ACE	ARG	conflict	UNP P10636		
Т	363	NH2	VAL	conflict	UNP P10636		
U	349	ACE	ARG	conflict	UNP P10636		
U	363	NH2	VAL	conflict	UNP P10636		
V	349	ACE	ARG	conflict	UNP P10636		

Continued from previous page...



Chain	Residue	Modelled	Actual	Comment	Reference
V	363	NH2	VAL	conflict	UNP P10636
W	349	ACE	ARG	conflict	UNP P10636
W	363	NH2	VAL	conflict	UNP P10636
Х	349	ACE	ARG	conflict	UNP P10636
Х	363	NH2	VAL	conflict	UNP P10636
Y	349	ACE	ARG	conflict	UNP P10636
Y	363	NH2	VAL	conflict	UNP P10636
Ζ	349	ACE	ARG	conflict	UNP P10636
Ζ	363	NH2	VAL	conflict	UNP P10636
a	349	ACE	ARG	conflict	UNP P10636
a	363	NH2	VAL	conflict	UNP P10636
b	349	ACE	ARG	conflict	UNP P10636
b	363	NH2	VAL	conflict	UNP P10636
с	349	ACE	ARG	conflict	UNP P10636
с	363	NH2	VAL	conflict	UNP P10636
d	349	ACE	ARG	conflict	UNP P10636
d	363	NH2	VAL	conflict	UNP P10636
е	349	ACE	ARG	conflict	UNP P10636
е	363	NH2	VAL	conflict	UNP P10636
f	349	ACE	ARG	conflict	UNP P10636
f	363	NH2	VAL	conflict	UNP P10636
g	349	ACE	ARG	conflict	UNP P10636
g	363	NH2	VAL	conflict	UNP P10636
h	349	ACE	ARG	conflict	UNP P10636
h	363	NH2	VAL	conflict	UNP P10636
i	349	ACE	ARG	conflict	UNP P10636
i	363	NH2	VAL	conflict	UNP P10636
j	349	ACE	ARG	conflict	UNP P10636
j	363	NH2	VAL	conflict	UNP P10636

Continued from previous page...

• Molecule 2 is water.

Mol	Chain	Residues	Atoms	AltConf
2	А	1	Total O 1 1	0
2	В	1	Total O 1 1	0
2	С	1	Total O 1 1	0
2	D	1	Total O 1 1	0
2	Е	1	Total O 1 1	0



Continued from previous page...

	*	Residues	Atoms	AltConf
2	G	1	Total O 1 1	0
2	Н	1	Total O 1 1	0
2	Ι	1	Total O 1 1	0
2	J	1	Total O 1 1	0
2	К	1	Total O 1 1	0
2	М	1	Total O 1 1	0
2	Ν	1	Total O 1 1	0
2	О	1	Total O 1 1	0
2	Р	1	Total O 1 1	0
2	Q	1	Total O 1 1	0
2	S	1	Total O 1 1	0
2	Т	1	Total O 1 1	0
2	U	1	Total O 1 1	0
2	V	1	Total O 1 1	0
2	W	1	Total O 1 1	0
2	Y	1	Total O 1 1	0
2	Z	1	Total O 1 1	0
2	a	1	Total O 1 1	0
2	b	1	Total O 1 1	0
2	с	1	Total O 1 1	0
2	е	1	Total O 1 1	0



Continued from previous page...

Mol	Chain	Residues	Atoms	AltConf
2	f	1	Total O 1 1	0
2	g	1	Total O 1 1	0
2	h	1	Total O 1 1	0
2	i	1	Total O 1 1	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Microtubule-associated protein tau

Chain A: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: Microtubule-associated protein tau
Chain B: 100%
• Molecule 1: Microtubule-associated protein tau
Chain C: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: Microtubule-associated protein tau
Chain D: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: Microtubule-associated protein tau
Chain E: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: Microtubule-associated protein tau
27%
Chain F: 100%
ACE349



• Molecule 1: Microtubule-associated protein tau
Chain G: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: Microtubule-associated protein tau
7% Chain H: 100%
Chain H: 100%
\bullet Molecule 1: Microtubule-associated protein tau
Chain I: 100%
There are no outlier residues recorded for this chain.
\bullet Molecule 1: Microtubule-associated protein tau
Chain J: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: Microtubule-associated protein tau
Chain K: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: Microtubule-associated protein tau 73%
Chain L: 100%
ACE349 V350 C355 C355 C355 C355 C355 C355 C355 C
\bullet Molecule 1: Microtubule-associated protein tau
Chain M: 100%
There are no outlier residues recorded for this chain.
\bullet Molecule 1: Microtubule-associated protein tau
Chain N: 100%



A CE349 NH22363 NH22363		
• Molecule 1: Microtubule	-associated protein tau	
Chain O:	100%	
There are no outlier residu	ues recorded for this chain.	
• Molecule 1: Microtubule	-associated protein tau	
Chain P:	100%	
There are no outlier residu	ues recorded for this chain.	
• Molecule 1: Microtubule	-associated protein tau	
Chain Q:	100%	
There are no outlier residu	ies recorded for this chain.	
• Molecule 1: Microtubule	-associated protein tau	
Chain R:	100%	
ACE349 V350 Q351 S352 K353 I354 Q355 C355 C355 C355 S356 L357 NH2363		
• Molecule 1: Microtubule	-associated protein tau	
Chain S:	100%	
There are no outlier residu	ies recorded for this chain.	
• Molecule 1: Microtubule	-associated protein tau	
Chain T:	100%	
ADER349 HI362 HI12365		
• Molecule 1: Microtubule	-associated protein tau	
Chain U:	100%	
There are no outlier residu	ies recorded for this chain.	
• Molecule 1: Microtubule	-associated protein tau	



Chain V:	100%	
There are no outlier r	residues recorded for this chain.	
• Molecule 1: Microtu	ubule-associated protein tau	
Chain W:	100%	
There are no outlier r	residues recorded for this chain.	
• Molecule 1: Microtu	ubule-associated protein tau	
	73%	
Chain X:	100%	
ACE349 V350 Q351 S352 K353 K353 C355 S356 C355 S356 L357 D358 N359		
• Molecule 1: Microtu	ubule-associated protein tau	
Chain Y:	100%	
There are no outlier r	residues recorded for this chain.	
• Molecule 1: Microtu	ubule-associated protein tau	
7%		
Chain Z:	100%	
ACE349 NH2363		
• Molecule 1: Microtu	ubule-associated protein tau	
Chain a:	100%	
	residues recorded for this chain.	
• Molecule 1: Micrott	ubule-associated protein tau	
Chain b:	100%	
There are no outlier r	residues recorded for this chain.	
• Molecule 1: Microtu	ubule-associated protein tau	
Chain c:	100%	
There are no outlier r	residues recorded for this chain.	
• Molecule 1: Microtu	ubule-associated protein tau	



40%
Chain d: 100%
ACE349 V350 C355 C355 C355 C355 C355 C355 C355 C
• Molecule 1: Microtubule-associated protein tau
Chain e: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: Microtubule-associated protein tau
Chain f: 100%
• Molecule 1: Microtubule-associated protein tau
Chain g: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: Microtubule-associated protein tau
Chain h: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: Microtubule-associated protein tau
Chain i: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: Microtubule-associated protein tau
67%
Chain j: 100%
ACE349 V 3550 C 3551 S 3525 S 3566 L 355 D 358 D 358 N H2363



4 Experimental information (i)

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=-0.68°, rise=4.86 Å, axial	Depositor
	sym=C1	
Number of segments used	17820	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	39	Depositor
Minimum defocus (nm)	1400	Depositor
Maximum defocus (nm)	2600	Depositor
Magnification	130000	Depositor
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	0.070	Depositor
Minimum map value	-0.030	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.004	Depositor
Recommended contour level	0.014	Depositor
Map size (Å)	285.0, 285.0, 285.0	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles $(^{\circ})$	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.95, 0.95, 0.95	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NH2, ACE $\,$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

N.T. 1		Bond	lengths	Bond	angles
Mol	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.38	0/99	0.49	0/133
1	В	0.32	0/99	0.48	0/133
1	С	0.29	0/99	0.47	0/133
1	D	0.32	0/99	0.46	0/133
1	Е	0.34	0/99	0.48	0/133
1	F	0.23	0/99	0.44	0/133
1	G	0.38	0/99	0.49	0/133
1	Н	0.32	0/99	0.48	0/133
1	Ι	0.29	0/99	0.47	0/133
1	J	0.32	0/99	0.46	0/133
1	Κ	0.34	0/99	0.48	0/133
1	L	0.23	0/99	0.44	0/133
1	М	0.38	0/99	0.49	0/133
1	N	0.32	0/99	0.48	0/133
1	0	0.29	0/99	0.47	0/133
1	Р	0.33	0/99	0.46	0/133
1	Q	0.35	0/99	0.48	0/133
1	R	0.23	0/99	0.44	0/133
1	S	0.38	0/99	0.49	0/133
1	Т	0.32	0/99	0.48	0/133
1	U	0.29	0/99	0.47	0/133
1	V	0.33	0/99	0.46	0/133
1	W	0.34	0/99	0.48	0/133
1	Х	0.23	0/99	0.44	0/133
1	Y	0.38	0/99	0.49	0/133
1	Ζ	0.32	0/99	0.48	0/133
1	a	0.29	0/99	0.47	0/133
1	b	0.33	0/99	0.46	0/133
1	с	0.34	0/99	0.48	0/133
1	d	0.23	0/99	0.44	0/133
1	е	0.38	0/99	0.49	0/133
1	f	0.32	0/99	0.47	0/133
L	I	I	/	1	/



Mol	Chain	Bond lengths		Bond angles		
	Unam	RMSZ $ $ $# Z > 5$		RMSZ	# Z > 5	
1	g	0.29	0/99	0.47	0/133	
1	h	0.33	0/99	0.46	0/133	
1	i	0.35	0/99	0.48	0/133	
1	j	0.23	0/99	0.44	0/133	
All	All	0.32	0/3564	0.47	0/4788	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	13/15~(87%)	13 (100%)	0	0	100 100
1	В	13/15~(87%)	13 (100%)	0	0	100 100
1	С	13/15~(87%)	13 (100%)	0	0	100 100
1	D	13/15~(87%)	13 (100%)	0	0	100 100
1	Ε	13/15~(87%)	13 (100%)	0	0	100 100
1	F	13/15~(87%)	13 (100%)	0	0	100 100
1	G	13/15~(87%)	13 (100%)	0	0	100 100
1	Н	13/15~(87%)	13 (100%)	0	0	100 100
1	Ι	13/15~(87%)	13 (100%)	0	0	100 100



Mol	Chain	n previous page Analysed	Favoured	Allowed	Outliers	-	Perce	ntiles
1	J	13/15~(87%)	13 (100%)	0	0		100	100
1	К	13/15~(87%)	13 (100%)	0	0		100	100
1	L	13/15~(87%)	13 (100%)	0	0		100	100
1	М	13/15~(87%)	13 (100%)	0	0		100	100
1	Ν	13/15~(87%)	13 (100%)	0	0		100	100
1	Ο	13/15~(87%)	13 (100%)	0	0		100	100
1	Р	13/15~(87%)	13 (100%)	0	0		100	100
1	Q	13/15~(87%)	13 (100%)	0	0		100	100
1	R	13/15~(87%)	13 (100%)	0	0		100	100
1	S	13/15~(87%)	13 (100%)	0	0		100	100
1	Т	13/15~(87%)	13 (100%)	0	0		100	100
1	U	13/15~(87%)	13 (100%)	0	0		100	100
1	V	13/15~(87%)	13 (100%)	0	0		100	100
1	W	13/15~(87%)	13 (100%)	0	0		100	100
1	Х	13/15~(87%)	13 (100%)	0	0		100	100
1	Y	13/15~(87%)	13 (100%)	0	0		100	100
1	Ζ	13/15~(87%)	13 (100%)	0	0		100	100
1	a	13/15~(87%)	13 (100%)	0	0		100	100
1	b	13/15~(87%)	13 (100%)	0	0		100	100
1	с	13/15~(87%)	13 (100%)	0	0		100	100
1	d	13/15~(87%)	13 (100%)	0	0		100	100
1	е	13/15~(87%)	13 (100%)	0	0		100	100
1	f	13/15~(87%)	13 (100%)	0	0		100	100
1	g	13/15~(87%)	13 (100%)	0	0		100	100
1	h	13/15~(87%)	13 (100%)	0	0		100	100
1	i	13/15~(87%)	13 (100%)	0	0		100	100
1	j	13/15~(87%)	13 (100%)	0	0		100	100
All	All	468/540 (87%)	468 (100%)	0	0		100	100

Continued from previous page...

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	А	12/12~(100%)	12 (100%)	0	100	100
1	В	12/12~(100%)	12 (100%)	0	100	100
1	С	12/12~(100%)	12 (100%)	0	100	100
1	D	12/12~(100%)	12 (100%)	0	100	100
1	Е	12/12~(100%)	12 (100%)	0	100	100
1	F	12/12~(100%)	12 (100%)	0	100	100
1	G	12/12~(100%)	12 (100%)	0	100	100
1	Н	12/12~(100%)	12 (100%)	0	100	100
1	Ι	12/12~(100%)	12 (100%)	0	100	100
1	J	12/12~(100%)	12 (100%)	0	100	100
1	К	12/12~(100%)	12 (100%)	0	100	100
1	L	12/12~(100%)	12 (100%)	0	100	100
1	М	12/12~(100%)	12 (100%)	0	100	100
1	Ν	12/12~(100%)	12 (100%)	0	100	100
1	Ο	12/12~(100%)	12 (100%)	0	100	100
1	Р	12/12~(100%)	12 (100%)	0	100	100
1	Q	12/12~(100%)	12 (100%)	0	100	100
1	R	12/12~(100%)	12 (100%)	0	100	100
1	S	12/12~(100%)	12 (100%)	0	100	100
1	Т	12/12~(100%)	12 (100%)	0	100	100
1	U	12/12~(100%)	12 (100%)	0	100	100
1	V	12/12~(100%)	12 (100%)	0	100	100
1	W	12/12~(100%)	12 (100%)	0	100	100
1	Х	12/12~(100%)	12 (100%)	0	100	100
1	Y	12/12~(100%)	12 (100%)	0	100	100
1	Ζ	12/12~(100%)	12 (100%)	0	100	100



Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	a	12/12~(100%)	12~(100%)	0	100	100
1	b	12/12~(100%)	12 (100%)	0	100	100
1	с	12/12~(100%)	12 (100%)	0	100	100
1	d	12/12~(100%)	12 (100%)	0	100	100
1	е	12/12~(100%)	12 (100%)	0	100	100
1	f	12/12~(100%)	12 (100%)	0	100	100
1	g	12/12~(100%)	12~(100%)	0	100	100
1	h	12/12~(100%)	12~(100%)	0	100	100
1	i	12/12~(100%)	12~(100%)	0	100	100
1	j	12/12~(100%)	12 (100%)	0	100	100
All	All	432/432~(100%)	432~(100%)	0	100	100

Continued from previous page...

There are no protein residues with a non-rotameric sidechain to report.

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such side chains are listed below:

Mol	Chain	Res	Type
1	Т	362	HIS
1	f	362	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-16876. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



6.1.2 Raw map




The images above show the map projected in three orthogonal directions.





6.2 Central slices (i)

6.2.1 Primary map




X Index: 150





Z Index: 150

6.2.2 Raw map



X Index: 150

Y Index: 150




The images above show central slices of the map in three orthogonal directions.





6.3 Largest variance slices (i)

6.3.1 Primary map




X Index: 151





Z Index: 130

6.3.2 Raw map



X Index: 151

Y Index: 148




The images above show the largest variance slices of the map in three orthogonal directions.




6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



6.4.2 Raw map




The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.014. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is 272 nm^3 ; this corresponds to an approximate mass of 246 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)




*Reported resolution corresponds to spatial frequency of 0.385 ${\rm \AA^{-1}}$



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.385 $\mathrm{\AA^{-1}}$



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
Resolution estimate (A)	0.143	0.5	Half-bit
Reported by author	2.60	-	-
Author-provided FSC curve	2.63	3.28	2.74
Unmasked-calculated*	3.39	6.00	3.47


*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.39 differs from the reported value 2.6 by more than 10 %



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-16876 and PDB model 80H2. Per-residue inclusion information can be found in section 3 on page 11.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.014 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.014).



9.4 Atom inclusion (i)



At the recommended contour level, 80% of all backbone atoms, 80% of all non-hydrogen atoms, are inside the map.



1.0

0.0 <0.0

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.014) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8010	0.6400
А	0.9510	0.6730
В	0.8220	0.6530
С	0.8610	0.6280
D	0.9010	0.6740
E	0.8810	0.6570
F	0.5540	0.6140
G	0.9410	0.6770
Н	0.7030	0.5970
Ι	0.8910	0.6520
J	0.9110	0.6600
K	0.8810	0.6490
L	0.3960	0.5690
М	0.9410	0.6750
N	0.8120	0.6450
0	0.8810	0.6260
Р	0.9210	0.6800
Q	0.8420	0.6430
R	0.5450	0.6120
S	0.9410	0.6800
Т	0.7130	0.6310
U	0.8810	0.6500
V	0.9210	0.6600
W	0.8810	0.6310
Х	0.3860	0.5790
Y	0.9410	0.6680
Z	0.7820	0.6390
a	0.8520	0.6130
b	0.9010	0.6800
с	0.8610	0.6420
d	0.5150	0.6110
e	0.9210	0.6740
f	0.6930	0.6020
g	0.8910	0.6530
h	0.9210	0.6560



Continued from previous page...

Chain	Atom inclusion	Q-score
i	0.8420	0.6210
j	0.3760	0.5600

