



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 8, 2023 – 05:35 am GMT

PDB ID : 1OI8
Title : 5'-Nucleotidase (E. coli) with an Engineered Disulfide Bridge (P90C, L424C)
Authors : Schultz-Heienbrok, R.; Maier, T.; Straeter, N.
Deposited on : 2003-06-10
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	532	
1	B	532	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9132 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN USHA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	526	4097	2585	704	789	19	0	0	1
1	B	526	4097	2585	704	789	19	0	0	1

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	90	CYS	PRO	engineered mutation	UNP P07024
A	424	CYS	LEU	engineered mutation	UNP P07024
B	90	CYS	PRO	engineered mutation	UNP P07024
B	424	CYS	LEU	engineered mutation	UNP P07024

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total 2	Mn 2	0	0
2	B	2	Total 2	Mn 2	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is CARBONATE ION (three-letter code: CO3) (formula: CO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	1	3		
4	B	1	Total	C	O	0	0
			4	1	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	475	Total 475	O 475	0	0
5	B	441	Total 441	O 441	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	97.69Å 97.69Å 312.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.96 – 2.10 19.89 – 2.10	Depositor EDS
% Data completeness (in resolution range)	95.3 (19.96-2.10) 95.5 (19.89-2.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.17 (at 2.09Å)	Xtrriage
Refinement program	REFMAC 5.1.19	Depositor
R, R_{free}	0.156 , 0.205 0.175 , 0.169	Depositor DCC
R_{free} test set	4275 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	28.6	Xtrriage
Anisotropy	0.001	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 49.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9132	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 48.67 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.3209e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CO3, SO4, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.95	3/4184 (0.1%)	1.01	19/5661 (0.3%)
1	B	0.99	6/4184 (0.1%)	0.96	15/5661 (0.3%)
All	All	0.97	9/8368 (0.1%)	0.99	34/11322 (0.3%)

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	345	MET	SD-CE	-9.83	1.22	1.77
1	B	235	GLU	CG-CD	7.13	1.62	1.51
1	B	362	VAL	CB-CG1	-6.50	1.39	1.52
1	B	438	MET	CG-SD	6.10	1.97	1.81
1	A	362	VAL	CB-CG2	-5.84	1.40	1.52

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	190	ARG	NE-CZ-NH1	13.84	127.22	120.30
1	A	190	ARG	NE-CZ-NH2	-13.46	113.57	120.30
1	A	410	ARG	NE-CZ-NH2	-10.49	115.05	120.30
1	A	410	ARG	NE-CZ-NH1	9.87	125.23	120.30
1	B	235	GLU	OE1-CD-OE2	-9.43	111.99	123.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4097	0	4009	22	0
1	B	4097	0	4009	32	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	5	0	0	1	0
3	B	5	0	0	0	0
4	A	4	0	0	0	0
4	B	4	0	0	1	0
5	A	475	0	0	4	0
5	B	441	0	0	10	0
All	All	9132	0	8018	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 55 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:217:HIS:CD2	1:B:252:HIS:HB2	2.14	0.81
1:A:217:HIS:CD2	1:A:252:HIS:HB2	2.15	0.81
1:B:359:GLN:O	1:B:362:VAL:HG12	1.82	0.80
1:A:34:ILE:HD11	1:A:163:LEU:CD1	2.18	0.73
1:A:190:ARG:HD2	1:A:195:GLU:OE1	1.91	0.70

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	524/532 (98%)	509 (97%)	13 (2%)	2 (0%)	34 32

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	524/532 (98%)	509 (97%)	13 (2%)	2 (0%)	34 32
All	All	1048/1064 (98%)	1018 (97%)	26 (2%)	4 (0%)	34 32

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	289	HIS
1	A	87	THR
1	B	87	THR
1	B	289	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	433/440 (98%)	416 (96%)	17 (4%)	32 33
1	B	433/440 (98%)	423 (98%)	10 (2%)	50 55
All	All	866/880 (98%)	839 (97%)	27 (3%)	40 43

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	481	LYS
1	B	78	LEU
1	B	471	LYS
1	A	539	SER
1	B	159	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	476	ASN
1	B	499	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	550	GLN
1	A	511	ASN
1	B	201	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	CO3	B	1554	2	2,3,3	0.41	0	2,3,3	1.97	1 (50%)
3	SO4	A	1553	-	4,4,4	0.67	0	6,6,6	1.74	2 (33%)
4	CO3	A	1554	2	2,3,3	0.18	0	2,3,3	0.81	0
3	SO4	B	1553	-	4,4,4	0.48	0	6,6,6	0.79	0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1553	SO4	O3-S-O2	-3.07	93.31	109.31
4	B	1554	CO3	O2-C-O1	2.69	126.53	119.55
3	A	1553	SO4	O4-S-O2	2.08	120.16	109.31

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1554	CO3	1	0
3	A	1553	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	526/532 (98%)	-0.49	6 (1%) 80 84	15, 23, 37, 53	0
1	B	526/532 (98%)	-0.42	8 (1%) 73 77	14, 24, 38, 54	0
All	All	1052/1064 (98%)	-0.45	14 (1%) 77 80	14, 23, 38, 54	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	551	GLU	5.9
1	B	472	ASP	4.7
1	A	550	GLN	3.7
1	B	162	ASP	3.5
1	A	267	VAL	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	CO3	A	1554	4/4	0.84	0.13	26,33,34,35	0
4	CO3	B	1554	4/4	0.86	0.12	35,36,37,38	0
3	SO4	A	1553	5/5	0.87	0.20	44,47,59,59	0
3	SO4	B	1553	5/5	0.88	0.24	51,58,64,68	0
2	MN	A	1551	1/1	1.00	0.06	21,21,21,21	0
2	MN	A	1552	1/1	1.00	0.02	29,29,29,29	0
2	MN	B	1551	1/1	1.00	0.01	32,32,32,32	0
2	MN	B	1552	1/1	1.00	0.07	21,21,21,21	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.