



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 29, 2023 – 11:37 PM EDT

PDB ID : 3OI7  
Title : Structure of the structure of the H13A mutant of Ykr043C in complex with s edoheptulose-1,7-bisphosphate  
Authors : Singer, A.U.; Xu, X.; Dong, A.; Cui, H.; Clasquin, M.F.; Caudy, A.A.; Edwards, A.M.; Savchenko, A.; Joachimiak, A.; Yakunin, A.F.; Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2010-08-18  
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	292	
1	B	292	
1	C	292	
1	D	292	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	EDO	D	274	-	-	-	X

## 2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 8833 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein YKR043C.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	260	2109	1327	381	394	4	3	0	1	0
1	B	260	2117	1331	383	396	4	3	0	2	0
1	C	260	2109	1327	381	394	4	3	0	1	0
1	D	260	2117	1332	384	394	4	3	0	2	0

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MSE	-	expression tag	UNP P36136
A	-19	GLY	-	expression tag	UNP P36136
A	-18	SER	-	expression tag	UNP P36136
A	-17	SER	-	expression tag	UNP P36136
A	-16	HIS	-	expression tag	UNP P36136
A	-15	HIS	-	expression tag	UNP P36136
A	-14	HIS	-	expression tag	UNP P36136
A	-13	HIS	-	expression tag	UNP P36136
A	-12	HIS	-	expression tag	UNP P36136
A	-11	HIS	-	expression tag	UNP P36136
A	-10	SER	-	expression tag	UNP P36136
A	-9	SER	-	expression tag	UNP P36136
A	-8	GLY	-	expression tag	UNP P36136
A	-7	ARG	-	expression tag	UNP P36136
A	-6	GLU	-	expression tag	UNP P36136
A	-5	ASN	-	expression tag	UNP P36136
A	-4	LEU	-	expression tag	UNP P36136
A	-3	TYR	-	expression tag	UNP P36136
A	-2	PHE	-	expression tag	UNP P36136
A	-1	GLN	-	expression tag	UNP P36136
A	0	GLY	-	expression tag	UNP P36136

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
A	13	ALA	HIS	engineered mutation	UNP P36136
B	-20	MSE	-	expression tag	UNP P36136
B	-19	GLY	-	expression tag	UNP P36136
B	-18	SER	-	expression tag	UNP P36136
B	-17	SER	-	expression tag	UNP P36136
B	-16	HIS	-	expression tag	UNP P36136
B	-15	HIS	-	expression tag	UNP P36136
B	-14	HIS	-	expression tag	UNP P36136
B	-13	HIS	-	expression tag	UNP P36136
B	-12	HIS	-	expression tag	UNP P36136
B	-11	HIS	-	expression tag	UNP P36136
B	-10	SER	-	expression tag	UNP P36136
B	-9	SER	-	expression tag	UNP P36136
B	-8	GLY	-	expression tag	UNP P36136
B	-7	ARG	-	expression tag	UNP P36136
B	-6	GLU	-	expression tag	UNP P36136
B	-5	ASN	-	expression tag	UNP P36136
B	-4	LEU	-	expression tag	UNP P36136
B	-3	TYR	-	expression tag	UNP P36136
B	-2	PHE	-	expression tag	UNP P36136
B	-1	GLN	-	expression tag	UNP P36136
B	0	GLY	-	expression tag	UNP P36136
B	13	ALA	HIS	engineered mutation	UNP P36136
C	-20	MSE	-	expression tag	UNP P36136
C	-19	GLY	-	expression tag	UNP P36136
C	-18	SER	-	expression tag	UNP P36136
C	-17	SER	-	expression tag	UNP P36136
C	-16	HIS	-	expression tag	UNP P36136
C	-15	HIS	-	expression tag	UNP P36136
C	-14	HIS	-	expression tag	UNP P36136
C	-13	HIS	-	expression tag	UNP P36136
C	-12	HIS	-	expression tag	UNP P36136
C	-11	HIS	-	expression tag	UNP P36136
C	-10	SER	-	expression tag	UNP P36136
C	-9	SER	-	expression tag	UNP P36136
C	-8	GLY	-	expression tag	UNP P36136
C	-7	ARG	-	expression tag	UNP P36136
C	-6	GLU	-	expression tag	UNP P36136
C	-5	ASN	-	expression tag	UNP P36136
C	-4	LEU	-	expression tag	UNP P36136
C	-3	TYR	-	expression tag	UNP P36136
C	-2	PHE	-	expression tag	UNP P36136

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	-1	GLN	-	expression tag	UNP P36136
C	0	GLY	-	expression tag	UNP P36136
C	13	ALA	HIS	engineered mutation	UNP P36136
D	-20	MSE	-	expression tag	UNP P36136
D	-19	GLY	-	expression tag	UNP P36136
D	-18	SER	-	expression tag	UNP P36136
D	-17	SER	-	expression tag	UNP P36136
D	-16	HIS	-	expression tag	UNP P36136
D	-15	HIS	-	expression tag	UNP P36136
D	-14	HIS	-	expression tag	UNP P36136
D	-13	HIS	-	expression tag	UNP P36136
D	-12	HIS	-	expression tag	UNP P36136
D	-11	HIS	-	expression tag	UNP P36136
D	-10	SER	-	expression tag	UNP P36136
D	-9	SER	-	expression tag	UNP P36136
D	-8	GLY	-	expression tag	UNP P36136
D	-7	ARG	-	expression tag	UNP P36136
D	-6	GLU	-	expression tag	UNP P36136
D	-5	ASN	-	expression tag	UNP P36136
D	-4	LEU	-	expression tag	UNP P36136
D	-3	TYR	-	expression tag	UNP P36136
D	-2	PHE	-	expression tag	UNP P36136
D	-1	GLN	-	expression tag	UNP P36136
D	0	GLY	-	expression tag	UNP P36136
D	13	ALA	HIS	engineered mutation	UNP P36136

- Molecule 2 is 1,7-di-O-phosphono-beta-D-altro-hept-2-ulofuranose (three-letter code: OI7) (formula: C<sub>7</sub>H<sub>16</sub>O<sub>13</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	O			P
2	A	1	22	7	13	2	0	0
2	B	1	22	7	13	2	0	0
2	C	1	22	7	13	2	0	0
2	D	1	22	7	13	2	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
3	A	1	1	1	0	0
3	B	1	1	1	0	0
3	C	1	1	1	0	0
3	D	1	1	1	0	0

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Na		
4	A	1	1	1	0	0


*Continued on next page...*



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Na 1 1	0	0
4	C	2	Total Na 2 2	0	0
4	D	1	Total Na 1 1	0	0

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	C	1	Total C O 4 2 2	0	0
5	C	1	Total C O 4 2 2	0	0
5	D	1	Total C O 4 2 2	0	0
5	D	1	Total C O 4 2 2	0	0
5	D	1	Total C O 4 2 2	0	0

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	C O	0	0
			6	3 3		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	60	Total	O	0	0
			60	60		
7	B	58	Total	O	0	0
			58	58		
7	C	65	Total	O	0	0
			65	65		
7	D	63	Total	O	0	0
			63	63		





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.56Å 74.94Å 83.61Å 90.04° 89.92° 77.22°	Depositor
Resolution (Å)	27.50 – 2.40 27.50 – 2.40	Depositor EDS
% Data completeness (in resolution range)	91.3 (27.50-2.40) 91.0 (27.50-2.40)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.98 (at 2.39Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.228 , 0.260 0.228 , 0.260	Depositor DCC
$R_{free}$ test set	2567 reflections (5.19%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.8	Xtrriage
Anisotropy	0.078	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 16.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.53$ , $\langle L^2 \rangle = 0.36$	Xtrriage
Estimated twinning fraction	0.448 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	8833	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 30.38 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.3206e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GOL, NA, EDO, OI7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.58	0/2153	0.66	0/2912
1	B	0.58	0/2161	0.67	1/2923 (0.0%)
1	C	0.58	0/2153	0.66	0/2912
1	D	0.59	0/2164	0.67	0/2926
All	All	0.58	0/8631	0.67	1/11673 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	104	ASP	CB-CG-OD1	5.22	123.00	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2109	0	2070	19	0
1	B	2117	0	2075	22	0
1	C	2109	0	2070	26	0
1	D	2117	0	2083	27	0
2	A	22	0	12	3	0
2	B	22	0	11	5	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	22	0	12	5	0
2	D	22	0	12	5	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	2	0	0	0	0
4	D	1	0	0	0	0
5	A	8	0	12	0	0
5	B	4	0	6	2	0
5	C	8	0	12	3	0
5	D	12	0	18	2	0
6	A	6	0	8	0	0
7	A	60	0	0	2	0
7	B	58	0	0	2	0
7	C	65	0	0	2	0
7	D	63	0	0	2	0
All	All	8833	0	8401	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 88 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:18:TRP:HE1	1:D:161:LYS:NZ	1.62	0.96
1:C:18:TRP:HE1	1:D:161:LYS:HZ3	1.06	0.95
1:D:62:ILE:CD1	1:D:78:LEU:HD11	2.07	0.84
1:B:62:ILE:CD1	1:B:78:LEU:HD11	2.09	0.83
1:A:62:ILE:CD1	1:A:78:LEU:HD11	2.08	0.82

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/292 (89%)	250 (96%)	8 (3%)	1 (0%)	34	48
1	B	260/292 (89%)	251 (96%)	8 (3%)	1 (0%)	34	48
1	C	259/292 (89%)	249 (96%)	9 (4%)	1 (0%)	34	48
1	D	260/292 (89%)	252 (97%)	7 (3%)	1 (0%)	34	48
All	All	1038/1168 (89%)	1002 (96%)	32 (3%)	4 (0%)	34	48

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	175	ALA
1	C	175	ALA
1	D	175	ALA
1	A	175	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/249 (91%)	221 (97%)	6 (3%)	46	66
1	B	228/249 (92%)	222 (97%)	6 (3%)	46	66
1	C	227/249 (91%)	221 (97%)	6 (3%)	46	66
1	D	228/249 (92%)	219 (96%)	9 (4%)	32	50
All	All	910/996 (91%)	883 (97%)	27 (3%)	42	61



5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	88	LYS
1	C	233	ASP
1	D	194	LYS
1	C	199	ILE
1	D	25	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	142	GLN
1	D	15	GLN
1	C	15	GLN
1	B	142	GLN
1	C	142	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 9 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	GOL	A	275	-	5,5,5	0.43	0	5,5,5	0.26	0
5	EDO	D	274	-	3,3,3	0.84	0	2,2,2	0.30	0
2	OI7	A	301	3	20,22,22	1.40	2 (10%)	27,35,35	2.40	5 (18%)
5	EDO	D	275	-	3,3,3	0.51	0	2,2,2	0.32	0
5	EDO	C	274	-	3,3,3	0.43	0	2,2,2	0.42	0
2	OI7	C	301	4,3	20,22,22	1.44	4 (20%)	27,35,35	2.35	5 (18%)
5	EDO	A	274	-	3,3,3	0.58	0	2,2,2	0.03	0
5	EDO	B	273	-	3,3,3	0.57	0	2,2,2	0.44	0
2	OI7	B	301	4	20,22,22	1.47	4 (20%)	27,35,35	2.34	5 (18%)
5	EDO	D	273	-	3,3,3	0.44	0	2,2,2	0.40	0
5	EDO	A	273	-	3,3,3	0.61	0	2,2,2	0.32	0
2	OI7	D	301	4,3	20,22,22	1.40	3 (15%)	27,35,35	2.36	4 (14%)
5	EDO	C	275	-	3,3,3	0.55	0	2,2,2	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	275	-	-	3/4/4/4	-
5	EDO	D	274	-	-	1/1/1/1	-
2	OI7	A	301	3	-	9/17/36/36	0/1/1/1
5	EDO	D	275	-	-	1/1/1/1	-
5	EDO	C	274	-	-	1/1/1/1	-
2	OI7	C	301	4,3	-	9/17/36/36	0/1/1/1
5	EDO	A	274	-	-	0/1/1/1	-
5	EDO	B	273	-	-	0/1/1/1	-
2	OI7	B	301	4	-	9/17/36/36	0/1/1/1
5	EDO	D	273	-	-	1/1/1/1	-
5	EDO	A	273	-	-	1/1/1/1	-
2	OI7	D	301	4,3	-	9/17/36/36	0/1/1/1
5	EDO	C	275	-	-	1/1/1/1	-

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	301	OI7	O3-C3	2.79	1.48	1.42
2	A	301	OI7	O3-C3	2.76	1.48	1.42
2	D	301	OI7	O3-C3	2.68	1.48	1.42

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	OI7	O3-C3	2.67	1.48	1.42
2	B	301	OI7	O4-C4	-2.65	1.36	1.43

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	OI7	C6-C5-C4	-9.49	103.17	115.86
2	D	301	OI7	C6-C5-C4	-9.20	103.55	115.86
2	B	301	OI7	C6-C5-C4	-9.19	103.57	115.86
2	C	301	OI7	C6-C5-C4	-9.15	103.62	115.86
2	B	301	OI7	O5-C5-C4	5.08	111.69	103.59

There are no chirality outliers.

5 of 45 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	OI7	C4-C5-C6-C7
2	A	301	OI7	C4-C5-C6-O6
2	A	301	OI7	O5-C5-C6-C7
2	A	301	OI7	O5-C5-C6-O6
2	A	301	OI7	O1-C1-C2-O5


There are no ring outliers.


7 monomers are involved in 25 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	OI7	3	0
5	C	274	EDO	3	0
2	C	301	OI7	5	0
5	B	273	EDO	2	0
2	B	301	OI7	5	0
5	D	273	EDO	2	0
2	D	301	OI7	5	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	257/292 (88%)	-0.12	2 (0%) 86 84	11, 21, 42, 57	0
1	B	257/292 (88%)	-0.11	2 (0%) 86 84	11, 21, 42, 57	0
1	C	257/292 (88%)	-0.13	0 100 100	11, 21, 42, 57	0
1	D	257/292 (88%)	-0.12	1 (0%) 92 91	11, 21, 42, 57	0
All	All	1028/1168 (88%)	-0.12	5 (0%) 91 89	11, 21, 42, 57	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	202	ILE	2.9
1	A	4	LEU	2.8
1	D	199	ILE	2.8
1	B	50	ASN	2.4
1	B	202	ILE	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,






median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	EDO	D	274	4/4	0.39	0.54	93,94,94,94	0
5	EDO	C	275	4/4	0.84	0.24	35,38,38,39	0
6	GOL	A	275	6/6	0.84	0.16	44,47,48,49	0
5	EDO	D	273	4/4	0.85	0.16	21,27,31,35	0
4	NA	C	273	1/1	0.86	0.14	27,27,27,27	0
5	EDO	A	273	4/4	0.88	0.15	20,25,28,31	0
5	EDO	C	274	4/4	0.91	0.14	15,20,23,28	0
5	EDO	D	275	4/4	0.92	0.16	36,36,37,38	0
5	EDO	B	273	4/4	0.92	0.15	16,23,26,29	0
2	OI7	C	301	22/22	0.93	0.16	21,41,48,51	0
2	OI7	D	301	22/22	0.93	0.19	21,41,48,52	0
3	MG	B	311	1/1	0.93	0.16	30,30,30,30	0
2	OI7	A	301	22/22	0.93	0.19	21,41,48,52	0
4	NA	D	272	1/1	0.93	0.12	25,25,25,25	0
2	OI7	B	301	22/22	0.93	0.18	21,41,48,51	0
5	EDO	A	274	4/4	0.93	0.14	31,34,34,35	0
4	NA	B	272	1/1	0.95	0.15	24,24,24,24	0
4	NA	C	272	1/1	0.96	0.21	25,25,25,25	0
3	MG	D	311	1/1	0.96	0.16	27,27,27,27	0
4	NA	A	272	1/1	0.97	0.15	29,29,29,29	0
3	MG	A	311	1/1	0.97	0.22	30,30,30,30	0
3	MG	C	311	1/1	0.98	0.18	29,29,29,29	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around OI7 C 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around OI7 D 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around OI7 A 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around OI7 B 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.