



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 12, 2023 – 08:15 PM EDT

PDB ID : 4OIF
Title : 3D structure of Gan42B, a GH42 beta-galactosidase from G.
Authors : Solomon, H.V.; Tabachnikov, O.; Feinberg, H.; Shoham, Y.; Shoham, G.
Deposited on : 2014-01-19
Resolution : 2.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1564 (2.46-2.42)
Clashscore	141614	1631 (2.46-2.42)
Ramachandran outliers	138981	1617 (2.46-2.42)
Sidechain outliers	138945	1617 (2.46-2.42)
RSRZ outliers	127900	1547 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	686	 88% 11%
1	B	686	 88% 11%
1	C	686	 89% 10%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 18838 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-galactosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	685	5735	3675	989	1048	23	0	14	0
1	B	684	5724	3667	985	1050	22	0	15	0
1	C	684	5722	3665	984	1051	22	0	14	0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		
2	C	1	Total	Zn	0	0
			1	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total 6	C 3	O 3	0	0
3	B	1	Total 6	C 3	O 3	0	0
3	B	1	Total 6	C 3	O 3	0	0
3	B	1	Total 6	C 3	O 3	0	0
3	B	1	Total 6	C 3	O 3	0	0
3	B	1	Total 6	C 3	O 3	0	0
3	B	1	Total 6	C 3	O 3	0	0
3	C	1	Total 6	C 3	O 3	0	0
3	C	1	Total 6	C 3	O 3	0	0
3	C	1	Total 6	C 3	O 3	0	0
3	C	1	Total 6	C 3	O 3	0	0
3	C	1	Total 6	C 3	O 3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	484	Total 484	O 484	0	0
4	B	541	Total 541	O 541	0	0
4	C	479	Total 479	O 479	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-galactosidase



- Molecule 1: Beta-galactosidase



- Molecule 1: Beta-galactosidase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.84Å 181.35Å 196.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.86 – 2.45 48.86 – 2.45	Depositor EDS
% Data completeness (in resolution range)	96.2 (48.86-2.45) 95.3 (48.86-2.45)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.42	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.90 (at 2.45Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309), REFMAC	Depositor
R, R_{free}	0.133 , 0.204 0.136 , 0.206	Depositor DCC
R_{free} test set	4595 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	12.4	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 54.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	18838	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	1/5939 (0.0%)	0.55	1/8055 (0.0%)
1	B	0.41	0/5933	0.55	0/8049
1	C	0.42	1/5929 (0.0%)	0.56	1/8045 (0.0%)
All	All	0.41	2/17801 (0.0%)	0.55	2/24149 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
All	All	0	3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	518	TRP	CB-CG	-7.36	1.37	1.50
1	C	518	TRP	CB-CG	-6.92	1.37	1.50

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	518	TRP	CA-CB-CG	-7.35	99.73	113.70
1	A	518	TRP	CA-CB-CG	-6.32	101.69	113.70

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	360	TYR	Peptide
1	B	360	TYR	Peptide
1	C	360	TYR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5735	0	5541	64	0
1	B	5724	0	5519	59	0
1	C	5722	0	5504	53	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	48	0	64	9	0
3	B	72	0	96	7	0
3	C	30	0	40	7	0
4	A	484	0	0	14	0
4	B	541	0	0	6	0
4	C	479	0	0	9	0
All	All	18838	0	16764	170	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 170 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:94:ARG:NH1	4:B:917:HOH:O	1.95	0.98
1:A:549:ARG:NH2	4:A:950:HOH:O	2.11	0.83
1:A:649:GLU:HA	1:A:677:LYS:HG3	1.62	0.81
1:A:120:ARG:HH12	3:A:703:GOL:H2	1.45	0.80
1:C:120:ARG:HH12	3:C:702:GOL:H2	1.48	0.78

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	697/686 (102%)	668 (96%)	28 (4%)	1 (0%)	51	64
1	B	697/686 (102%)	674 (97%)	22 (3%)	1 (0%)	51	64
1	C	696/686 (102%)	669 (96%)	26 (4%)	1 (0%)	51	64
All	All	2090/2058 (102%)	2011 (96%)	76 (4%)	3 (0%)	51	64

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	262	HIS
1	A	4	LYS
1	C	262	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	615/602 (102%)	604 (98%)	11 (2%)	59	71
1	B	614/602 (102%)	605 (98%)	9 (2%)	65	76
1	C	613/602 (102%)	604 (98%)	9 (2%)	65	76
All	All	1842/1806 (102%)	1813 (98%)	29 (2%)	62	74

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	328	LEU
1	C	518	TRP
1	B	507	VAL
1	C	328	LEU
1	B	420	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 28 ligands modelled in this entry, 3 are monoatomic - leaving 25 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	B	707	-	5,5,5	0.42	0	5,5,5	0.73	0
3	GOL	C	705	-	5,5,5	0.36	0	5,5,5	0.28	0
3	GOL	A	704	-	5,5,5	0.35	0	5,5,5	0.33	0
3	GOL	A	702	-	5,5,5	0.34	0	5,5,5	0.55	0
3	GOL	C	703	-	5,5,5	0.38	0	5,5,5	0.48	0
3	GOL	C	706	-	5,5,5	0.43	0	5,5,5	0.33	0
3	GOL	A	703	-	5,5,5	0.43	0	5,5,5	0.58	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	705	-	5,5,5	0.30	0	5,5,5	0.43	0
3	GOL	A	708	-	5,5,5	0.38	0	5,5,5	0.33	0
3	GOL	B	704	-	5,5,5	0.32	0	5,5,5	0.27	0
3	GOL	B	713	-	5,5,5	0.52	0	5,5,5	0.37	0
3	GOL	C	704	-	5,5,5	0.52	0	5,5,5	0.28	0
3	GOL	B	712	-	5,5,5	0.39	0	5,5,5	0.32	0
3	GOL	B	702	-	5,5,5	0.37	0	5,5,5	0.57	0
3	GOL	C	702	-	5,5,5	0.41	0	5,5,5	0.67	0
3	GOL	A	705	-	5,5,5	0.36	0	5,5,5	0.53	0
3	GOL	A	709	-	5,5,5	0.42	0	5,5,5	0.19	0
3	GOL	B	711	-	5,5,5	0.36	0	5,5,5	0.51	0
3	GOL	A	707	-	5,5,5	0.35	0	5,5,5	0.42	0
3	GOL	B	708	-	5,5,5	0.39	0	5,5,5	0.48	0
3	GOL	A	706	-	5,5,5	0.32	0	5,5,5	0.82	0
3	GOL	B	706	-	5,5,5	0.34	0	5,5,5	0.44	0
3	GOL	B	709	-	5,5,5	0.29	0	5,5,5	0.50	0
3	GOL	B	710	-	5,5,5	0.34	0	5,5,5	0.29	0
3	GOL	B	703	-	5,5,5	0.39	0	5,5,5	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	707	-	-	2/4/4/4	-
3	GOL	C	705	-	-	2/4/4/4	-
3	GOL	A	704	-	-	2/4/4/4	-
3	GOL	A	702	-	-	0/4/4/4	-
3	GOL	C	703	-	-	2/4/4/4	-
3	GOL	C	706	-	-	2/4/4/4	-
3	GOL	A	703	-	-	2/4/4/4	-
3	GOL	B	705	-	-	0/4/4/4	-
3	GOL	A	708	-	-	2/4/4/4	-
3	GOL	B	704	-	-	0/4/4/4	-
3	GOL	B	713	-	-	3/4/4/4	-
3	GOL	C	704	-	-	2/4/4/4	-
3	GOL	B	712	-	-	2/4/4/4	-
3	GOL	B	702	-	-	0/4/4/4	-
3	GOL	C	702	-	-	3/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	705	-	-	4/4/4/4	-
3	GOL	A	709	-	-	2/4/4/4	-
3	GOL	B	711	-	-	2/4/4/4	-
3	GOL	A	707	-	-	2/4/4/4	-
3	GOL	B	708	-	-	2/4/4/4	-
3	GOL	A	706	-	-	4/4/4/4	-
3	GOL	B	706	-	-	4/4/4/4	-
3	GOL	B	709	-	-	0/4/4/4	-
3	GOL	B	710	-	-	3/4/4/4	-
3	GOL	B	703	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 47 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	703	GOL	O1-C1-C2-O2
3	A	704	GOL	O1-C1-C2-C3
3	A	706	GOL	C1-C2-C3-O3
3	A	708	GOL	O1-C1-C2-C3
3	A	709	GOL	O1-C1-C2-O2

There are no ring outliers.

16 monomers are involved in 23 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	707	GOL	1	0
3	C	705	GOL	1	0
3	A	702	GOL	1	0
3	C	703	GOL	1	0
3	C	706	GOL	1	0
3	A	703	GOL	2	0
3	A	708	GOL	1	0
3	B	713	GOL	1	0
3	C	704	GOL	1	0
3	B	712	GOL	2	0
3	C	702	GOL	3	0
3	A	705	GOL	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	707	GOL	1	0
3	B	708	GOL	1	0
3	A	706	GOL	2	0
3	B	710	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	685/686 (99%)	-0.71	7 (1%) 82 81	5, 12, 30, 79	0
1	B	684/686 (99%)	-0.85	2 (0%) 94 94	3, 10, 26, 65	0
1	C	684/686 (99%)	-0.83	2 (0%) 94 94	5, 12, 31, 62	0
All	All	2053/2058 (99%)	-0.80	11 (0%) 91 91	3, 11, 29, 79	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	3	PRO	4.8
1	A	4	LYS	3.9
1	C	518	TRP	3.5
1	A	518	TRP	3.0
1	B	518	TRP	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	C	706	6/6	0.86	0.18	23,33,34,43	0
3	GOL	B	712	6/6	0.89	0.17	21,27,31,37	0
3	GOL	A	709	6/6	0.89	0.21	23,38,56,56	0
3	GOL	C	705	6/6	0.91	0.15	21,39,43,44	0
3	GOL	C	704	6/6	0.91	0.18	13,38,42,43	0
3	GOL	A	708	6/6	0.94	0.15	20,24,33,43	0
3	GOL	A	705	6/6	0.94	0.15	15,19,30,38	0
3	GOL	A	706	6/6	0.94	0.16	15,28,34,38	0
3	GOL	A	704	6/6	0.95	0.13	27,29,48,61	0
3	GOL	B	713	6/6	0.95	0.22	16,26,37,47	0
3	GOL	A	707	6/6	0.95	0.13	12,31,34,36	0
3	GOL	B	710	6/6	0.95	0.18	13,25,26,30	0
3	GOL	B	711	6/6	0.95	0.15	16,26,36,62	0
3	GOL	B	708	6/6	0.96	0.09	16,17,24,27	0
3	GOL	C	702	6/6	0.97	0.12	10,10,22,23	0
3	GOL	A	703	6/6	0.97	0.11	11,15,17,21	0
3	GOL	B	709	6/6	0.97	0.16	17,22,25,34	0
3	GOL	B	706	6/6	0.97	0.15	10,28,34,44	0
3	GOL	A	702	6/6	0.98	0.10	9,14,15,17	0
3	GOL	C	703	6/6	0.98	0.11	11,13,24,27	0
3	GOL	B	707	6/6	0.98	0.11	7,13,14,23	0
3	GOL	B	703	6/6	0.98	0.11	6,10,15,16	0
3	GOL	B	704	6/6	0.98	0.13	8,11,15,15	0
3	GOL	B	705	6/6	0.99	0.09	6,13,14,16	0
3	GOL	B	702	6/6	0.99	0.10	6,8,10,12	0
2	ZN	B	701	1/1	1.00	0.05	12,12,12,12	0
2	ZN	C	701	1/1	1.00	0.04	16,16,16,16	0
2	ZN	A	701	1/1	1.00	0.04	13,13,13,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.