



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 12, 2023 – 09:33 PM EDT

PDB ID : 4OIN
Title : Crystal structure of Thermus thermophilus transcription initiation complex soaked with GE23077
Authors : Zhang, Y.; Ebright, R.H.; Arnold, E.
Deposited on : 2014-01-20
Resolution : 2.80 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	 2% 59% 14% 27%
1	B	315	 2% 60% 12% 28%
2	C	1119	 3% 83% 16% ..
3	D	1524	 7% 80% 17% ..
4	E	99	 85% 10% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	F	443	
6	G	19	
7	H	27	
8	I	7	

2 Entry composition [i](#)

There are 12 unique types of molecules in this entry. The entry contains 29180 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	231	Total	C	N	O	S	0	0	0
			1809	1155	315	337	2			
1	B	227	Total	C	N	O	S	0	0	0
			1789	1143	310	334	2			

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	1112	Total	C	N	O	S	0	0	0
			8774	5550	1565	1635	24			

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	D	1485	Total	C	N	O	S	0	1	0
			11739	7442	2069	2193	35			

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	E	94	Total	C	N	O	S	0	0	0
			758	483	132	139	4			

- Molecule 5 is a protein called DNA directed RNA polymerase sigma factor A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	F	346	Total	C	N	O	S	0	0	0
			2807	1770	509	524	4			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	-19	MET	-	expression tag	UNP Q5SKW1
F	-18	GLY	-	expression tag	UNP Q5SKW1
F	-17	SER	-	expression tag	UNP Q5SKW1
F	-16	SER	-	expression tag	UNP Q5SKW1
F	-15	HIS	-	expression tag	UNP Q5SKW1
F	-14	HIS	-	expression tag	UNP Q5SKW1
F	-13	HIS	-	expression tag	UNP Q5SKW1
F	-12	HIS	-	expression tag	UNP Q5SKW1
F	-11	HIS	-	expression tag	UNP Q5SKW1
F	-10	HIS	-	expression tag	UNP Q5SKW1
F	-9	SER	-	expression tag	UNP Q5SKW1
F	-8	SER	-	expression tag	UNP Q5SKW1
F	-7	GLY	-	expression tag	UNP Q5SKW1
F	-6	LEU	-	expression tag	UNP Q5SKW1
F	-5	VAL	-	expression tag	UNP Q5SKW1
F	-4	PRO	-	expression tag	UNP Q5SKW1
F	-3	ARG	-	expression tag	UNP Q5SKW1
F	-2	GLY	-	expression tag	UNP Q5SKW1
F	-1	SER	-	expression tag	UNP Q5SKW1
F	0	HIS	-	expression tag	UNP Q5SKW1

- Molecule 6 is a DNA chain called 5'-D(*CP*CP*T*GP*CP*AP*TP*CP*CP*GP*TP*GP*AP*GP*TP*CP*GP*AP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	G	16	Total	C	N	O	P	0	0	0
			328	156	63	94	15			

- Molecule 7 is a DNA chain called 5'-D(*TP*AP*TP*AP*AP*TP*GP*GP*GP*AP*GP*C P*TP*GP*TP*CP*AP*CP*GP*GP*AP*TP*GP*CP*AP*GP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	H	24	Total	C	N	O	P	0	0	0
			495	236	94	142	23			

- Molecule 8 is a protein called GE23077.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
8	I	7	Total	C	N	O	0	0	0
			50	26	9	15			


- Molecule 9 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	B	1	Total Mg 1 1	0	0
9	D	3	Total Mg 3 3	0	0
9	F	1	Total Mg 1 1	0	0

- Molecule 10 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	D	2	Total Zn 2 2	0	0

- Molecule 11 is (2Z)-2-methylbut-2-enoic acid (three-letter code: MB8) (formula: C₅H₈O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	I	1	Total C O 2 1 1	0	0


- Molecule 12 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
12	A	27	Total O 27 27	0	0
12	B	25	Total O 25 25	0	0



Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	C	221	Total 221	O 221	0	0
12	D	269	Total 269	O 269	0	0
12	E	22	Total 22	O 22	0	0
12	F	36	Total 36	O 36	0	0
12	G	10	Total 10	O 10	0	0
12	H	4	Total 4	O 4	0	0
12	I	8	Total 8	O 8	0	0



• Molecule 3: DNA-directed RNA polymerase subunit beta'






- Molecule 4: DNA-directed RNA polymerase subunit omega

Chain E: 85% 10% 5%




- Molecule 5: DNA directed RNA polymerase sigma factor A

Chain F: 3% 68% 10% 22%




- Molecule 6: 5'-D(*CP*CP*T*GP*CP*AP*TP*CP*CP*GP*TP*GP*AP*GP*TP*CP*GP*AP*G)-3'

Chain G: 16% 58% 21% 5% 16%



- Molecule 7: 5'-D(*TP*AP*TP*AP*AP*TP*GP*GP*GP*AP*GP*CP*TP*GP*TP*CP*AP*CP*GP*GP*AP*TP*GP*CP*AP*GP*G)-3'

Chain H: 11% 56% 33% 11%



● Molecule 8: GE23077

Chain I: 



4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	183.77Å 103.20Å 294.77Å 90.00° 99.18° 90.00°	Depositor
Resolution (Å)	48.50 – 2.80 48.84 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.0 (48.50-2.80) 98.1 (48.84-2.80)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.49 (at 2.81Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.206 , 0.252 0.207 , 0.252	Depositor DCC
R_{free} test set	6582 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	47.8	Xtrriage
Anisotropy	0.026	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 41.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.019 for $1/2^*h-3/2^*k,-1/2^*h-1/2^*k,-1/2^*h+1/2^*k-1$ 0.017 for $1/2^*h+3/2^*k,1/2^*h-1/2^*k,-1/2^*h-1/2^*k-1$	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	29180	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 2TL, ZN, MB8, MG, DVA, FGL, 0QZ, R2T, DSN, 2RA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/1841	0.46	0/2504
1	B	0.22	0/1821	0.44	0/2476
2	C	0.24	0/8941	0.45	0/12092
3	D	0.24	0/11948	0.46	0/16153
4	E	0.24	0/772	0.41	0/1040
5	F	0.23	0/2852	0.39	0/3837
6	G	0.47	0/368	1.00	1/567 (0.2%)
7	H	0.45	0/556	1.04	0/858
All	All	0.25	0/29099	0.48	1/39527 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
8	I	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	G	5	DC	O4'-C4'-C3'	-7.59	101.44	106.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
8	I	5	2TL	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1809	0	1863	28	0
1	B	1789	0	1841	23	0
2	C	8774	0	8877	109	0
3	D	11739	0	11977	165	0
4	E	758	0	770	7	0
5	F	2807	0	2882	35	0
6	G	328	0	181	5	0
7	H	495	0	272	9	0
8	I	50	0	37	3	0
9	B	1	0	0	0	0
9	D	3	0	0	0	0
9	F	1	0	0	0	0
10	D	2	0	0	0	0
11	I	2	0	0	0	0
12	A	27	0	0	0	0
12	B	25	0	0	0	0
12	C	221	0	0	5	0
12	D	269	0	0	7	0
12	E	22	0	0	0	0
12	F	36	0	0	0	0
12	G	10	0	0	1	0
12	H	4	0	0	2	0
12	I	8	0	0	0	0
All	All	29180	0	28700	342	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 342 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:802:ARG:HB2	2:C:826:TYR:HB2	1.62	0.81
3:D:1254:GLN:HB3	3:D:1258:ARG:HB2	1.66	0.75
2:C:428:ARG:NH2	2:C:447:ALA:O	2.20	0.75
2:C:409:ARG:HH11	2:C:454:SER:HB2	1.53	0.74

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:758:ARG:HH21	2:C:788:THR:HB	1.53	0.74

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	229/315 (73%)	226 (99%)	3 (1%)	0	100	100
1	B	225/315 (71%)	222 (99%)	3 (1%)	0	100	100
2	C	1108/1119 (99%)	1088 (98%)	20 (2%)	0	100	100
3	D	1482/1524 (97%)	1451 (98%)	31 (2%)	0	100	100
4	E	92/99 (93%)	89 (97%)	3 (3%)	0	100	100
5	F	344/443 (78%)	340 (99%)	4 (1%)	0	100	100
All	All	3480/3815 (91%)	3416 (98%)	64 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/273 (73%)	196 (98%)	4 (2%)	55	84
1	B	200/273 (73%)	196 (98%)	4 (2%)	55	84

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	C	936/941 (100%)	915 (98%)	21 (2%)	52	83
3	D	1253/1279 (98%)	1232 (98%)	21 (2%)	60	87
4	E	82/88 (93%)	82 (100%)	0	100	100
5	F	301/388 (78%)	297 (99%)	4 (1%)	69	91
All	All	2972/3242 (92%)	2918 (98%)	54 (2%)	59	86

5 of 54 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	1078	GLU
3	D	415	VAL
3	D	1488	ASP
3	D	67	ARG
3	D	276	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
3	D	1195	GLN
5	F	83	GLN
3	D	1441	GLN
3	D	560	GLN
3	D	1124	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

7 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	2RA	I	1	8,11	3,5,6	0.54	0	1,5,7	0.39	0
8	0QZ	I	6	8	4,5,6	1.65	1 (25%)	2,5,7	0.66	0
8	2TL	I	5	8	5,6,7	1.04	0	6,7,9	0.91	1 (16%)
8	R2T	I	4	8	8,10,11	1.88	2 (25%)	6,13,15	0.90	0
8	FGL	I	7	8	5,6,7	1.03	0	1,7,9	1.98	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	2RA	I	1	8,11	-	0/2/4/6	-
8	0QZ	I	6	8	-	1/3/4/6	-
8	2TL	I	5	8	-	1/5/6/8	-
8	R2T	I	4	8	-	7/13/14/16	-
8	FGL	I	7	8	-	0/4/6/8	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	I	4	R2T	CD-NE2	4.20	1.43	1.32
8	I	6	0QZ	OB-CA	-3.15	1.37	1.43
8	I	4	R2T	OB1-CB	-2.34	1.37	1.43

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	I	5	2TL	O-C-CA	-2.03	119.45	124.78

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	I	4	R2T	CA-CB-CG-OG1
8	I	4	R2T	OB1-CB-CG-CD
8	I	4	R2T	OB1-CB-CG-OG1
8	I	4	R2T	OE1-CD-CG-OG1
8	I	4	R2T	NE2-CD-CG-OG1

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	I	4	R2T	2	0
8	I	7	FGL	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 7 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
11	MB8	I	101	8	0,1,6	-	-	-		

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	231/315 (73%)	-0.12	6 (2%) 56 46	16, 33, 61, 102	0
1	B	227/315 (72%)	0.05	6 (2%) 56 46	21, 43, 78, 108	0
2	C	1112/1119 (99%)	-0.07	34 (3%) 49 39	4, 25, 84, 113	0
3	D	1485/1524 (97%)	0.17	105 (7%) 16 9	2, 33, 94, 121	0
4	E	94/99 (94%)	-0.29	0 100 100	7, 28, 63, 73	0
5	F	346/443 (78%)	0.12	13 (3%) 40 30	14, 47, 87, 113	0
6	G	16/19 (84%)	0.62	3 (18%) 1 1	44, 77, 145, 154	0
7	H	24/27 (88%)	0.34	3 (12%) 3 2	43, 83, 136, 156	0
8	I	0/7	-	-	-	-
All	All	3535/3868 (91%)	0.05	170 (4%) 30 21	2, 34, 89, 156	0

The worst 5 of 170 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	1499	ARG	5.6
3	D	191	LEU	5.4
2	C	207	LEU	5.1
3	D	422	ALA	4.9
2	C	365	ASP	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
8	2RA	I	1	6/7	0.92	0.17	8,10,16,18	0
8	FGL	I	7	7/8	0.97	0.13	8,9,10,11	0
8	DSN	I	2	6/7	0.97	0.15	7,9,15,15	0
8	2TL	I	5	7/8	0.97	0.17	6,7,8,9	0
8	R2T	I	4	11/12	0.98	0.15	7,8,9,11	0
8	DVA	I	3	7/8	0.98	0.17	8,9,13,13	0
8	0QZ	I	6	6/7	0.98	0.18	7,7,10,11	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
9	MG	B	2001	1/1	0.82	0.25	50,50,50,50	0
9	MG	D	2005	1/1	0.83	0.10	31,31,31,31	0
9	MG	F	2001	1/1	0.95	0.10	22,22,22,22	0
11	MB8	I	101	2/7	0.95	0.15	6,6,6,11	0
9	MG	D	2004	1/1	0.96	0.08	31,31,31,31	0
9	MG	D	2003	1/1	0.97	0.17	5,5,5,5	0
10	ZN	D	2002	1/1	0.98	0.05	56,56,56,56	0
10	ZN	D	2001	1/1	1.00	0.13	10,10,10,10	0

6.5 Other polymers [i](#)

There are no such residues in this entry.