



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 26, 2024 – 02:15 PM EDT

PDB ID : 6OI7  
Title : Se-Met structure of apo- Escherichia coli dGTPase  
Authors : Calero, G.; Barnes, C.O.; Wu, Y.  
Deposited on : 2019-04-08  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2335 (2.90-2.90)
Clashscore	180529	2564 (2.90-2.90)
Ramachandran outliers	177936	2514 (2.90-2.90)
Sidechain outliers	177891	2516 (2.90-2.90)
RSRZ outliers	164620	2337 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	505	 71% 23% 6% 0% 0%
1	B	505	 70% 22% 6% 2% 0%
1	C	505	 73% 22% 5% 0% 0%
1	D	505	 68% 25% 6% 2% 0%
1	E	505	 76% 21% 3% 0% 0%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	505	 75% 21%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	A	602	-	-	X	-
3	SO4	C	603	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 24823 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Deoxyguanosinetriphosphate triphosphohydrolase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	498	4135	2639	737	743	6	10	0	0	0
1	B	488	4050	2591	721	722	6	10	0	0	0
1	C	497	4125	2633	736	740	6	10	0	0	0
1	D	497	4132	2638	737	741	6	10	0	0	0
1	E	498	4139	2641	738	744	6	10	0	0	0
1	F	503	4178	2663	747	752	6	10	0	0	0

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Mn 1	0	0
2	B	1	Total 1	Mn 1	0	0
2	C	1	Total 1	Mn 1	0	0
2	D	1	Total 1	Mn 1	0	0
2	E	1	Total 1	Mn 1	0	0
2	F	1	Total 1	Mn 1	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0
3	E	1	Total O S 5 4 1	0	0
3	F	1	Total O S 5 4 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	5	Total O 5 5	0	0
4	C	3	Total O 3 3	0	0
4	D	1	Total O 1 1	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
4	E	2	Total O 2 2	0	0
4	F	2	Total O 2 2	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Deoxyguanosinetriphosphate triphosphohydrolase




- Molecule 1: Deoxyguanosinetriphosphate triphosphohydrolase




GLN


• Molecule 1: Deoxyguanosinetriphosphate triphosphohydrolase




• Molecule 1: Deoxyguanosinetriphosphate triphosphohydrolase



• Molecule 1: Deoxyguanosinetriphosphate triphosphohydrolase








● Molecule 1: Deoxyguanosinetriphosphate triphosphohydrolase

Chain F: 75% 21%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	192.18Å 192.18Å 287.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.97 – 2.90 49.97 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.97-2.90) 99.9 (49.97-2.90)	Depositor EDS
$R_{merge}$	0.28	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.53 (at 2.91Å)	Xtrriage
Refinement program	PHENIX (1.15.1_3469: ???)	Depositor
R, $R_{free}$	0.185 , 0.234 0.200 , 0.240	Depositor DCC
$R_{free}$ test set	3562 reflections (3.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	94.1	Xtrriage
Anisotropy	0.178	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 98.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	24823	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	110.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.42% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.57	4/4227 (0.1%)	0.75	4/5699 (0.1%)
1	B	0.41	0/4140	0.62	2/5581 (0.0%)
1	C	0.53	1/4217 (0.0%)	0.72	3/5685 (0.1%)
1	D	0.43	1/4224 (0.0%)	0.65	3/5695 (0.1%)
1	E	0.53	0/4231	0.70	1/5704 (0.0%)
1	F	0.52	1/4271 (0.0%)	0.72	3/5759 (0.1%)
All	All	0.50	7/25310 (0.0%)	0.69	16/34123 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	2
1	F	0	1
All	All	0	4

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	54	VAL	C-N	6.86	1.49	1.34
1	A	54	VAL	C-N	6.25	1.48	1.34
1	F	273	CYS	CB-SG	-6.01	1.72	1.82
1	A	415	ARG	CG-CD	5.61	1.66	1.51
1	D	54	VAL	C-N	5.32	1.46	1.34

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	150	LEU	CA-CB-CG	7.99	133.67	115.30
1	F	17	ARG	NE-CZ-NH1	6.91	123.75	120.30
1	B	57	LEU	CA-CB-CG	6.70	130.70	115.30
1	F	57	LEU	CA-CB-CG	6.39	130.00	115.30
1	C	433	ARG	CG-CD-NE	-6.33	98.51	111.80

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	167	GLU	Peptide
1	C	221	GLU	Peptide
1	C	58	GLU	Peptide
1	F	504	GLU	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4135	0	4091	96	1
1	B	4050	0	4011	83	0
1	C	4125	0	4081	85	0
1	D	4132	0	4086	115	1
1	E	4139	0	4094	72	0
1	F	4178	0	4135	98	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	5	0	0	2	0
3	B	5	0	0	0	0
3	C	20	0	0	4	0
3	D	5	0	0	0	0
3	E	5	0	0	0	0
3	F	5	0	0	0	0
4	A	5	0	0	1	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	3	0	0	3	0
4	D	1	0	0	0	0
4	E	2	0	0	3	0
4	F	2	0	0	1	0
All	All	24823	0	24498	514	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 514 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:347:TYR:CD1	1:D:375:CYS:SG	2.07	1.48
1:D:347:TYR:CG	1:D:375:CYS:SG	2.15	1.39
1:D:347:TYR:CE1	1:D:375:CYS:SG	2.44	1.10
1:D:57:LEU:HD12	1:D:57:LEU:H	1.19	1.06
1:F:498:ARG:HH21	1:F:503:VAL:HG12	1.24	1.01

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:148:GLN:NE2	1:D:374:GLU:OE1[3_544]	2.17	0.03

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	494/505 (98%)	458 (93%)	24 (5%)	12 (2%)	<b>5</b> <b>19</b>
1	B	482/505 (95%)	435 (90%)	33 (7%)	14 (3%)	<b>3</b> <b>15</b>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	493/505 (98%)	460 (93%)	24 (5%)	9 (2%)	7	25
1	D	493/505 (98%)	457 (93%)	29 (6%)	7 (1%)	9	31
1	E	494/505 (98%)	464 (94%)	22 (4%)	8 (2%)	8	28
1	F	501/505 (99%)	462 (92%)	29 (6%)	10 (2%)	6	23
All	All	2957/3030 (98%)	2736 (92%)	161 (5%)	60 (2%)	6	23

5 of 60 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	304	LYS
1	A	370	GLU
1	A	371	ASP
1	A	372	ALA
1	B	58	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	443/439 (101%)	412 (93%)	31 (7%)	12	36
1	B	433/439 (99%)	401 (93%)	32 (7%)	11	34
1	C	442/439 (101%)	413 (93%)	29 (7%)	14	39
1	D	442/439 (101%)	396 (90%)	46 (10%)	5	18
1	E	444/439 (101%)	417 (94%)	27 (6%)	15	43
1	F	449/439 (102%)	415 (92%)	34 (8%)	11	32
All	All	2653/2634 (101%)	2454 (92%)	199 (8%)	11	33

5 of 199 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	310	LEU
1	E	166	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	328	THR
1	D	415	ARG
1	E	273	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	296	HIS
1	C	302	HIS
1	D	251	ASN
1	D	365	ASN
1	E	60	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 6 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	D	602	-	4,4,4	0.26	0	6,6,6	0.18	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	F	602	-	4,4,4	0.20	0	6,6,6	0.11	0
3	SO4	A	602	-	4,4,4	0.21	0	6,6,6	0.12	0
3	SO4	E	602	-	4,4,4	0.23	0	6,6,6	0.15	0
3	SO4	B	602	-	4,4,4	0.25	0	6,6,6	0.20	0
3	SO4	C	605	-	4,4,4	0.30	0	6,6,6	0.25	0
3	SO4	C	603	-	4,4,4	0.26	0	6,6,6	0.43	0
3	SO4	C	602	-	4,4,4	0.26	0	6,6,6	0.57	0
3	SO4	C	604	-	4,4,4	0.27	0	6,6,6	0.37	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	602	SO4	2	0
3	C	605	SO4	1	0
3	C	603	SO4	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	488/505 (96%)	-0.58	4 (0%) 82 78	48, 87, 154, 220	0
1	B	478/505 (94%)	-0.39	5 (1%) 79 74	55, 132, 191, 225	0
1	C	487/505 (96%)	-0.61	4 (0%) 82 78	50, 88, 141, 198	0
1	D	487/505 (96%)	-0.29	8 (1%) 70 64	30, 136, 192, 225	0
1	E	488/505 (96%)	-0.60	6 (1%) 76 71	30, 91, 144, 200	0
1	F	493/505 (97%)	-0.57	2 (0%) 89 86	30, 93, 154, 231	0
All	All	2921/3030 (96%)	-0.51	29 (0%) 79 74	30, 102, 177, 231	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	505	GLN	7.3
1	D	373	SER	6.1
1	E	505	GLN	6.1
1	D	505	GLN	5.5
1	D	368(A)	LEU	4.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	A	602	5/5	0.06	0.29	426,444,448,450	0
3	SO4	F	602	5/5	0.72	0.07	168,172,177,178	0
3	SO4	D	602	5/5	0.77	0.07	169,176,178,181	0
3	SO4	B	602	5/5	0.79	0.07	165,166,169,171	0
3	SO4	C	603	5/5	0.85	0.08	136,146,150,150	0
3	SO4	E	602	5/5	0.86	0.07	142,143,146,158	0
2	MN	E	601	1/1	0.88	0.07	123,123,123,123	1
2	MN	F	601	1/1	0.89	0.07	101,101,101,101	1
3	SO4	C	605	5/5	0.90	0.12	109,120,129,137	0
3	SO4	C	602	5/5	0.92	0.07	84,90,129,133	0
2	MN	D	601	1/1	0.93	0.06	129,129,129,129	0
2	MN	C	601	1/1	0.94	0.05	102,102,102,102	1
2	MN	B	601	1/1	0.95	0.06	101,101,101,101	1
3	SO4	C	604	5/5	0.96	0.06	83,86,93,104	0
2	MN	A	601	1/1	0.97	0.04	114,114,114,114	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.