

wwPDB X-ray Structure Validation Summary Report (i)

Oct 3, 2023 – 04:50 AM EDT

PDB ID : 6OJB

Title : Crystal Structure of Aspergillus fumigatus Ega3 complex with galactosamine

Authors: Bamford, N.C.; Howell, P.L.

Deposited on : 2019-04-11

Resolution : 2.09 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : FAILED

Mogul : 1.8.5 (274361), CSD as 541be (2020)

Xtriage (Phenix) : 1.13

EDS : FAILED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.09 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 2372 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Endo alpha-1,4 polygalactosaminidase.

Mol	Chain	Residues		\mathbf{At}	oms			ZeroOcc	AltConf	Trace
1	۸	251	Total	С	N	О	S	0	0	0
1	A	201	1958	1222	332	393	11	0	0	

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	25	MET	-	initiating methionine	UNP Q4WX16
A	26	GLY	-	expression tag	UNP Q4WX16
A	27	SER	-	expression tag	UNP Q4WX16
A	28	SER	-	expression tag	UNP Q4WX16
A	29	HIS	-	expression tag	UNP Q4WX16
A	30	HIS	-	expression tag	UNP Q4WX16
A	31	HIS	-	expression tag	UNP Q4WX16
A	32	HIS	-	expression tag	UNP Q4WX16
A	33	HIS	-	expression tag	UNP Q4WX16
A	34	HIS	-	expression tag	UNP Q4WX16
A	35	SER	-	expression tag	UNP Q4WX16
A	36	SER	-	expression tag	UNP Q4WX16
A	37	GLY	-	expression tag	UNP Q4WX16
A	38	LEU	-	expression tag	UNP Q4WX16
A	39	VAL	-	expression tag	UNP Q4WX16
A	40	PRO	-	expression tag	UNP Q4WX16
A	41	ARG	-	expression tag	UNP Q4WX16
A	42	GLY	-	expression tag	UNP Q4WX16
A	43	SER	-	expression tag	UNP Q4WX16
A	44	LEU		expression tag	UNP Q4WX16
A	45	MET	-	expression tag	UNP Q4WX16

• Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	В	8	Total 94	C 52	N 2	O 40	0	0	0


• Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyran ose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-3)-beta-D-mannopyran ose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	С	7	Total 83			O 35	0	0	0

• Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	2	Total 28	C 16	N 2	O 10	0	0	0


• Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf	
5	А	1	Total	С	N	О	0	0	
	11	1	14	8	1	5	U		
5	Λ	1	Total	С	N	О	0	0	
9	A	1	14	8	1	5	0	U	



• Molecule 6 is 2-amino-2-deoxy-alpha-D-galactopyranose (three-letter code: X6X) (formula: $C_6H_{13}NO_5$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C N O 12 6 1 5	0	0

• Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	169	Total O 169 169	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	44.89Å 48.17Å 163.44Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.13 - 2.09	Depositor
% Data completeness	99.8 (28.13-2.09)	Depositor
(in resolution range)	, , ,	•
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.79 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.168 , 0.210	Depositor
Wilson B-factor (\mathring{A}^2)	30.6	Xtriage
Anisotropy	0.602	Xtriage
L-test for twinning ²	$ < L > = 0.50, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2372	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 9.04% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

17 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Во	ond leng	ths	В	ond ang	les
IVIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	В	1	2,1	14,14,15	0.78	1 (7%)	17,19,21	0.59	0
2	NAG	В	2	2	14,14,15	0.45	0	17,19,21	0.59	0
2	BMA	В	3	2	11,11,12	0.84	0	15,15,17	0.96	1 (6%)
2	MAN	В	4	2	11,11,12	1.09	1 (9%)	15,15,17	1.67	2 (13%)
2	MAN	В	5	2	11,11,12	0.79	0	15,15,17	1.05	2 (13%)
2	MAN	В	6	2	11,11,12	0.90	0	15,15,17	1.07	1 (6%)
2	MAN	В	7	2	11,11,12	1.64	2 (18%)	15,15,17	1.43	3 (20%)
2	MAN	В	8	2	11,11,12	0.99	0	15,15,17	1.08	2 (13%)
3	NAG	С	1	1,3	14,14,15	0.42	0	17,19,21	0.93	1 (5%)
3	NAG	С	2	3	14,14,15	0.53	0	17,19,21	0.86	0
3	BMA	С	3	3	11,11,12	1.92	3 (27%)	15,15,17	1.14	1 (6%)
3	MAN	С	4	3	11,11,12	1.31	2 (18%)	15,15,17	1.41	2 (13%)
3	MAN	С	5	3	11,11,12	0.75	0	15,15,17	1.20	2 (13%)
3	MAN	С	6	3	11,11,12	1.35	2 (18%)	15,15,17	1.01	1 (6%)
3	MAN	С	7	3	11,11,12	1.37	2 (18%)	15,15,17	1.43	3 (20%)
4	NAG	D	1	1,4	14,14,15	0.44	0	17,19,21	0.57	0
4	NAG	D	2	4	14,14,15	0.27	0	17,19,21	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	В	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	В	2	2	-	2/6/23/26	0/1/1/1
2	BMA	В	3	2	-	2/2/19/22	0/1/1/1
2	MAN	В	4	2	-	2/2/19/22	0/1/1/1
2	MAN	В	5	2	-	2/2/19/22	0/1/1/1
2	MAN	В	6	2	-	0/2/19/22	1/1/1/1
2	MAN	В	7	2	-	1/2/19/22	0/1/1/1
2	MAN	В	8	2	-	0/2/19/22	0/1/1/1
3	NAG	С	1	1,3	-	3/6/23/26	0/1/1/1
3	NAG	С	2	3	-	2/6/23/26	0/1/1/1
3	BMA	С	3	3	-	2/2/19/22	0/1/1/1
3	MAN	С	4	3	-	0/2/19/22	0/1/1/1
3	MAN	С	5	3	-	2/2/19/22	0/1/1/1

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAN	С	6	3	-	0/2/19/22	0/1/1/1
3	MAN	С	7	3	-	0/2/19/22	0/1/1/1
4	NAG	D	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	D	2	4	-	2/6/23/26	0/1/1/1

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(\text{\AA})$
3	С	3	BMA	C1-C2	5.12	1.63	1.52
2	В	7	MAN	C2-C3	3.89	1.58	1.52
3	С	7	MAN	C1-C2	3.15	1.59	1.52
2	В	4	MAN	O5-C5	2.83	1.49	1.43
3	С	4	MAN	C1-C2	2.70	1.58	1.52

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}(^{o})$
2	В	4	MAN	C1-O5-C5	5.40	119.52	112.19
3	С	7	MAN	C1-O5-C5	3.83	117.39	112.19
3	С	4	MAN	C1-C2-C3	3.73	114.25	109.67
3	С	5	MAN	C1-O5-C5	3.47	116.89	112.19
2	В	7	MAN	C1-C2-C3	3.17	113.56	109.67

There are no chirality outliers.

5 of 22 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	С	5	MAN	C4-C5-C6-O6
3	С	5	MAN	O5-C5-C6-O6
3	С	3	BMA	O5-C5-C6-O6
2	В	4	MAN	O5-C5-C6-O6
3	С	1	NAG	C4-C5-C6-O6


All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	6	MAN	C1-C2-C3-C4-C5-O5


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.














4.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Res	Link	Во	ond leng	ths	Bond angles		
IVIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	A	418	1	14,14,15	0.24	0	17,19,21	0.56	0
5	NAG	A	419	1	14,14,15	0.49	0	17,19,21	0.43	0
6	X6X	A	420	-	12,12,12	1.42	1 (8%)	16,17,17	1.45	5 (31%)



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	418	1	-	2/6/23/26	0/1/1/1
5	NAG	A	419	1	-	0/6/23/26	0/1/1/1
6	X6X	A	420	-	-	1/2/22/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	Ideal(A)
6	A	420	X6X	C1-C2	2.50	1.55	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\mathrm{Ideal}(^{o})$
6	A	420	X6X	O5-C1-C2	2.75	112.72	109.51
6	A	420	X6X	C1-C2-C3	-2.34	107.53	110.60
6	A	420	X6X	O3-C3-C2	-2.12	106.40	110.22
6	A	420	X6X	O6-C6-C5	-2.08	104.16	111.29
6	A	420	X6X	C3-C2-N2	-2.05	106.85	111.05

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	418	NAG	O5-C5-C6-O6
5	A	418	NAG	C4-C5-C6-O6
6	A	420	X6X	O5-C5-C6-O6


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and



any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

