

wwPDB X-ray Structure Validation Summary Report (i)

Jan 7, 2024 – 02:08 pm GMT

PDB ID	:	50LZ
Title	:	Structure of the A2A-StaR2-bRIL562-Compound 4e complex at 1.9A obtained
		from bespoke co-crystallisation experiments.
Authors	:	Rucktooa, P.; Cheng, R.K.Y.; Segala, E.; Geng, T.; Errey, J.C.; Brown, G.A.;
		Cooke, R.; Marshall, F.H.; Dore, A.S.
Deposited on		
Resolution	:	1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as 541 be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
			13%						
1	А	434	80%	9%	11%				

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	PGE	А	1233	-	-	-	Х



2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 3809 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Adenosine receptor A2a,Soluble cytochrome b562,Adenosine receptor A2a.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	387	Total 3104	C 2024	N 525	O 532	S 23	0	14	0

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-9	ALA	-	expression tag	UNP P29274
А	-8	ASP	- expression tag		UNP P29274
А	-7	TYR	-	expression tag	UNP P29274
А	-6	LYS	-	expression tag	UNP P29274
А	-5	ASP	-	expression tag	UNP P29274
А	-4	ASP	-	expression tag	UNP P29274
А	-3	ASP	-	expression tag	UNP P29274
А	-2	ASP	-	expression tag	UNP P29274
А	-1	GLY	-	expression tag	UNP P29274
А	0	ALA	-	expression tag	UNP P29274
А	1	PRO	-	expression tag	UNP P29274
А	54	LEU	ALA	engineered mutation	UNP P29274
A	88	ALA	THR	engineered mutation	UNP P29274
А	107	ALA	ARG	engineered mutation	UNP P29274
А	122	ALA	LYS	engineered mutation	UNP P29274
А	154	ALA	ASN	engineered mutation	UNP P29274
А	202	ALA	LEU	engineered mutation	UNP P29274
А	1007	TRP	MET	engineered mutation	UNP P0ABE7
А	1102	ILE	HIS	engineered mutation	UNP P0ABE7
А	1106	LEU	ARG	engineered mutation	UNP P0ABE7
А	235	ALA	LEU	engineered mutation	UNP P29274
А	239	ALA	VAL	engineered mutation	UNP P29274
А	277	ALA	SER	engineered mutation	UNP P29274
А	318	ALA	-	expression tag	UNP P29274
А	319	HIS	-	expression tag	UNP P29274
А	320	HIS	-	expression tag	UNP P29274


Continued on next page...



Chain	Residue	Modelled	Actual	Comment	Reference
А	321	HIS	-	expression tag	UNP P29274
А	322	HIS	-	expression tag	UNP P29274
А	323	HIS	-	expression tag	UNP P29274
А	324	HIS	-	expression tag	UNP P29274
А	325	HIS	-	expression tag	UNP P29274
А	326	HIS	-	expression tag	UNP P29274
А	327	HIS	-	expression tag	UNP P29274
A	328	HIS	-	expression tag	UNP P29274

Continued from previous page...


• Molecule 2 is 4-(3-amino-5-phenyl-1,2,4-triazin-6-yl)-2-chlorophenol (three-letter code: T4E) (formula: $C_{15}H_{11}ClN_4O$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	А	1	Total	С	Cl	Ν	Ο	0	0
	11	Ĩ	21	15	1	4	1	Ŭ	Ŭ


• Molecule 3 is CHOLESTEROL (three-letter code: CLR) (formula: $C_{27}H_{46}O$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total C O 28 27 1	0	0
3	А	1	Total C O 28 27 1	0	0
3	А	1	Total C O 28 27 1	0	0
3	А	1	Total C O 28 27 1	0	0

• Molecule 4 is OLEIC ACID (three-letter code: OLA) (formula: $C_{18}H_{34}O_2$).






50LZ

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	Total C O 20 18 2	0	0
4	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 9 & 7 & 2 \end{array}$	0	0
4	А	1	Total C O 20 18 2	0	0
4	А	1	Total C O 18 16 2	0	0
4	А	1	Total C O 15 13 2	0	0
4	А	1	Total C O 12 10 2	0	0
4	А	1	Total C O 12 10 2	0	0
4	А	1	Total C 10 10	0	0
4	А	1	Total C 10 10	0	0
4	А	1	Total C 9 9	0	0
4	А	1	Total C O 19 17 2	0	0
4	А	1	Total C 12 12	0	0
4	А	1	Total C 9 9	0	0
4	А	1	Total C O 16 14 2	0	0
4	А	1	Total C O 20 18 2	0	0

• Molecule 5 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula: $C_{21}H_{40}O_4$).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	Total C O 17 13 4	0	0
5	А	1	Total C O 19 15 4	0	0
5	А	1	Total C O 16 12 4	0	0
5	А	1	Total C O 22 18 4	0	0
5	А	1	Total C O 25 21 4	0	0
5	А	1	Total C O 25 21 4	0	0
5	А	1	Total C O 24 20 4	0	0
5	А	1	Total C O 18 14 4	0	0
5	А	1	Total C O 25 21 4	0	0
5	А	1	Total C O 17 13 4	0	0
5	А	1	Total C 14 14	0	0


• Molecule 6 is D(-)-TARTARIC ACID (three-letter code: TAR) (formula: $C_4H_6O_6$).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	А	1	Total 10	$\begin{array}{c} \mathrm{C} \\ 4 \end{array}$	O 6	0	0

• Molecule 7 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
7	А	1	Total 10	C 6	0 4	0	0

• Molecule 8 is SODIUM ION (three-letter code: NA) (formula: Na).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	А	1	Total Na 1 1	0	0

• Molecule 9 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	А	118	Total O 118 118	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Adenosine receptor A2a,Soluble cytochrome b562,Adenosine receptor A2a





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants	39.37Å 179.25Å 140.07Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.71 - 1.90	Depositor
Resolution (A)	33.71 - 1.90	EDS
% Data completeness	99.3 (33.71-1.90)	Depositor
(in resolution range)	99.1 (33.71-1.90)	EDS
R _{merge}	0.08	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.52 (at 1.89 \text{\AA})$	Xtriage
Refinement program	PHENIX (1.12rc2_2821: ???)	Depositor
D D.	0.173 , 0.196	Depositor
R, R_{free}	0.172 , 0.193	DCC
R_{free} test set	1954 reflections $(4.96%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	30.3	Xtriage
Anisotropy	0.045	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.37,77.1	EDS
L-test for twinning ²	$ \langle L \rangle = 0.47, \langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3809	wwPDB-VP
Average B, all atoms $(Å^2)$	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.78% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, TAR, T4E, NA, CLR, OLC, OLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
	Unain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.28	0/3173	0.47	3/4315~(0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	208	LEU	O-C-N	-12.16	103.24	122.70
1	А	208	LEU	CA-C-N	6.76	132.08	117.20
1	А	208	LEU	C-N-CA	5.88	136.41	121.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	208	LEU	Mainchain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3104	0	3181	35	0
2	А	21	0	10	0	0
3	А	112	0	184	6	0
4	А	211	0	315	20	0
5	А	222	0	321	26	0
6	А	10	0	4	0	0
7	А	10	0	14	1	0
8	А	1	0	0	0	0
9	А	118	0	0	2	0
All	All	3809	0	4029	55	0

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 55 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1203:CLR:C27	5:A:1225:OLC:H15	1.92	0.99
3:A:1203:CLR:H272	5:A:1225:OLC:H15	1.47	0.93
3:A:1203:CLR:C27	5:A:1225:OLC:C15	2.49	0.90
1:A:71:CYS:O	5:A:1225:OLC:H24A	1.72	0.89
4:A:1208:OLA:H151	5:A:1224:OLC:H12	1.62	0.80

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Analysed Favoured		Outliers	Percentiles	
1	А	397/434~(92%)	395~(100%)	2~(0%)	0	100 100	

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	325/353~(92%)	325 (100%)	0	100 100	

There are no protein residues with a non-rotameric sidechain to report.

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. There are no such side chains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 34 ligands modelled in this entry, 1 is monoatomic - leaving 33 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type	Chain	Res	Res Link	Bond lengths			Bond angles			
				Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2	
4	OLA	А	1211	-	11,11,19	0.81	0	$11,\!11,\!19$	1.02	1 (9%)



N T 1	m		Ъ	T • 1	Bo	ond leng	ths	В	ond ang	les
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	OLC	А	1226	-	23,23,24	0.92	1 (4%)	24,24,25	1.09	2 (8%)
4	OLA	А	1214	-	9,9,19	0.60	0	8,8,19	0.25	0
5	OLC	А	1221	-	18,18,24	0.98	1(5%)	18,19,25	1.26	2 (11%)
4	OLA	А	1219	-	$15,\!15,\!19$	0.69	0	15,15,19	0.83	1 (6%)
4	OLA	А	1208	-	19,19,19	0.68	0	19,19,19	0.70	1 (5%)
4	OLA	А	1218	-	8,8,19	0.45	0	7,7,19	0.20	0
3	CLR	А	1204	-	31,31,31	0.76	0	48,48,48	0.84	0
5	OLC	А	1220	-	16, 16, 24	1.06	1(6%)	17,17,25	1.40	2 (11%)
4	OLA	А	1206	-	19,19,19	0.66	0	19,19,19	0.71	1 (5%)
4	OLA	А	1210	-	14,14,19	0.71	0	14,14,19	0.86	1 (7%)
5	OLC	А	1224	-	24,24,24	0.90	1 (4%)	25,25,25	0.98	2 (8%)
7	PGE	А	1233	-	$9,\!9,\!9$	0.30	0	8,8,8	0.31	0
4	OLA	А	1216	-	18,18,19	0.68	0	18,18,19	0.82	1 (5%)
3	CLR	А	1205	-	31,31,31	0.71	0	48,48,48	0.88	0
6	TAR	А	1231	-	$9,\!9,\!9$	0.96	0	12,12,12	1.58	2 (16%)
4	OLA	А	1212	-	11,11,19	0.82	0	11,11,19	1.04	1 (9%)
4	OLA	А	1232	-	19,19,19	0.68	0	19,19,19	0.75	1 (5%)
3	CLR	А	1202	-	31,31,31	0.67	0	48,48,48	1.11	5 (10%)
4	OLA	А	1217	-	11,11,19	0.39	0	9,10,19	0.35	0
3	CLR	А	1203	-	31,31,31	0.71	0	48,48,48	1.08	3 (6%)
5	OLC	А	1227	-	17,17,24	1.01	1 (5%)	18,18,25	1.26	2 (11%)
5	OLC	А	1229	-	16,16,24	1.07	1 (6%)	17,17,25	1.24	2 (11%)
5	OLC	А	1225	-	24,24,24	0.89	1 (4%)	25,25,25	1.28	2 (8%)
2	T4E	А	1201	-	23,23,23	2.06	8 (34%)	32,32,32	2.94	9 (28%)
5	OLC	А	1230	-	13,13,24	0.23	0	12,12,25	0.45	0
4	OLA	А	1215	-	8,8,19	0.48	0	7,7,19	0.54	0
4	OLA	А	1209	-	17,17,19	0.53	0	17,17,19	0.90	2 (11%)
4	OLA	А	1213	-	9,9,19	0.44	0	7,8,19	0.34	0
4	OLA	А	1207	-	8,8,19	0.85	0	8,8,19	1.17	2 (25%)
5	OLC	А	1223	-	21,21,24	0.80	1 (4%)	22,22,25	0.95	2 (9%)
5	OLC	А	1222	-	15,15,24	1.09	1 (6%)	16,16,25	1.08	1 (6%)
5	OLC	А	1228	-	24,24,24	0.87	1 (4%)	25,25,25	0.95	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	OLA	А	1211	-	-	7/9/9/17	-
5	OLC	А	1226	-	-	6/23/23/24	-
4	OLA	А	1214	-	-	2/7/7/17	-
5	OLC	А	1221	-	-	12/18/18/24	-
4	OLA	А	1219	-	-	9/13/13/17	-
4	OLA	А	1208	-	-	5/17/17/17	-
4	OLA	А	1218	_	-	5/6/6/17	_
3	CLR	А	1204	-	-	0/10/68/68	0/4/4/4
5	OLC	А	1220	-	-	4/16/16/24	-
4	OLA	А	1206	-	-	10/17/17/17	-
4	OLA	А	1210	-	-	2/12/12/17	-
5	OLC	А	1224	-	-	8/24/24/24	-
7	PGE	А	1233	-	-	2/7/7/7	-
4	OLA	А	1216	-	-	5/16/16/17	-
3	CLR	А	1205	-	-	2/10/68/68	0/4/4/4
6	TAR	А	1231	-	-	8/12/12/12	-
4	OLA	А	1212	-	-	7/9/9/17	_
4	OLA	А	1232	_	-	9/17/17/17	-
3	CLR	А	1202	_	-	5/10/68/68	0/4/4/4
4	OLA	А	1217	-	-	5/9/9/17	_
3	CLR	А	1203	-	-	0/10/68/68	0/4/4/4
5	OLC	А	1227	-	-	4/17/17/24	-
5	OLC	А	1229	-	-	7/16/16/24	-
5	OLC	А	1225	-	-	7/24/24/24	-
2	T4E	А	1201	-	-	0/8/8/8	0/3/3/3
5	OLC	А	1230	-	-	8/11/11/24	-
4	OLA	А	1215	-	-	2/6/6/17	-
4	OLA	А	1209	-	-	9/15/15/17	-
4	OLA	А	1213	-	-	5/7/7/17	-
4	OLA	А	1207	-	-	1/6/6/17	-
5	OLC	А	1223	-	-	6/21/21/24	-
5	OLC	А	1222	-	-	9/15/15/24	-
5	OLC	А	1228	-	-	8/24/24/24	-

The worst 5 of 18 bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\mathrm{Ideal}(\mathrm{\AA})$
2	А	1201	T4E	C3-N2	4.90	1.40	1.35
5	А	1224	OLC	O20-C1	4.05	1.45	1.33
5	А	1226	OLC	O20-C1	4.00	1.45	1.33
5	А	1228	OLC	O20-C1	3.99	1.45	1.33
5	А	1229	OLC	O20-C1	3.93	1.44	1.33

The worst 5 of 50 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms		$Observed(^{o})$	$Ideal(^{o})$
2	А	1201	T4E	N4-C3-N2	-9.13	119.33	125.57
2	А	1201	T4E	C5-C6-N1	-8.15	115.04	119.87
2	А	1201	T4E	C3-N2-N1	5.00	121.01	117.38
2	А	1201	T4E	C8-C9-C10	4.49	123.52	120.91
2	А	1201	T4E	C7-C6-N1	4.44	120.36	114.44

There are no chirality outliers.

5 of 179 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	А	1217	OLA	C9-C10-C11-C12
5	А	1222	OLC	O20-C21-C22-C24
5	А	1224	OLC	O20-C21-C22-C24
6	А	1231	TAR	O3-C3-C4-O4
6	А	1231	TAR	O3-C3-C4-O41

There are no ring outliers.

22 monomers are involved in 41 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	А	1226	OLC	3	0
4	А	1214	OLA	2	0
5	А	1221	OLC	1	0
4	А	1219	OLA	1	0
4	А	1208	OLA	8	0
4	А	1218	OLA	1	0
5	А	1220	OLC	1	0
5	А	1224	OLC	6	0
7	А	1233	PGE	1	0
4	А	1216	OLA	3	0
4	А	1212	OLA	1	0
3	А	1202	CLR	1	0
4	А	1217	OLA	3	0


Continued on next page...




Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	А	1203	CLR	5	0
5	А	1229	OLC	2	0
5	А	1225	OLC	8	0
4	А	1215	OLA	1	0
4	А	1209	OLA	1	0
4	А	1213	OLA	1	0
5	А	1223	OLC	1	0
5	А	1222	OLC	3	0
5	А	1228	OLC	3	0

Continued from previous page...


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


















































5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	387/434~(89%)	0.67	55 (14%) 2 2	18, 37, 86, 122	0

The worst 5 of 55 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	-1	GLY	9.2
1	А	1061	PHE	7.4
1	А	290	TYR	7.0
1	А	1017	ILE	6.1
1	А	1059	LYS	6.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	\mathbf{RSR}	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
7	PGE	А	1233	10/10	0.34	0.56	99,103,109,109	0
5	OLC	А	1224	25/25	0.56	0.24	53,70,91,93	0

Continued on next page...




50LZ

Mol	Type	Chain	Res	Atoms	RSCC	\mathbf{RSR}	B-factors(Å ²)	Q<0.9
4	OLA	А	1211	12/20	0.62	0.24	$55,\!66,\!81,\!82$	0
4	OLA	А	1208	20/20	0.65	0.21	46,70,76,76	0
5	OLC	А	1220	17/25	0.67	0.24	71,76,82,83	0
5	OLC	А	1228	25/25	0.71	0.22	63,76,96,98	0
4	OLA	А	1212	12/20	0.71	0.18	65,68,70,71	0
4	OLA	А	1210	15/20	0.72	0.20	68,72,85,86	0
3	CLR	А	1202	28/28	0.73	0.35	69,76,78,79	0
5	OLC	А	1229	17/25	0.74	0.28	58,71,88,89	0
5	OLC	А	1223	22/25	0.74	0.21	48,55,82,84	0
4	OLA	А	1213	10/20	0.75	0.20	67,69,71,71	0
4	OLA	А	1214	10/20	0.76	0.20	$50,\!55,\!63,\!63$	0
5	OLC	А	1221	19/25	0.80	0.22	32,55,68,69	0
4	OLA	А	1232	20/20	0.80	0.21	64,65,69,69	20
6	TAR	А	1231	10/10	0.80	0.29	92,92,92,92	10
4	OLA	А	1216	19/20	0.80	0.24	55,67,77,77	0
5	OLC	А	1226	24/25	0.81	0.20	43,64,76,78	0
5	OLC	А	1227	18/25	0.81	0.20	52,60,75,76	0
5	OLC	А	1222	16/25	0.82	0.19	47,58,65,67	0
4	OLA	А	1218	9/20	0.82	0.20	47,59,65,66	0
4	OLA	А	1217	12/20	0.82	0.14	61,63,67,67	0
4	OLA	А	1209	18/20	0.84	0.18	54,63,75,75	0
4	OLA	А	1207	9/20	0.85	0.17	44,47,61,63	0
4	OLA	А	1206	20/20	0.88	0.15	40,57,69,69	0
4	OLA	А	1219	16/20	0.88	0.21	61,71,74,75	0
5	OLC	А	1225	25/25	0.88	0.17	40,56,80,82	0
5	OLC	А	1230	14/25	0.90	0.12	48,56,61,63	0
4	OLA	А	1215	9/20	0.93	0.18	45,48,57,59	0
3	CLR	А	1204	28/28	0.95	0.09	28,33,41,45	0
3	CLR	А	1203	28/28	0.96	0.08	26,29,56,61	0
3	CLR	А	1205	28/28	0.96	0.09	23,29,57,58	0
8	NA	А	1234	1/1	0.97	0.13	40,40,40,40	0
2	T4E	А	1201	21/21	0.98	0.23	17,19,21,27	0


Continued from previous page...

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.















































































6.5 Other polymers (i)

There are no such residues in this entry.

