



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 29, 2024 – 03:03 PM EDT

PDB ID : 4OPD
Title : Constructing tailored isoprenoid products by structure-guided modification of geranylgeranyl reductase.
Authors : McAndrew, R.P.; Kung, Y.; Xie, X.; Liu, C.; Pereira, J.H.; Keasling, J.D.; Adams, P.D.
Deposited on : 2014-02-05
Resolution : 1.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	9242 (1.84-1.80)
Clashscore	180529	1080 (1.82-1.82)
Ramachandran outliers	177936	1073 (1.82-1.82)
Sidechain outliers	177891	1073 (1.82-1.82)
RSRZ outliers	164620	9241 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	453	 90% 10%
1	B	453	 91% 8%

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 8334 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Conserved Archaeal protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	452	Total 3649	C 2339	N 609	O 686	S 15	0	15	0
1	B	452	Total 3638	C 2331	N 608	O 684	S 15	0	13	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	HIS	-	expression tag	UNP Q4JA33
B	0	HIS	-	expression tag	UNP Q4JA33

- Molecule 2 is DIHYDROFLAVINE-ADENINE DINUCLEOTIDE (three-letter code: FDA) (formula: $C_{27}H_{35}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total 53	C 27	N 9	O 15	P 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	B	1	53	27	9	15	2	0	0

- Molecule 3 is GERANYLGERANYL DIPHOSPHATE (three-letter code: GRG) (formula: $C_{20}H_{36}O_7P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
3	A	1	29	20	7	2	0	0
3	A	1	25	20	4	1	0	0
3	A	1	21	20	1		0	0
3	B	1	29	20	7	2	0	0
3	B	1	25	20	4	1	0	0
3	B	1	21	20	1		0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	389	Total 389 389	0	0
4	B	402	Total 402 402	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Conserved Archaeal protein



- Molecule 1: Conserved Archaeal protein



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	63.08Å 63.23Å 65.10Å 120.98° 89.97° 88.55°	Depositor
Resolution (Å)	42.10 – 1.81 42.10 – 1.81	Depositor EDS
% Data completeness (in resolution range)	90.7 (42.10-1.81) 91.3 (42.10-1.81)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.19 (at 1.81Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1525)	Depositor
R, R_{free}	0.146 , 0.191 0.149 , 0.190	Depositor DCC
R_{free} test set	1925 reflections (2.68%)	wwPDB-VP
Wilson B-factor (Å ²)	16.9	Xtriage
Anisotropy	0.421	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 36.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.000 for h,k+1,-k 0.000 for h,-l,k+1 0.000 for h,l,-k-l 0.000 for h,-k-l,k 0.009 for h,-k,-l 0.010 for -h,k,-k-l 0.013 for -h,l,k 0.020 for -h,k+1,-l 0.397 for -h,-k-l,l 0.000 for -h,-k,k+1 0.000 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	8334	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GRG, FDA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/3764	0.53	0/5076
1	B	0.35	0/3747	0.52	0/5054
All	All	0.35	0/7511	0.52	0/10130

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3649	0	3683	24	0
1	B	3638	0	3666	23	0
2	A	53	0	33	4	0
2	B	53	0	33	3	0
3	A	75	0	99	6	0
3	B	75	0	99	7	0
4	A	389	0	0	3	0
4	B	402	0	0	1	0
All	All	8334	0	7613	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:381:GLU:HG3	1:B:382:LYS:HG3	1.70	0.74
3:B:502:GRG:H111	3:B:503:GRG:HC42	1.78	0.66
1:B:350:PHE:CZ	3:B:504:GRG:HC91	2.34	0.62
1:A:126:ILE:HD11	1:A:147:THR:HG23	1.82	0.61
1:B:126:ILE:HD11	1:B:147:THR:HG23	1.82	0.61
1:A:140[B]:ARG:NH2	1:B:58:LYS:O	2.36	0.59
1:A:159:TYR:CE2	1:A:268:LEU:HD12	2.37	0.58
1:A:189:VAL:HB	1:A:263:VAL:HG12	1.86	0.58
1:A:191:LEU:HD12	1:A:260[A]:LYS:HD3	1.87	0.56
1:B:68:GLU:OE2	1:B:104:ARG:HD2	2.07	0.54
1:B:159:TYR:CE2	1:B:268:LEU:HD12	2.42	0.53
1:A:78:LEU:HD11	1:A:206:ILE:HD11	1.90	0.53
3:A:502:GRG:H111	3:A:503:GRG:HC42	1.91	0.52
1:B:378:GLU:O	1:B:381:GLU:HG2	2.10	0.51
1:A:68:GLU:OE2	1:A:104:ARG:HD2	2.12	0.49
1:A:300:GLY:H	2:A:501:FDA:HN1	1.60	0.48
1:B:426:TYR:CG	1:B:427:PRO:HD2	2.49	0.48
1:B:78:LEU:HD11	1:B:206:ILE:HD11	1.95	0.47
2:B:501:FDA:H2'	2:B:501:FDA:HN1	1.80	0.47
3:B:503:GRG:H101	3:B:503:GRG:HC62	1.64	0.47
1:A:420:LYS:O	1:A:424:LEU:HG	2.14	0.46
2:A:501:FDA:HN1	2:A:501:FDA:H2'	1.80	0.46
1:A:259:SER:O	1:A:260[B]:LYS:HD3	2.15	0.46
1:B:64:PRO:HG2	1:B:96:LEU:HD21	1.98	0.46
1:B:300:GLY:H	2:B:501:FDA:HN1	1.64	0.45
3:A:502:GRG:HC2	3:A:502:GRG:HC62	1.65	0.45
1:B:117:TRP:CZ3	1:B:146:LEU:HD13	2.51	0.45
1:A:346[B]:SER:OG	1:A:420:LYS:HD3	2.16	0.45
1:A:427:PRO:HB3	1:A:432:SER:HB2	1.98	0.44
3:A:503:GRG:H101	3:A:503:GRG:HC62	1.70	0.44
2:A:501:FDA:C5X	3:A:502:GRG:H192	2.46	0.44
1:A:107:LYS:HD2	4:A:831:HOH:O	2.17	0.44
1:B:206:ILE:HD12	3:B:503:GRG:H17	1.99	0.44
1:B:98:ALA:HB3	1:B:99:PRO:HD3	1.99	0.44
1:A:98:ALA:HB3	1:A:99:PRO:HD3	2.00	0.44
1:B:175:GLU:HB3	1:B:275:LEU:HD22	1.99	0.43
1:A:110:GLN:HA	1:A:114:VAL:O	2.17	0.43
1:B:375:ASP:OD2	4:B:715:HOH:O	2.21	0.43
1:A:64:PRO:HG2	1:A:96:LEU:HD21	1.99	0.43
3:B:502:GRG:H101	3:B:502:GRG:HC61	1.62	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:502:GRG:H101	3:A:502:GRG:HC61	1.57	0.42
3:A:504:GRG:HC11	3:A:504:GRG:HC41	1.73	0.42
1:B:189:VAL:HB	1:B:263:VAL:HG12	2.00	0.42
1:A:250:ASP:OD1	4:A:639:HOH:O	2.21	0.42
3:B:502:GRG:HC62	3:B:502:GRG:HC2	1.69	0.42
1:B:203:ARG:HB2	1:B:218:TYR:HB3	2.02	0.42
1:B:259:SER:O	1:B:260:LYS:HD3	2.19	0.42
1:A:428:ARG:NH1	4:A:708:HOH:O	2.45	0.41
1:B:394:MET:CE	1:B:409:LEU:HB3	2.51	0.41
1:A:47:CYS:HA	2:A:501:FDA:C6	2.50	0.41
1:A:387:LEU:HD23	1:A:388:SER:N	2.36	0.41
1:B:53:LYS:HE2	1:B:63:TYR:OH	2.21	0.41
1:B:427:PRO:HB3	1:B:432:SER:HB2	2.01	0.41
2:B:501:FDA:C5X	3:B:502:GRG:H192	2.51	0.41
1:A:158:GLY:HA2	1:A:288:ASP:HB2	2.03	0.40
1:A:175:GLU:HB3	1:A:275:LEU:HD22	2.03	0.40
1:A:138:PHE:CE2	1:B:58:LYS:HG3	2.57	0.40
1:A:426:TYR:CG	1:A:427:PRO:HD2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	465/453 (103%)	456 (98%)	8 (2%)	1 (0%)	44	33
1	B	463/453 (102%)	454 (98%)	8 (2%)	1 (0%)	44	33
All	All	928/906 (102%)	910 (98%)	16 (2%)	2 (0%)	44	33

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	340	TYR
1	B	340	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	397/383 (104%)	392 (99%)	5 (1%)	65	52
1	B	395/383 (103%)	390 (99%)	5 (1%)	65	52
All	All	792/766 (103%)	782 (99%)	10 (1%)	65	52

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	52	SER
1	A	102	ASN
1	A	155	GLU
1	A	187	ARG
1	A	381	GLU
1	B	102	ASN
1	B	146	LEU
1	B	155	GLU
1	B	187	ARG
1	B	382	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	90	ASN
1	A	344	GLN
1	B	90	ASN
1	B	344	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GRG	A	504	-	20,20,28	0.73	0	23,23,37	1.48	5 (21%)
3	GRG	B	504	-	20,20,28	0.76	0	23,23,37	1.42	5 (21%)
3	GRG	B	502	-	27,28,28	0.67	0	33,37,37	1.57	6 (18%)
3	GRG	A	503	-	24,24,28	0.88	0	28,30,37	1.36	4 (14%)
2	FDA	A	501	-	53,58,58	0.55	0	64,89,89	0.68	1 (1%)
3	GRG	A	502	-	27,28,28	0.67	0	33,37,37	1.54	7 (21%)
3	GRG	B	503	-	24,24,28	0.88	0	28,30,37	1.34	4 (14%)
2	FDA	B	501	-	53,58,58	0.55	0	64,89,89	0.71	1 (1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GRG	A	504	-	-	2/21/21/31	-
3	GRG	B	504	-	-	3/21/21/31	-
3	GRG	B	502	-	-	6/31/31/31	-
3	GRG	A	503	-	-	8/25/25/31	-
2	FDA	A	501	-	-	1/30/50/50	0/6/6/6
3	GRG	A	502	-	-	4/31/31/31	-
3	GRG	B	503	-	-	8/25/25/31	-
2	FDA	B	501	-	-	1/30/50/50	0/6/6/6

There are no bond length outliers.

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	502	GRG	C1-C2-C3	-3.78	120.00	126.20
3	A	502	GRG	C6-C7-C8	-3.69	119.19	127.62
3	A	502	GRG	C1-C2-C3	-3.41	120.60	126.20
3	A	503	GRG	C11-C12-C13	-3.40	119.85	127.62
3	B	502	GRG	C6-C7-C8	-3.38	119.88	127.62
3	B	503	GRG	C11-C12-C13	-3.28	120.13	127.62
3	B	502	GRG	C14-C13-C15	3.15	120.69	115.23
3	A	502	GRG	C10-C8-C9	3.14	120.67	115.23
3	B	502	GRG	C10-C8-C9	2.81	120.11	115.23
3	B	504	GRG	C14-C13-C15	2.77	120.04	115.23
3	A	504	GRG	C14-C13-C15	2.66	119.84	115.23
3	B	503	GRG	C10-C8-C9	2.60	119.74	115.23
3	A	503	GRG	C10-C8-C9	2.58	119.70	115.23
3	A	503	GRG	C4-C3-C5	2.56	119.67	115.23
3	B	503	GRG	C6-C7-C8	-2.52	121.86	127.62
3	A	502	GRG	C14-C13-C15	2.52	119.59	115.23
3	B	502	GRG	C4-C3-C5	2.45	119.47	115.23
3	B	502	GRG	C19-C18-C20	2.42	120.16	114.59
3	A	502	GRG	C4-C3-C5	2.38	119.36	115.23
3	A	504	GRG	C11-C12-C13	-2.33	122.29	127.62
3	A	504	GRG	C4-C3-C5	2.33	119.27	115.23
3	B	504	GRG	C4-C3-C5	2.32	119.26	115.23
2	B	501	FDA	C5A-C6A-N6A	2.30	123.82	120.31
2	A	501	FDA	C5A-C6A-N6A	2.30	123.81	120.31
3	A	504	GRG	C19-C18-C20	2.30	119.88	114.59
3	B	504	GRG	C19-C18-C20	2.26	119.80	114.59
3	A	502	GRG	C19-C18-C20	2.23	119.71	114.59
3	B	503	GRG	C4-C3-C5	2.19	119.03	115.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	GRG	C16-C17-C18	-2.16	120.45	127.64
3	B	504	GRG	C6-C7-C8	-2.13	122.74	127.62
3	B	504	GRG	C1-C2-C3	-2.11	119.37	126.38
3	A	503	GRG	C6-C7-C8	-2.04	122.95	127.62
3	A	504	GRG	C16-C17-C18	-2.00	120.97	127.64

There are no chirality outliers.

All (33) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	FDA	C2'-C1'-N10-C10
3	A	502	GRG	C1-O1-PA-O1A
3	A	503	GRG	C1-O1-PA-O1A
3	A	503	GRG	C1-O1-PA-O3A
3	A	503	GRG	C1-O1-PA-O2A
3	B	502	GRG	C1-O1-PA-O1A
3	B	503	GRG	C13-C15-C16-C17
3	B	503	GRG	C1-O1-PA-O1A
3	B	503	GRG	C1-O1-PA-O3A
3	B	503	GRG	C1-O1-PA-O2A
3	A	503	GRG	C13-C15-C16-C17
3	A	503	GRG	C12-C11-C9-C8
3	A	504	GRG	C12-C11-C9-C8
3	B	503	GRG	C12-C11-C9-C8
3	B	504	GRG	C12-C11-C9-C8
3	B	503	GRG	O1-C1-C2-C3
3	B	504	GRG	C7-C8-C9-C11
3	B	504	GRG	C10-C8-C9-C11
3	A	503	GRG	O1-C1-C2-C3
2	B	501	FDA	C2'-C1'-N10-C10
3	A	502	GRG	C1-O1-PA-O3A
3	A	502	GRG	C1-O1-PA-O2A
3	B	502	GRG	C1-O1-PA-O3A
3	B	502	GRG	C1-O1-PA-O2A
3	B	503	GRG	C3-C5-C6-C7
3	B	502	GRG	C13-C15-C16-C17
3	A	504	GRG	C14-C13-C15-C16
3	B	502	GRG	C2-C3-C5-C6
3	A	503	GRG	C9-C11-C12-C13
3	B	503	GRG	C9-C11-C12-C13
3	B	502	GRG	C4-C3-C5-C6
3	A	503	GRG	C3-C5-C6-C7

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	502	GRG	C15-C16-C17-C18


There are no ring outliers.


8 monomers are involved in 18 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	504	GRG	1	0
3	B	504	GRG	1	0
3	B	502	GRG	4	0
3	A	503	GRG	2	0
2	A	501	FDA	4	0
3	A	502	GRG	4	0
3	B	503	GRG	3	0
2	B	501	FDA	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	452/453 (99%)	-1.50	0 100 100	8, 20, 39, 61	15 (3%)
1	B	452/453 (99%)	-1.50	0 100 100	8, 20, 40, 61	13 (2%)
All	All	904/906 (99%)	-1.50	0 100 100	8, 20, 40, 61	28 (3%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GRG	B	504	21/29	0.97	0.07	22,30,41,46	0
3	GRG	A	504	21/29	0.98	0.05	20,31,43,48	0
3	GRG	A	502	29/29	0.98	0.05	12,25,103,123	0
2	FDA	A	501	53/53	0.99	0.02	10,13,18,18	0
3	GRG	B	502	29/29	0.99	0.05	11,25,101,104	0
3	GRG	B	503	25/29	0.99	0.05	26,53,60,63	0
3	GRG	A	503	25/29	0.99	0.05	25,49,59,64	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	FDA	B	501	53/53	1.00	0.02	6,13,16,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around GRG A 504:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GRG A 502:**




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)









Electron density around GRG B 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GRG A 503:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.