



# wwPDB X-ray Structure Validation Summary Report

Jan 27, 2022 – 06:33 pm GMT

PDB ID : 7OP7  
Title : Bacteroides thetaiotaomicron mannosidase GH2 with beta-manno-configured N-alkyl cyclophellitol aziridine  
Authors : McGregor, N.G.S.; Beenakker, T.J.M.; Kuo, C.; Wong, C.; Offen, W.A.; Armstrong, Z.; Codee, J.D.C.; Aerts, J.M.F.G.; Florea, B.I.; Overkleeft, H.S.; Davies, G.J.  
Deposited on : 2021-05-29  
Resolution : 1.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.26  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.26


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	849	 87% 10% ..
1	B	849	 89% 10% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	EDO	B	936	-	-	X	-

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 14952 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-mannosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	837	6985	4467	1174	1310	34	0	36	0
1	B	840	6958	4462	1169	1292	35	0	27	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	MET	-	initiating methionine	UNP Q8AAK6
A	25	GLY	-	expression tag	UNP Q8AAK6
A	865	LEU	-	expression tag	UNP Q8AAK6
A	866	GLU	-	expression tag	UNP Q8AAK6
A	867	HIS	-	expression tag	UNP Q8AAK6
A	868	HIS	-	expression tag	UNP Q8AAK6
A	869	HIS	-	expression tag	UNP Q8AAK6
A	870	HIS	-	expression tag	UNP Q8AAK6
A	871	HIS	-	expression tag	UNP Q8AAK6
A	872	HIS	-	expression tag	UNP Q8AAK6
B	24	MET	-	initiating methionine	UNP Q8AAK6
B	25	GLY	-	expression tag	UNP Q8AAK6
B	865	LEU	-	expression tag	UNP Q8AAK6
B	866	GLU	-	expression tag	UNP Q8AAK6
B	867	HIS	-	expression tag	UNP Q8AAK6
B	868	HIS	-	expression tag	UNP Q8AAK6
B	869	HIS	-	expression tag	UNP Q8AAK6
B	870	HIS	-	expression tag	UNP Q8AAK6
B	871	HIS	-	expression tag	UNP Q8AAK6
B	872	HIS	-	expression tag	UNP Q8AAK6

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 8 4 4	0	1
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is (1R,2S,3R,4S,5R,6R)-5-(8-azidooctylamino)-6-(hydroxymethyl)cyclohexane-1,2,3,4-tetrol (three-letter code: VEE) (formula: C<sub>15</sub>H<sub>30</sub>N<sub>4</sub>O<sub>4</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	12	1	4		
3	B	1	Total	C	N	O	0	0
			17	12	1	4		

- Molecule 4 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	7	Total	Br	0	0
			7	7		
4	B	6	Total	Br	0	0
			6	6		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	6	Total	Cl	0	0
			6	6		
5	B	12	Total	Cl	0	0
			12	12		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	378	Total	O	0	0
			378	378		

*Continued on next page...*

*Continued from previous page...*


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
6	B	450	Total 450	O 450	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-mannosidase



- Molecule 1: Beta-mannosidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.76Å 114.71Å 98.64Å 90.00° 112.89° 90.00°	Depositor
Resolution (Å)	67.66 – 1.85 67.57 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.9 (67.66-1.85) 99.3 (67.57-1.85)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.21 (at 1.86Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.189 , 0.244 0.197 , 0.248	Depositor DCC
$R_{free}$ test set	7826 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.7	Xtriage
Anisotropy	0.482	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	14952	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, BR, VEE, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.88	2/7212 (0.0%)	1.18	12/9803 (0.1%)
1	B	0.90	4/7207 (0.1%)	1.01	2/9795 (0.0%)
All	All	0.89	6/14419 (0.0%)	1.10	14/19598 (0.1%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	591[A]	ILE	CB-CG2	12.37	1.91	1.52
1	A	591[B]	ILE	CB-CG2	12.37	1.91	1.52
1	B	555[A]	GLU	CD-OE2	10.32	1.36	1.25
1	B	555[B]	GLU	CD-OE2	10.32	1.36	1.25
1	B	861	GLU	CD-OE2	6.64	1.32	1.25

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	591[A]	ILE	CA-CB-CG1	-32.45	49.34	111.00
1	A	591[B]	ILE	CA-CB-CG1	-32.45	49.34	111.00
1	A	591[A]	ILE	CA-CB-CG2	-23.37	64.16	110.90
1	A	591[B]	ILE	CA-CB-CG2	-23.37	64.16	110.90
1	A	591[A]	ILE	CB-CG1-CD1	-16.45	67.83	113.90

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6985	0	6677	63	0
1	B	6958	0	6706	74	0
2	A	48	0	71	7	0
2	B	68	0	96	14	0
3	A	17	0	0	1	0
3	B	17	0	0	2	0
4	A	7	0	0	0	0
4	B	6	0	0	1	0
5	A	6	0	0	1	0
5	B	12	0	0	0	0
6	A	378	0	0	12	0
6	B	450	0	0	10	0
All	All	14952	0	13550	138	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 138 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:151[A]:HIS:HE1	2:B:936:EDO:H12	1.20	1.03
1:B:151[A]:HIS:HE1	2:B:936:EDO:C1	1.72	1.01
1:B:151[A]:HIS:CE1	2:B:936:EDO:H12	1.94	1.01
1:B:762:LYS:HD2	6:B:1006:HOH:O	1.64	0.96
1:B:793:MET:SD	1:B:800[B]:CYS:SG	2.64	0.95

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	871/849 (103%)	842 (97%)	25 (3%)	4 (0%)	29	15
1	B	866/849 (102%)	840 (97%)	25 (3%)	1 (0%)	51	36
All	All	1737/1698 (102%)	1682 (97%)	50 (3%)	5 (0%)	47	26

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	536	TRP
1	A	536	TRP
1	A	576[A]	TYR
1	A	576[B]	TYR
1	A	77	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	754/756 (100%)	737 (98%)	17 (2%)	50	34
1	B	755/756 (100%)	743 (98%)	12 (2%)	62	49
All	All	1509/1512 (100%)	1480 (98%)	29 (2%)	59	43

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	848[A]	LYS
1	B	754[B]	ARG
1	B	45	SER
1	B	550	PRO
1	A	850	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	181	HIS

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type
1	B	237	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 62 ligands modelled in this entry, 31 are monoatomic - leaving 31 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	EDO	A	903[B]	-	3,3,3	0.44	0	2,2,2	0.53	0
2	EDO	A	903[A]	-	3,3,3	0.26	0	2,2,2	0.46	0
2	EDO	B	910	-	3,3,3	0.45	0	2,2,2	0.89	0
3	VEE	B	913	1	17,17,23	0.35	0	21,22,28	1.80	4 (19%)
2	EDO	B	908	-	3,3,3	0.20	0	2,2,2	0.30	0
2	EDO	B	903	-	3,3,3	0.36	0	2,2,2	0.34	0
2	EDO	B	933	-	3,3,3	0.63	0	2,2,2	0.41	0
2	EDO	A	906	-	3,3,3	0.71	0	2,2,2	0.71	0
2	EDO	B	936	-	3,3,3	0.15	0	2,2,2	0.12	0
3	VEE	A	911	1	17,17,23	0.74	0	21,22,28	1.49	5 (23%)
2	EDO	B	912	-	3,3,3	0.17	0	2,2,2	0.25	0
2	EDO	A	902	-	3,3,3	0.39	0	2,2,2	0.30	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	EDO	B	905	-	3,3,3	1.33	0	2,2,2	0.82	0
2	EDO	B	911	-	3,3,3	0.25	0	2,2,2	0.52	0
2	EDO	B	904	-	3,3,3	0.37	0	2,2,2	0.57	0
2	EDO	A	907	-	3,3,3	0.30	0	2,2,2	0.58	0
2	EDO	B	909	-	3,3,3	0.66	0	2,2,2	0.25	0
2	EDO	A	901	-	3,3,3	0.37	0	2,2,2	0.65	0
2	EDO	B	935	-	3,3,3	0.17	0	2,2,2	0.17	0
2	EDO	A	905	-	3,3,3	0.41	0	2,2,2	0.36	0
2	EDO	A	910	-	3,3,3	0.15	0	2,2,2	0.18	0
2	EDO	B	902	-	3,3,3	0.30	0	2,2,2	0.40	0
2	EDO	B	906	-	3,3,3	0.74	0	2,2,2	1.31	0
2	EDO	B	934	-	3,3,3	0.42	0	2,2,2	0.21	0
2	EDO	A	925	-	3,3,3	0.21	0	2,2,2	0.65	0
2	EDO	B	907	-	3,3,3	0.36	0	2,2,2	0.12	0
2	EDO	A	908	-	3,3,3	0.47	0	2,2,2	0.66	0
2	EDO	B	901	-	3,3,3	0.21	0	2,2,2	0.29	0
2	EDO	B	932	-	3,3,3	0.37	0	2,2,2	0.48	0
2	EDO	A	904	-	3,3,3	0.48	0	2,2,2	0.18	0
2	EDO	A	909	-	3,3,3	0.63	0	2,2,2	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	903[B]	-	-	0/1/1/1	-
2	EDO	A	903[A]	-	-	0/1/1/1	-
2	EDO	B	910	-	-	0/1/1/1	-
3	VEE	B	913	1	-	2/8/28/34	0/1/1/1
2	EDO	B	908	-	-	0/1/1/1	-
2	EDO	B	903	-	-	0/1/1/1	-
2	EDO	B	933	-	-	0/1/1/1	-
2	EDO	A	906	-	-	1/1/1/1	-
2	EDO	B	936	-	-	1/1/1/1	-
3	VEE	A	911	1	-	3/8/28/34	0/1/1/1
2	EDO	B	912	-	-	0/1/1/1	-
2	EDO	A	902	-	-	1/1/1/1	-
2	EDO	B	905	-	-	0/1/1/1	-
2	EDO	B	911	-	-	1/1/1/1	-
2	EDO	B	904	-	-	0/1/1/1	-
2	EDO	A	907	-	-	1/1/1/1	-

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	B	909	-	-	1/1/1/1	-
2	EDO	A	901	-	-	0/1/1/1	-
2	EDO	B	935	-	-	1/1/1/1	-
2	EDO	A	905	-	-	0/1/1/1	-
2	EDO	A	910	-	-	0/1/1/1	-
2	EDO	B	902	-	-	1/1/1/1	-
2	EDO	B	906	-	-	0/1/1/1	-
2	EDO	B	934	-	-	0/1/1/1	-
2	EDO	A	925	-	-	1/1/1/1	-
2	EDO	B	907	-	-	1/1/1/1	-
2	EDO	A	908	-	-	1/1/1/1	-
2	EDO	B	901	-	-	1/1/1/1	-
2	EDO	B	932	-	-	0/1/1/1	-
2	EDO	A	904	-	-	0/1/1/1	-
2	EDO	A	909	-	-	1/1/1/1	-

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	913	VEE	C5-C6-N7	-5.76	104.09	111.36
3	B	913	VEE	C2-C1-C6	-3.13	105.86	112.42
3	A	911	VEE	C1-C6-C5	-2.92	108.14	111.49
3	A	911	VEE	O12-C3-C2	-2.74	104.75	109.99
3	B	913	VEE	C1-C6-N7	2.53	117.73	112.11

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	907	EDO	O1-C1-C2-O2
2	A	907	EDO	O1-C1-C2-O2
2	B	911	EDO	O1-C1-C2-O2
2	B	936	EDO	O1-C1-C2-O2
3	A	911	VEE	C15-C16-C17-C18

There are no ring outliers.

11 monomers are involved in 24 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	903[B]	EDO	1	0


*Continued on next page...*




*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	903[A]	EDO	3	0
3	B	913	VEE	2	0
2	B	933	EDO	2	0
2	A	906	EDO	2	0
2	B	936	EDO	8	0
3	A	911	VEE	1	0
2	B	905	EDO	2	0
2	A	908	EDO	1	0
2	B	901	EDO	1	0
2	B	932	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	837/849 (98%)	-0.48	6 (0%) 87 88	22, 34, 55, 70	59 (7%)
1	B	840/849 (98%)	-0.46	2 (0%) 95 94	20, 32, 47, 61	76 (9%)
All	All	1677/1698 (98%)	-0.47	8 (0%) 91 91	20, 33, 51, 70	135 (8%)

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	864	LYS	3.5
1	A	849	GLY	3.4
1	B	472	PHE	2.9
1	A	472	PHE	2.7
1	A	470	TRP	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	EDO	B	934	4/4	0.74	0.22	30,31,32,36	4
2	EDO	A	906	4/4	0.80	0.15	37,43,44,45	0
2	EDO	B	905	4/4	0.81	0.13	30,30,31,36	4
2	EDO	B	912	4/4	0.83	0.18	36,41,42,43	4
2	EDO	A	908	4/4	0.83	0.27	21,21,27,32	4
2	EDO	B	935	4/4	0.84	0.14	41,45,46,51	4
2	EDO	B	907	4/4	0.87	0.23	38,42,44,49	4
2	EDO	A	905	4/4	0.88	0.16	26,29,29,35	4
2	EDO	B	936	4/4	0.88	0.21	33,36,36,38	4
5	CL	B	924	1/1	0.88	0.07	60,60,60,60	0
3	VEE	B	913	17/23	0.89	0.15	21,36,42,45	17
2	EDO	A	907	4/4	0.89	0.12	31,41,44,46	4
2	EDO	A	925	4/4	0.90	0.18	26,26,28,29	4
2	EDO	B	901	4/4	0.90	0.08	43,44,45,48	0
5	CL	B	926	1/1	0.90	0.11	68,68,68,68	0
3	VEE	A	911	17/23	0.91	0.13	32,39,48,50	4
2	EDO	B	909	4/4	0.91	0.23	22,24,26,30	4
5	CL	A	923	1/1	0.91	0.24	60,60,60,60	0
2	EDO	B	910	4/4	0.91	0.10	41,43,45,48	0
2	EDO	B	906	4/4	0.91	0.27	13,18,18,23	4
2	EDO	A	903[A]	4/4	0.92	0.13	22,28,29,32	4
2	EDO	B	933	4/4	0.92	0.18	22,24,24,27	4
2	EDO	A	903[B]	4/4	0.92	0.13	24,25,27,28	4
2	EDO	B	902	4/4	0.92	0.23	32,35,36,36	4
2	EDO	A	910	4/4	0.92	0.23	37,39,41,48	4
2	EDO	B	911	4/4	0.93	0.20	40,43,45,46	4
2	EDO	A	901	4/4	0.93	0.19	12,13,14,14	4
5	CL	B	920	1/1	0.93	0.06	62,62,62,62	0
2	EDO	B	932	4/4	0.93	0.14	29,30,31,31	4
2	EDO	B	908	4/4	0.93	0.25	49,54,62,66	0
2	EDO	A	902	4/4	0.94	0.08	37,39,44,47	0
5	CL	A	924	1/1	0.94	0.06	50,50,50,50	0
5	CL	A	919	1/1	0.94	0.16	56,56,56,56	0
5	CL	A	921	1/1	0.94	0.07	46,46,46,46	0
5	CL	A	922	1/1	0.94	0.05	62,62,62,62	0
5	CL	B	929	1/1	0.94	0.04	60,60,60,60	0
5	CL	B	930	1/1	0.94	0.12	62,62,62,62	0
5	CL	B	923	1/1	0.95	0.07	58,58,58,58	0
5	CL	B	928	1/1	0.95	0.10	66,66,66,66	0
4	BR	B	918	1/1	0.96	0.05	52,52,52,52	1
4	BR	B	919	1/1	0.97	0.06	48,48,48,48	1
5	CL	B	925	1/1	0.97	0.12	61,61,61,61	0
2	EDO	A	909	4/4	0.97	0.12	46,49,58,64	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	EDO	B	903	4/4	0.97	0.08	30,34,41,43	0
5	CL	B	922	1/1	0.97	0.06	57,57,57,57	0
2	EDO	A	904	4/4	0.97	0.09	29,34,39,43	0
5	CL	B	921	1/1	0.98	0.06	42,42,42,42	0
4	BR	A	914	1/1	0.98	0.05	48,48,48,48	1
5	CL	B	927	1/1	0.98	0.04	56,56,56,56	0
5	CL	B	931	1/1	0.98	0.13	38,38,38,38	0
4	BR	B	914	1/1	0.99	0.03	37,37,37,37	0
4	BR	B	915	1/1	0.99	0.03	38,38,38,38	1
4	BR	B	916	1/1	0.99	0.03	56,56,56,56	0
4	BR	B	917	1/1	0.99	0.03	46,46,46,46	0
4	BR	A	912	1/1	0.99	0.02	41,41,41,41	0
4	BR	A	913	1/1	0.99	0.03	35,35,35,35	0
2	EDO	B	904	4/4	0.99	0.08	22,25,25,26	0
5	CL	A	920	1/1	0.99	0.06	63,63,63,63	0
4	BR	A	915	1/1	0.99	0.04	30,30,30,30	0
4	BR	A	916	1/1	0.99	0.07	45,45,45,45	0
4	BR	A	917	1/1	0.99	0.08	58,58,58,58	0
4	BR	A	918	1/1	0.99	0.11	49,49,49,49	1




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around VEE B 913:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around VEE A 911:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.