



wwPDB X-ray Structure Validation Summary Report ⓘ

May 14, 2020 – 03:18 am BST

PDB ID : 5OQF
Title : Crystal structure of a single chain trimer composed of the MHC 1 heavy chain H2-Kb WT, beta-2microglobulin, and ovalbumin derived peptide
Authors : Mikolajek, H.; Werner, J.M.
Deposited on : 2017-08-11
Resolution : 2.27 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6980 (2.30-2.26)
Clashscore	141614	7711 (2.30-2.26)
Ramachandran outliers	138981	7597 (2.30-2.26)
Sidechain outliers	138945	7598 (2.30-2.26)
RSRZ outliers	127900	6849 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	431	
1	B	431	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 6727 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-2-microglobulin,H-2 class I histocompatibility antigen, K-B alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	388	3150	1993	548	591	18	0	0	0
1	B	388	3158	1999	549	592	18	0	1	0

There are 102 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP P01887
A	2	ILE	-	expression tag	UNP P01887
A	3	ILE	-	expression tag	UNP P01887
A	4	ASN	-	expression tag	UNP P01887
A	5	PHE	-	expression tag	UNP P01887
A	6	GLU	-	expression tag	UNP P01887
A	7	LYS	-	expression tag	UNP P01887
A	8	LEU	-	expression tag	UNP P01887
A	9	GLY	-	expression tag	UNP P01887
A	10	CYS	-	expression tag	UNP P01887
A	11	GLY	-	expression tag	UNP P01887
A	12	ALA	-	expression tag	UNP P01887
A	13	SER	-	expression tag	UNP P01887
A	?	GLY	-	expression tag	UNP P01887
A	?	GLY	-	expression tag	UNP P01887
A	?	GLY	-	expression tag	UNP P01887
A	?	GLY	-	expression tag	UNP P01887
A	?	SER	-	expression tag	UNP P01887
A	?	GLY	-	expression tag	UNP P01887
A	?	GLY	-	expression tag	UNP P01887
A	?	GLY	-	expression tag	UNP P01887
A	?	GLY	-	expression tag	UNP P01887
A	?	GLY	-	expression tag	UNP P01887
A	?	GLY	-	expression tag	UNP P01887
A	?	SER	-	expression tag	UNP P01887
A	?	GLY	-	linker	UNP P01887

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	SER	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	SER	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	SER	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	SER	-	linker	UNP P01887
A	143	GLY	-	linker	UNP P01887
A	226	CYS	TYR	engineered mutation	UNP P01901
A	426	HIS	-	expression tag	UNP P01901
A	427	HIS	-	expression tag	UNP P01901
A	428	HIS	-	expression tag	UNP P01901
A	429	HIS	-	expression tag	UNP P01901
A	430	HIS	-	expression tag	UNP P01901
A	431	HIS	-	expression tag	UNP P01901
B	1	SER	-	expression tag	UNP P01887
B	2	ILE	-	expression tag	UNP P01887
B	3	ILE	-	expression tag	UNP P01887
B	4	ASN	-	expression tag	UNP P01887
B	5	PHE	-	expression tag	UNP P01887
B	6	GLU	-	expression tag	UNP P01887
B	7	LYS	-	expression tag	UNP P01887
B	8	LEU	-	expression tag	UNP P01887
B	9	GLY	-	expression tag	UNP P01887
B	10	CYS	-	expression tag	UNP P01887
B	11	GLY	-	expression tag	UNP P01887
B	12	ALA	-	expression tag	UNP P01887
B	13	SER	-	expression tag	UNP P01887
B	?	GLY	-	expression tag	UNP P01887
B	?	GLY	-	expression tag	UNP P01887

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	200	Total 200	O 200	0	0

4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.65Å 90.08Å 89.33Å 90.00° 111.23° 90.00°	Depositor
Resolution (Å)	83.27 - 2.27 83.27 - 2.27	Depositor EDS
% Data completeness (in resolution range)	98.5 (83.27-2.27) 98.5 (83.27-2.27)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 2.27Å)	Xtrriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.200 , 0.250 0.200 , 0.250	Depositor DCC
R_{free} test set	2266 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	35.1	Xtrriage
Anisotropy	0.838	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6727	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 38.56 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.6214e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/3237	0.59	0/4392
1	B	0.43	0/3245	0.58	0/4403
All	All	0.43	0/6482	0.59	0/8795

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3150	0	3016	46	0
1	B	3158	0	3026	42	0
2	A	219	0	0	5	0
2	B	200	0	0	11	0
All	All	6727	0	6042	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 87 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:187:TYR:HE1	1:A:209:ALA:HB2	1.35	0.89

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2:ILE:O	1:A:208:LYS:NZ	2.06	0.87
1:B:8:LEU:HD23	1:B:285:THR:HG21	1.59	0.85
1:B:336:ARG:O	1:B:338:GLU:N	2.16	0.78
1:A:166:GLU:OE1	1:A:187:TYR:OH	2.04	0.75

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	382/431 (89%)	370 (97%)	12 (3%)	0	100	100
1	B	383/431 (89%)	370 (97%)	11 (3%)	2 (0%)	29	34
All	All	765/862 (89%)	740 (97%)	23 (3%)	2 (0%)	41	49

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	337	PRO
1	B	70	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	337/356 (95%)	333 (99%)	4 (1%)	71	82
1	B	338/356 (95%)	332 (98%)	6 (2%)	59	72
All	All	675/712 (95%)	665 (98%)	10 (2%)	69	77

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	93	PHE
1	B	181	ASP
1	B	215	SER
1	A	181	ASP
1	B	187	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	388/431 (90%)	0.09	12 (3%) 49 54	22, 37, 57, 79	0
1	B	388/431 (90%)	0.09	8 (2%) 63 69	22, 36, 55, 77	0
All	All	776/862 (90%)	0.09	20 (2%) 56 62	22, 37, 57, 79	0

The worst 5 of 20 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	367	ILE	6.2
1	B	369	ASP	4.0
1	B	368	GLN	3.9
1	A	337	PRO	3.9
1	B	317	GLY	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.