



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 23, 2021 – 01:14 PM EDT

PDB ID : 1OSP
Title : CRYSTAL STRUCTURE OF OUTER SURFACE PROTEIN A OF BORRELIA BURGdorFERI COMPLEXED WITH A MURINE MONOCLONAL ANTIBODY FAB
Authors : Li, H.; Lawson, C.L.
Deposited on : 1996-11-23
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	L	214	
2	H	218	
3	O	257	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7418 atoms, of which 1881 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FAB 184.1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	L	214	2057	1036	392	278	344	7	29	0	0

- Molecule 2 is a protein called FAB 184.1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	H	218	2015	1048	364	264	332	7	49	0	0

- Molecule 3 is a protein called OUTER SURFACE PROTEIN A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
3	O	251	2362	1170	469	313	408	2	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
O	17	ALA	CYS	conflict	UNP P14013
O	39	LYS	ASN	variant	UNP P14013
O	84	CYS	SER	engineered mutation	UNP P14013
O	149	GLY	GLU	variant	UNP P14013
O	164	GLY	SER	variant	UNP P14013

- Molecule 4 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	L	119	Total	H	O	0	0
			357	238	119		
4	H	99	Total	H	O	0	0
			297	198	99		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	H	O		
4	O	110	330	220	110	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: FAB 184.1



- Molecule 2: FAB 184.1



- Molecule 3: OUTER SURFACE PROTEIN A



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	89.10Å 91.70Å 100.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 1.95	Depositor
% Data completeness (in resolution range)	(Not available) (6.00-1.95)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.229 , 0.295	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	7418	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	L	0.62	3/1704 (0.2%)	0.78	3/2315 (0.1%)
2	H	0.56	1/1698 (0.1%)	0.88	4/2323 (0.2%)
3	O	0.37	0/1902	0.66	0/2550
All	All	0.52	4/5304 (0.1%)	0.78	7/7188 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	L	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	133	PRO	C-N	14.55	1.59	1.33
1	L	213	GLU	C-N	14.54	1.67	1.34
1	L	211	ARG	C-N	-8.24	1.15	1.34
1	L	212	ASN	C-N	-7.93	1.15	1.34

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	133	PRO	O-C-N	15.69	149.87	123.20
2	H	133	PRO	CA-C-N	-13.59	89.02	116.20
1	L	211	ARG	O-C-N	-10.91	105.24	122.70
2	H	4	LEU	CA-CB-CG	6.27	129.73	115.30
2	H	133	PRO	C-N-CA	-6.15	109.39	122.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	L	211	ARG	Mainchain
1	L	212	ASN	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1665	392	1589	32	3
2	H	1651	364	1598	49	0
3	O	1893	469	1965	21	6
4	H	99	198	0	2	1
4	L	119	238	0	2	6
4	O	110	220	0	1	0
All	All	5537	1881	5152	98	8

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 98 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:170:VAL:HG22	2:H:188:VAL:HG22	1.39	1.04
3:O:159:LYS:HG2	3:O:168:GLU:HG2	1.53	0.88
1:L:18:ARG:NH2	1:L:76:THR:HG22	1.93	0.84
2:H:2:VAL:HG13	2:H:27:GLU:HB3	1.59	0.84
3:O:24:LEU:HG	3:O:55:ILE:HD12	1.62	0.80

The worst 5 of 8 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:O:230:LYS:HZ3	4:L:275:HOH:H1[3_446]	0.65	0.95
3:O:254:LYS:HZ1	4:L:258:HOH:H2[3_446]	0.66	0.94
1:L:188:ARG:HH12	4:L:281:HOH:H1[4_456]	1.09	0.51
1:L:61:ARG:HH22	4:H:306:HOH:H1[4_556]	1.14	0.46
3:O:254:LYS:HZ1	4:L:258:HOH:O[3_446]	1.27	0.33

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	212/214 (99%)	201 (95%)	11 (5%)	0	100	100
2	H	216/218 (99%)	204 (94%)	9 (4%)	3 (1%)	11	3
3	O	249/257 (97%)	245 (98%)	4 (2%)	0	100	100
All	All	677/689 (98%)	650 (96%)	24 (4%)	3 (0%)	34	22

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	153	PHE
2	H	156	SER
2	H	195	TRP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	192/192 (100%)	191 (100%)	1 (0%)	88	88
2	H	190/190 (100%)	181 (95%)	9 (5%)	26	13
3	O	220/225 (98%)	216 (98%)	4 (2%)	59	53
All	All	602/607 (99%)	588 (98%)	14 (2%)	50	42

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	139	THR
2	H	155	GLU
3	O	213	THR
3	O	128	GLU
3	O	175	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	137	ASN
1	L	198	HIS
2	H	43	ASN
2	H	171	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	L	3

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	L	213:GLU	C	214:CYS	N	1.67
1	L	211:ARG	C	212:ASN	N	1.15
1	L	212:ASN	C	213:GLU	N	1.15

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.