

wwPDB X-ray Structure Validation Summary Report (i)

Aug 6, 2020 – 02:46 PM BST

PDB ID : 5OUG

Title : Humanized alpha-AChBP (acetylcholine binding protein) in complex with lo-

beline and allosteric binder fragment 4.

Authors : Delbart, F.; Gruss, F.; Ulens, C.

Deposited on : 2017-08-23

Resolution : 2.57 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.13.1 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.13.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.57 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution			
Metric	$(\# {\rm Entries})$	$(\# ext{Entries}, ext{resolution range}(ext{Å}))$			
R_{free}	130704	3676 (2.60-2.56)			
Clashscore	141614	4049 (2.60-2.56)			
Ramachandran outliers	138981	3979 (2.60-2.56)			
Sidechain outliers	138945	3979 (2.60-2.56)			
RSRZ outliers	127900	3614 (2.60-2.56)			

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	Δ.	00.5	4%	
1	A	205	91%	9%
1	В	205	90%	9% •
_			5%	370 -
1	С	205	89%	11%
1	D	205	91%	9%
			5%	
1	Е	205	91%	9%
2	F	2	100%	
	1		100%	





Continued from previous page...

Mol	Chain	Length	Quality of chain							
2	J	2	50%	50%						
3	G	5	20%	80%						
4	Н	3		100%						
4	I	3		100%						

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	L0B	A	1106	-	-	-	X
5	L0B	В	308	-	-	-	X
5	L0B	С	306	-	-	-	X
5	L0B	D	306	-	-	-	X
6	9Z0	A	1104	-	-	-	X
6	9Z0	В	306	-	-	-	X
6	9Z0	В	307	-	-	-	X
6	9Z0	С	304	-	-	-	X
6	9Z0	D	304	-	-	-	X
6	9Z0	D	305	-	-	-	X
6	9Z0	E	302	-	-	-	X



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 8950 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Humanized alpha-AChBP.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	205	Total	С	N	О	S	0	0	0
1	A	200	1680	1077	282	314	7	0	0	U
1	В	205	Total	С	N	О	S	0	0	0
1	Ъ	200	1680	1077	282	314	7	0	0	U
1	С	205	Total	С	N	О	S	0	0	0
1		200	1680	1077	282	314	7	0		
1	D	205	Total	С	N	О	S	0	0	0
1	ש	200	1680	1077	282	314	7	0	0	"
1	Е	205	Total	С	N	О	S	0	0	0
1	Ľ	200	1676	1075	282	312	7	U	U	U

• Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace		
9	Ē	9	Total	С	N	О	0	0	0
	1		28	16	2	10	U	U	
9	ī	9	Total	С	N	О	0	0	0
	J		28	16	2	10	0	0	U

• Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.






Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace
3	G	5	Total C N 61 34 2	O 25	0	0	0

• Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-b eta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
4	Н	3	Total C N 39 22 2	0	0	0
4	I	3	Total C N 39 22 2	0	0	0

• Molecule 5 is Alpha-Lobeline (three-letter code: L0B) (formula: $C_{22}H_{27}NO_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C N O 25 22 1 2	0	0
5	A	1	Total C N O 25 22 1 2	0	0
5	В	1	Total C N O 25 22 1 2	0	0
5	С	1	Total C N O 25 22 1 2	0	0


Continued on next page...



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	D	1	Total	С	N	О	0	0
)	D	1	25	22	1	2	0	0


• Molecule 6 is 4,5-dibromo-N-(3-hydroxypropyl)-1H-pyrrole-2-carboxamide (three-letter code: 9Z0) (formula: $C_8H_{10}Br_2N_2O_2$).



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	
6	A	1	Total	Br	С	N	О	0	0	
0	A	1	14	2	8	2	2	0	0	
6	A	1	Total	Br	С	N	О	0	0	
0	Λ	1	14	2	8	2	2		0	
6	В	1	Total	Br	С	N	О	0	0	
0	Ъ	1	14	2	8	2	2	U	U	
6	В	1	Total	Br	С	N	О	0	0	
0	Ъ	1	14	2	8	2	2		0	
6	С	1	Total	Br	С	N	О	0	0	
0		1	14	2	8	2	2	0	U	
6	С	1	Total	Br	С	Ν	О	0	0	
0		1	14	2	8	2	2	0	U	
6	D	1	Total	Br	С	N	Ο	0	0	
	D	1	14	2	8	2	2	0	U	
6	$_{ m D}$	1	Total	Br	С	Ν	Ο	0	0	
	D	1	14	2	8	2	2	U	U	
6	E	1	Total	Br	С	Ν	Ο	0	0	
	Ľ	1	14	2	8	2	2		U	
6	E	1	Total	Br	С	N	О	0	0	
	127	1	14	2	8	2	2		U	



 \bullet Molecule 7 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: $\mathrm{C_2H_6OS}).$



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	E	1	Total 4	C 2	O 1	S 1	0	0

• Molecule 8 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	17	Total O 17 17	0	0
8	В	26	Total O 26 26	0	0
8	С	19	Total O 19 19	0	0
8	D	18	Total O 18 18	0	0
8	Е	10	Total O 10 10	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Humanized alpha-AChBP





• Molecule 2 opyranose	: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido	o-2-deoxy-beta-D-gluc
Chain F:	100%	
NAG2 NAG2		
• Molecule 2 opyranose	: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido	o-2-deoxy-beta-D-gluc
Chain J:	50% 50%	
NAG1 NAG2		
	: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6) etamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-de	
Chain G:	20% 80%	
NAG1 NAG2 BMA3 MAN4 MAN5		
	: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glueoxy-beta-D-glucopyranose	acopyranose-(1-4)-2-ac
Chain H:	100%	
NAG1 NAG2 BMA3		
	: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glueoxy-beta-D-glucopyranose	acopyranose-(1-4)-2-ac
Chain I:	100%	
NAG1 NAG2 BMA3		



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	68.67Å 119.81Å 84.54Å	Depositor
a, b, c, α , β , γ	90.00° 107.29° 90.00°	Depositor
Resolution (Å)	80.72 - 2.57	Depositor
Resolution (A)	66.94 - 2.57	EDS
% Data completeness	99.7 (80.72-2.57)	Depositor
(in resolution range)	99.7 (66.94-2.57)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.54 (at 2.58Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
D D.	0.214 , 0.240	Depositor
R, R_{free}	0.222 , 0.245	DCC
R_{free} test set	2109 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	63.6	Xtriage
Anisotropy	0.272	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.32 , 43.1	EDS
L-test for twinning ²	$ < L > = 0.50, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8950	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.79% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 9Z0, BMA, NAG, DMS, L0B, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	В	ond angles
		RMSZ	# Z >5	RMSZ	# Z > 5
1	A	0.64	0/1724	0.82	$4/2344 \ (0.2\%)$
1	В	0.64	0/1724	0.86	5/2344 (0.2%)
1	С	0.61	0/1724	0.82	2/2344~(0.1%)
1	D	0.64	0/1724	0.83	3/2344 (0.1%)
1	E	0.64	0/1720	0.84	$4/2339 \ (0.2\%)$
All	All	0.63	0/8616	0.83	$18/11715 \ (0.2\%)$

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	В	0	1
1	E	0	1
All	All	0	3

There are no bond length outliers.

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
1	В	131	ASP	N-CA-CB	8.34	125.61	110.60
1	D	131	ASP	N-CA-CB	8.33	125.59	110.60
1	E	122	ARG	NE-CZ-NH2	7.01	123.81	120.30
1	В	122	ARG	NE-CZ-NH2	6.94	123.77	120.30
1	D	120	ARG	NE-CZ-NH1	6.32	123.46	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	A	186	CYS	Peptide
1	В	186	CYS	Peptide
1	Е	186	CYS	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Α	1680	0	1651	10	0
1	В	1680	0	1651	11	0
1	С	1680	0	1651	13	0
1	D	1680	0	1651	12	0
1	E	1676	0	1647	10	0
2	F	28	0	25	0	0
2	J	28	0	25	0	0
3	G	61	0	52	1	0
4	Н	39	0	34	0	0
4	I	39	0	34	0	0
5	A	50	0	54	6	0
5	В	25	0	27	0	0
5	С	25	0	27	1	0
5	D	25	0	27	2	0
6	A	28	0	20	4	0
6	В	28	0	20	4	0
6	С	28	0	20	4	0
6	D	28	0	20	4	0
6	E	28	0	20	3	0
7	Ε	4	0	6	0	0
8	A	17	0	0	0	0
8	В	26	0	0	0	0
8	С	19	0	0	1	0
8	D	18	0	0	0	0
8	E	10	0	0	0	0
All	All	8950	0	8662	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 56 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	$egin{aligned} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{aligned}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:D:182:ARG:NH1	1:D:193:ASP:OD2	2.16	0.77
1:C:78:VAL:CG1	1:C:83:LEU:HD11	2.25	0.66
1:C:78:VAL:HG12	1:C:83:LEU:HD11	1.80	0.64
1:D:7:TYR:HA	6:D:305:9Z0:BR2	2.52	0.63
5:C:306:L0B:C5	1:D:116:LEU:HD13	2.30	0.62

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	203/205~(99%)	199 (98%)	3 (2%)	1 (0%)	29	50
1	В	203/205~(99%)	198 (98%)	4 (2%)	1 (0%)	29	50
1	С	203/205~(99%)	199 (98%)	3 (2%)	1 (0%)	29	50
1	D	203/205~(99%)	198 (98%)	4 (2%)	1 (0%)	29	50
1	E	203/205~(99%)	199 (98%)	3 (2%)	1 (0%)	29	50
All	All	1015/1025~(99%)	993 (98%)	17 (2%)	5 (0%)	29	50

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	24	ASP
1	В	24	ASP
1	С	24	ASP
1	E	24	ASP
1	D	24	ASP



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percer	$_{ m tiles}$
1	A	$192/192 \ (100\%)$	189 (98%)	3 (2%)	62	81
1	В	192/192 (100%)	189 (98%)	3 (2%)	62	81
1	С	192/192 (100%)	188 (98%)	4 (2%)	53	75
1	D	$192/192 \; (100\%)$	190 (99%)	2 (1%)	76	89
1	E	191/192 (100%)	190 (100%)	1 (0%)	88	96
All	All	959/960 (100%)	946 (99%)	13 (1%)	67	84

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	133	GLU
1	С	25	ARG
1	D	23	ARG
1	В	131	ASP
1	С	182	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	${f Res}$	\mathbf{Type}
1	С	177	GLN
1	D	37	GLN
1	E	61	HIS
1	С	61	HIS
1	Ε	103	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

15 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

N / - 1	TD.	C1 '	D	т. 1	Во	nd leng	ths	Bond angles		
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	F	1	1,2	14,14,15	0.42	0	17,19,21	1.09	2 (11%)
2	NAG	F	2	2	14,14,15	0.54	0	17,19,21	1.36	2 (11%)
3	NAG	G	1	1,3	14,14,15	0.47	0	17,19,21	0.96	0
3	NAG	G	2	3	14,14,15	0.43	0	17,19,21	0.79	0
3	BMA	G	3	3	11,11,12	0.46	0	15,15,17	1.22	2 (13%)
3	MAN	G	4	3	11,11,12	0.53	0	15,15,17	1.03	1 (6%)
3	MAN	G	5	3	11,11,12	0.32	0	15,15,17	1.06	1 (6%)
4	NAG	Н	1	1,4	14,14,15	0.40	0	17,19,21	1.34	3 (17%)
4	NAG	Н	2	4	14,14,15	0.49	0	17,19,21	1.04	1 (5%)
4	BMA	Н	3	4	11,11,12	0.62	0	15,15,17	1.00	1 (6%)
4	NAG	I	1	1,4	14,14,15	0.50	0	17,19,21	1.27	1 (5%)
4	NAG	I	2	4	14,14,15	0.66	0	17,19,21	1.38	3 (17%)
4	BMA	I	3	4	11,11,12	0.67	0	15,15,17	1.48	2 (13%)
2	NAG	J	1	1,2	14,14,15	0.48	0	17,19,21	1.31	2 (11%)
2	NAG	J	2	2	14,14,15	0.36	0	17,19,21	1.02	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	F	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	F	2	2	-	0/6/23/26	0/1/1/1

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	G	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	G	2	3	-	0/6/23/26	0/1/1/1
3	BMA	G	3	3	-	0/2/19/22	0/1/1/1
3	MAN	G	4	3	-	0/2/19/22	0/1/1/1
3	MAN	G	5	3	-	0/2/19/22	0/1/1/1
4	NAG	Н	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	Н	2	4	-	2/6/23/26	0/1/1/1
4	BMA	Н	3	4	-	0/2/19/22	0/1/1/1
4	NAG	I	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	I	2	4	-	0/6/23/26	0/1/1/1
4	BMA	I	3	4	-	0/2/19/22	0/1/1/1
2	NAG	J	1	1,2	=	0/6/23/26	0/1/1/1
2	NAG	J	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^o)$
2	J	1	NAG	O5-C1-C2	-3.48	105.79	111.29
4	Н	1	NAG	O5-C1-C2	-3.38	105.94	111.29
4	I	3	BMA	C3-C4-C5	3.24	116.02	110.24
4	I	1	NAG	O5-C1-C2	-3.23	106.19	111.29
3	G	3	BMA	C1-O5-C5	3.19	116.52	112.19

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	J	2	NAG	O5-C5-C6-O6
2	J	2	NAG	C4-C5-C6-O6
4	Н	2	NAG	C4-C5-C6-O6
4	Н	2	NAG	O5-C5-C6-O6

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	G	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,




bond angles, torsion angles, and ring geometry for oligosaccharide.



















5.6 Ligand geometry (i)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tune	Chain	Res	Link	Вс	ond leng	ths	Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	9Z0	В	307	-	12,14,14	0.88	1 (8%)	10,18,18	0.96	0
6	9Z0	D	305	-	12,14,14	0.80	1 (8%)	10,18,18	0.95	0
6	9Z0	A	1104	-	12,14,14	0.91	1 (8%)	10,18,18	0.80	0
6	9Z0	С	305	-	12,14,14	0.87	1 (8%)	10,18,18	1.31	2 (20%)
5	L0B	D	306	-	27,27,27	0.50	0	33,36,36	1.25	2 (6%)
6	9Z0	С	304	-	12,14,14	1.01	1 (8%)	10,18,18	0.93	0
6	9Z0	D	304	-	12,14,14	1.07	1 (8%)	10,18,18	0.91	1 (10%)
5	L0B	С	306	-	27,27,27	0.49	0	33,36,36	1.20	2 (6%)



Mol	Tune	Chain	Res Link		Во	ond leng	ths	Bond angles			
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
6	9Z0	A	1105	-	12,14,14	0.87	1 (8%)	10,18,18	0.95	0	
5	L0B	В	308	-	27,27,27	0.47	0	33,36,36	1.17	3 (9%)	
5	L0B	A	1106	-	27,27,27	0.54	0	33,36,36	1.19	1 (3%)	
7	DMS	E	303	-	3,3,3	0.54	0	3,3,3	0.74	0	
6	9Z0	В	306	_	12,14,14	0.98	1 (8%)	10,18,18	0.90	0	
6	9Z0	E	302	-	12,14,14	0.92	1 (8%)	10,18,18	0.85	0	
6	9Z0	Е	301	-	12,14,14	1.00	1 (8%)	10,18,18	0.98	0	
5	L0B	A	1101	-	27,27,27	0.49	0	33,36,36	1.27	3 (9%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	9Z0	В	307	_	-	2/6/9/9	0/1/1/1
6	9Z0	D	305	_	-	1/6/9/9	0/1/1/1
6	9Z0	A	1104	-	-	2/6/9/9	0/1/1/1
6	9Z0	С	305	_	-	2/6/9/9	0/1/1/1
5	L0B	D	306	-	-	8/16/30/30	0/3/3/3
6	9Z0	С	304	-	-	0/6/9/9	0/1/1/1
6	9Z0	D	304	-	-	3/6/9/9	0/1/1/1
5	L0B	С	306	_	-	6/16/30/30	0/3/3/3
6	9Z0	A	1105	_	-	2/6/9/9	0/1/1/1
5	L0B	В	308	-	-	9/16/30/30	0/3/3/3
5	L0B	A	1106	-	-	4/16/30/30	0/3/3/3
6	9Z0	В	306	-	-	1/6/9/9	0/1/1/1
6	9Z0	Е	302		-	1/6/9/9	0/1/1/1
6	9Z0	Е	301	-	-	2/6/9/9	0/1/1/1
5	L0B	A	1101	-	-	4/16/30/30	0/3/3/3

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(ext{\AA})$
6	С	304	9Z0	BR2-C12	-2.81	1.85	1.90
6	E	301	9Z0	BR2-C12	-2.77	1.85	1.90
6	В	306	9Z0	BR2-C12	-2.67	1.86	1.90
6	A	1104	9Z0	BR2-C12	-2.60	1.86	1.90

Continued on next page...



Continued from previous page...

\mathbf{Mol}	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$Ideal(\AA)$
6	D	304	9Z0	BR2-C12	-2.54	1.86	1.90

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
5	A	1106	L0B	C22-N1-C11	4.89	117.11	113.16
5	D	306	L0B	C22-N1-C11	4.74	116.99	113.16
5	В	308	L0B	C22-N1-C11	4.39	116.71	113.16
5	A	1101	L0B	C22-N1-C11	4.30	116.63	113.16
5	С	306	L0B	C22-N1-C11	3.96	116.36	113.16

There are no chirality outliers.

5 of 47 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	В	307	9Z0	N03-C04-C05-C06
6	D	305	9Z0	N03-C04-C05-C06
6	A	1104	9Z0	C08-C02-N03-C04
6	С	305	9Z0	C04-C05-C06-O07
5	D	306	L0B	C11-C14-C16-C17

There are no ring outliers.


14 monomers are involved in 28 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	В	307	9Z0	1	0
6	D	305	9Z0	1	0
6	A	1104	9Z0	2	0
6	С	305	9Z0	1	0
5	D	306	L0B	2	0
6	С	304	9Z0	3	0
6	D	304	9Z0	3	0
5	С	306	L0B	1	0
6	A	1105	9Z0	2	0
5	A	1106	L0B	3	0
6	В	306	9Z0	3	0
6	Е	302	9Z0	1	0
6	Е	301	9Z0	2	0
5	A	1101	L0B	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,




bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











































Torsions









5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2		$\mathrm{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	$205/205 \; (100\%)$	0.43	9 (4%) 34	30	42, 68, 111, 131	0
1	В	$205/205 \; (100\%)$	0.47	9 (4%) 34	30	44, 65, 115, 133	0
1	С	$205/205 \; (100\%)$	0.56	11 (5%) 25	22	43, 71, 121, 146	0
1	D	$205/205 \; (100\%)$	0.50	13 (6%) 20	17	46, 72, 122, 134	0
1	E	$205/205 \; (100\%)$	0.49	10 (4%) 29	26	46, 68, 114, 126	0
All	All	$1025/1025 \; (100\%)$	0.49	52 (5%) 28	24	42, 69, 117, 146	0

The worst 5 of 52 RSRZ outliers are listed below:

Mol	Chain	${f Res}$	Type	RSRZ
1	D	161	ILE	6.4
1	С	204	GLY	5.8
1	D	68	SER	4.8
1	Е	11	VAL	4.8
1	С	159	ALA	4.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mo	l Type	Chain	Res	Atoms	RSCC	RSR	${f B\text{-factors}}({f A}^2)$	Q<0.9
4	BMA	I	3	11/12	0.42	0.36	104,122,125,126	0


Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{\textbf{B-factors}}(\mbox{\c A}^2)$	Q < 0.9
2	NAG	F	2	14/15	0.75	0.28	70,110,116,118	0
4	BMA	Н	3	11/12	0.78	0.19	90,104,111,114	0
4	NAG	I	2	14/15	0.83	0.24	87,108,118,120	0
4	NAG	Н	2	14/15	0.84	0.31	96,111,115,116	0
4	NAG	Н	1	14/15	0.85	0.15	73,86,98,106	0
2	NAG	J	2	14/15	0.87	0.18	94,120,129,130	0
3	MAN	G	4	11/12	0.90	0.21	77,88,94,96	0
3	MAN	G	5	11/12	0.91	0.21	86,87,92,93	0
2	NAG	J	1	14/15	0.92	0.10	73,82,88,97	0
3	BMA	G	3	11/12	0.92	0.12	81,89,90,91	0
3	NAG	G	1	14/15	0.93	0.14	58,63,71,76	0
3	NAG	G	2	14/15	0.93	0.15	64,78,84,87	0
4	NAG	I	1	14/15	0.94	0.12	68,77,86,92	0
2	NAG	F	1	14/15	0.94	0.09	62,74,83,88	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.










Electron density around Chain G: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray mF_o - DF_c (at 3 rmsd) in purple (negative) and green (positive) Electron density around Chain H: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)





6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B\text{-}factors}({f \AA}^2)$	Q<0.9
6	9Z0	A	1104	14/14	0.49	0.50	72,93,125,130	14
6	9Z0	D	305	14/14	0.51	0.43	92,103,159,160	14
6	9Z0	E	302	14/14	0.57	0.83	115,161,214,246	14
6	9Z0	D	304	14/14	0.62	0.52	69,94,106,125	14
5	L0B	В	308	25/25	0.66	0.51	82,108,123,126	0
6	9Z0	В	307	14/14	0.68	0.55	92,112,136,174	14
6	9 Z 0	В	306	14/14	0.69	0.42	64,91,107,118	14
5	L0B	D	306	25/25	0.69	0.48	$92,\!110,\!118,\!120$	0
5	L0B	A	1106	25/25	0.71	0.57	84,104,114,119	0
6	9Z0	Ε	301	14/14	0.71	0.39	$60,\!95,\!115,\!115$	14
6	9Z0	$^{\mathrm{C}}$	304	14/14	0.73	0.42	$70,\!87,\!101,\!117$	14
5	L0B	С	306	25/25	0.74	0.59	82,102,111,120	0
5	L0B	A	1101	25/25	0.75	0.40	77,98,116,118	0
6	9Z0	A	1105	14/14	0.77	0.38	81,94,117,125	14


Continued on next page...



 $Continued\ from\ previous\ page...$

Mol	Type	Chain	${f Res}$	Atoms	RSCC	RSR	${f B-factors}({f A}^2)$	Q < 0.9
6	9Z0	С	305	14/14	0.82	0.33	72,79,105,121	14
7	DMS	E	303	4/4	0.91	0.53	110,112,113,123	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around 9Z0 D 305:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around 9Z0 E 302:











Electron density around 9Z0 D 304: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around L0B B 308:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around 9Z0 B 307:












Electron density around 9Z0 B 306:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around L0B D 306:












Electron density around L0B A 1106:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around 9Z0 E 301:












Electron density around 9Z0 C 304:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around L0B C 306:












Electron density around L0B A 1101:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around 9Z0 A 1105:











6.5 Other polymers (i)

There are no such residues in this entry.

