



# Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 02:58 am BST

PDB ID : 3OW4  
Title : Discovery of dihydrothieno- and dihydrofuropyrimidines as potent pan Akt inhibitors  
Authors : Dizon, F.; Wu, W.; Vigers, G.P.A.; Brandhuber, B.J.  
Deposited on : 2010-09-17  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	341	
1	B	341	
2	C	10	
2	D	10	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5597 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RAC-alpha serine/threonine-protein kinase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	318	2604	1672	438	478	1	15	0	0	0
1	B	318	2604	1672	438	478	1	15	0	0	0


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	140	GLY	-	EXPRESSION TAG	UNP P31749
A	141	ALA	-	EXPRESSION TAG	UNP P31749
A	142	MET	-	EXPRESSION TAG	UNP P31749
A	143	ALA	-	EXPRESSION TAG	UNP P31749
A	473	ASP	SER	ENGINEERED MUTATION	UNP P31749
A	478	SER	GLY	SEE REMARK 999	UNP P31749
B	140	GLY	-	EXPRESSION TAG	UNP P31749
B	141	ALA	-	EXPRESSION TAG	UNP P31749
B	142	MET	-	EXPRESSION TAG	UNP P31749
B	143	ALA	-	EXPRESSION TAG	UNP P31749
B	473	ASP	SER	ENGINEERED MUTATION	UNP P31749
B	478	SER	GLY	SEE REMARK 999	UNP P31749

- Molecule 2 is a protein called GSK 3 beta peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	10	79	47	16	16	0	0	0
2	D	10	79	47	16	16	0	0	0

- Molecule 3 is (2R)-3-(1H-indol-3-yl)-1-{4-[(5S)-5-methyl-5,7-dihydrothieno[3,4-d]pyrimidin-4-yl]piperazin-1-yl}-1-oxopropan-2-amine (three-letter code: SMY) (formula: C<sub>22</sub>H<sub>26</sub>N<sub>6</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	A	1	30	22	6	1	1	0	0
3	B	1	30	22	6	1	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	95	95	95	0	0
4	B	72	72	72	0	0
4	C	3	3	3	0	0
4	D	1	1	1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RAC-alpha serine/threonine-protein kinase



- Molecule 1: RAC-alpha serine/threonine-protein kinase



- Molecule 2: GSK 3 beta peptide



- Molecule 2: GSK 3 beta peptide



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	86.43Å 55.82Å 91.35Å 90.00° 103.93° 90.00°	Depositor
Resolution (Å)	27.47 – 2.60 27.47 – 2.60	Depositor EDS
% Data completeness (in resolution range)	86.3 (27.47-2.60) 86.4 (27.47-2.60)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	8.47 (at 2.61Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.193 , 0.268 0.191 , 0.262	Depositor DCC
$R_{free}$ test set	1110 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.3	Xtrriage
Anisotropy	0.088	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 49.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5597	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 56.98 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.5306e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, SMY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.72	1/2654 (0.0%)	0.79	0/3568
1	B	0.70	0/2654	0.77	0/3568
2	C	0.78	0/80	0.89	0/105
2	D	0.67	0/80	0.65	0/105
All	All	0.71	1/5468 (0.0%)	0.78	0/7346

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	364	GLU	CG-CD	5.76	1.60	1.51

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2604	0	2568	39	0
1	B	2604	0	2568	41	0
2	C	79	0	77	6	0
2	D	79	0	77	6	0
3	A	30	0	26	5	0
3	B	30	0	26	4	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	95	0	0	3	0
4	B	72	0	0	1	0
4	C	3	0	0	0	0
4	D	1	0	0	0	0
All	All	5597	0	5342	89	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:413:TRP:H	1:B:352:GLN:HE22	1.17	0.88
1:B:229:TYR:CE2	1:B:231:ASN:HB3	2.09	0.87
1:A:207:HIS:HD2	1:A:209:PHE:H	1.31	0.77
1:A:428:GLN:HE21	1:A:428:GLN:HA	1.49	0.75
2:C:2:ARG:CG	2:C:2:ARG:HH11	2.02	0.73
1:B:339:MET:HA	1:B:342:MET:HE2	1.70	0.72
1:A:231:ASN:H	1:A:231:ASN:ND2	1.87	0.71
3:A:1:SMY:H12	3:A:1:SMY:H9	1.75	0.69
1:B:366:ILE:HD13	1:B:368:PHE:CZ	2.29	0.67
1:A:339:MET:HA	1:A:342:MET:HE2	1.77	0.67
1:A:413:TRP:N	1:B:352:GLN:HE22	1.92	0.66
1:B:239:LEU:HD23	1:B:277:LEU:HD11	1.79	0.65
1:A:231:ASN:ND2	1:A:284:LYS:HG2	2.13	0.64
3:B:2:SMY:H9	3:B:2:SMY:H12	1.79	0.63
1:B:372:LEU:HD21	1:B:380:LEU:HD12	1.80	0.62
2:C:2:ARG:HG2	2:C:2:ARG:HH11	1.66	0.61
1:A:428:GLN:HE21	1:A:428:GLN:CA	2.14	0.61
1:B:207:HIS:HD2	1:B:209:PHE:H	1.49	0.60
1:A:428:GLN:O	1:A:436:ARG:NE	2.33	0.60
1:B:231:ASN:HD21	1:B:284:LYS:HB3	1.65	0.60
1:B:397:GLU:H	1:B:397:GLU:CD	2.03	0.60
1:A:165:ILE:HD13	1:A:180:ILE:CD1	2.33	0.59
1:B:368:PHE:CZ	1:B:381:SER:HB3	2.37	0.59
1:A:250:ALA:HB1	1:A:342:MET:HE3	1.85	0.59
1:A:428:GLN:NE2	1:A:428:GLN:HA	2.18	0.58
1:B:165:ILE:HD12	1:B:180:ILE:HD12	1.86	0.58
1:A:250:ALA:CB	1:A:342:MET:HE3	2.33	0.57
1:B:368:PHE:HZ	1:B:381:SER:HB3	1.68	0.57
1:A:231:ASN:H	1:A:231:ASN:HD22	1.52	0.57

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:375:GLU:HG2	4:A:4:HOH:O	2.04	0.56
1:B:194:HIS:HE1	2:D:9:ALA:HB3	1.71	0.55
3:A:1:SMY:H12	3:A:1:SMY:C9	2.36	0.55
1:A:165:ILE:HD13	1:A:180:ILE:HD11	1.88	0.55
1:B:466:ARG:N	1:B:467:PRO:HD3	2.23	0.54
1:B:234:GLU:OE1	2:D:4:ARG:NH2	2.40	0.53
1:A:434:ASP:OD1	1:A:436:ARG:HG3	2.08	0.53
1:B:207:HIS:HB3	1:B:210:LEU:HG	1.91	0.53
1:B:231:ASN:HD21	1:B:284:LYS:CB	2.24	0.51
1:A:392:LEU:HD12	1:A:401:GLU:HG2	1.93	0.51
3:B:2:SMY:H12	3:B:2:SMY:C9	2.42	0.50
1:A:234:GLU:OE2	3:A:1:SMY:N30	2.45	0.50
1:B:366:ILE:HD13	1:B:368:PHE:HZ	1.77	0.49
1:B:207:HIS:CD2	1:B:209:PHE:H	2.29	0.49
1:A:397:GLU:CD	1:A:397:GLU:H	2.15	0.49
1:B:158:LYS:O	3:B:2:SMY:H29	2.11	0.49
1:B:194:HIS:HE1	2:D:9:ALA:CB	2.26	0.49
4:A:483:HOH:O	1:B:353:ASP:HB2	2.14	0.48
1:A:205:SER:HA	1:A:263:TYR:OH	2.14	0.48
1:A:216:SER:O	1:A:473:ASP:HA	2.14	0.48
1:B:234:GLU:CD	2:D:4:ARG:HH22	2.16	0.48
3:A:1:SMY:HN3A	2:C:4:ARG:HH21	1.62	0.47
1:B:231:ASN:HD21	1:B:284:LYS:HA	1.79	0.47
1:B:210:LEU:HD21	1:B:260:ALA:HB1	1.97	0.47
1:B:183:LYS:HE2	1:B:469:PHE:CE1	2.50	0.46
1:A:301:LYS:O	1:A:302:ASP:C	2.54	0.46
1:B:185:VAL:O	1:B:189:LYS:HB2	2.16	0.46
1:B:258:VAL:HG22	1:B:335:LEU:HD21	1.98	0.46
1:A:366:ILE:C	1:A:367:ARG:HG2	2.32	0.46
1:A:366:ILE:H	1:A:366:ILE:HG13	1.41	0.46
1:B:195:THR:HG21	1:B:223:LEU:HD13	1.98	0.46
3:B:2:SMY:N30	2:D:4:ARG:NH2	2.64	0.46
1:B:231:ASN:HD21	1:B:284:LYS:CA	2.29	0.45
1:B:417:TYR:CD2	1:B:417:TYR:C	2.89	0.45
1:B:318:PRO:HG2	1:B:386:LYS:HA	1.99	0.45
1:A:271:VAL:HG12	1:A:273:ARG:HG3	1.99	0.45
3:A:1:SMY:H13A	3:A:1:SMY:H19	1.74	0.44
1:A:312:THR:CG2	2:C:5:THR:HB	2.47	0.44
1:B:436:ARG:HD3	4:B:482:HOH:O	2.18	0.44
1:A:275:LEU:HD21	1:A:280:LEU:HD21	2.00	0.44
1:A:165:ILE:HD13	1:A:180:ILE:HD12	2.00	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:207:HIS:CG	1:B:208:PRO:HD2	2.54	0.43
1:A:275:LEU:HB2	4:A:53:HOH:O	2.19	0.43
1:B:223:LEU:HD21	1:B:469:PHE:CD1	2.54	0.42
1:A:301:LYS:O	1:A:302:ASP:O	2.37	0.42
1:A:219:THR:OG1	1:A:222:ARG:HB2	2.19	0.42
1:A:318:PRO:HG2	1:A:386:LYS:HA	2.01	0.42
1:B:154:LYS:O	1:B:165:ILE:HG23	2.19	0.42
1:A:180:ILE:HG23	1:A:224:CYS:SG	2.60	0.42
1:B:301:LYS:HE3	1:B:301:LYS:HB2	1.93	0.42
1:A:297:LYS:HG2	1:A:306:MET:HG2	2.01	0.42
1:A:213:LEU:HA	1:A:227:MET:HG2	2.03	0.41
1:A:239:LEU:HD13	1:A:239:LEU:HA	1.94	0.41
1:B:361:ILE:O	1:B:386:LYS:HD3	2.21	0.41
1:A:153:LEU:HD11	1:A:168:LYS:HB2	2.01	0.41
2:C:2:ARG:HG3	2:C:2:ARG:HH11	1.79	0.41
1:B:309:PHE:CE1	2:D:8:PHE:CZ	3.09	0.41
1:B:231:ASN:ND2	1:B:284:LYS:HB3	2.34	0.41
1:A:235:LEU:HA	1:A:235:LEU:HD12	1.89	0.40
2:C:2:ARG:HG2	2:C:2:ARG:NH1	2.34	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/341 (92%)	295 (94%)	16 (5%)	2 (1%)	25	47
1	B	313/341 (92%)	295 (94%)	17 (5%)	1 (0%)	41	64
2	C	8/10 (80%)	8 (100%)	0	0	100	100
2	D	8/10 (80%)	8 (100%)	0	0	100	100
All	All	642/702 (92%)	606 (94%)	33 (5%)	3 (0%)	29	52

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	302	ASP
1	A	240	SER
1	B	430	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	278/297 (94%)	254 (91%)	24 (9%)	10	20
1	B	278/297 (94%)	256 (92%)	22 (8%)	12	24
2	C	8/8 (100%)	7 (88%)	1 (12%)	4	8
2	D	8/8 (100%)	6 (75%)	2 (25%)	0	1
All	All	572/610 (94%)	523 (91%)	49 (9%)	10	20

All (49) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	144	ARG
1	A	146	THR
1	A	148	ASN
1	A	154	LYS
1	A	158	LYS
1	A	182	LYS
1	A	206	ARG
1	A	216	SER
1	A	220	HIS
1	A	231	ASN
1	A	239	LEU
1	A	275	LEU
1	A	284	LYS
1	A	297	LYS
1	A	301	LYS
1	A	346	ARG
1	A	365	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	366	ILE
1	A	367	ARG
1	A	375	GLU
1	A	397	GLU
1	A	404	GLN
1	A	428	GLN
1	A	466	ARG
1	B	149	GLU
1	B	184	GLU
1	B	206	ARG
1	B	228	GLU
1	B	231	ASN
1	B	236	PHE
1	B	275	LEU
1	B	297	LYS
1	B	301	LYS
1	B	324	ASN
1	B	366	ILE
1	B	378	SER
1	B	397	GLU
1	B	402	ILE
1	B	406	ARG
1	B	421	LEU
1	B	426	LYS
1	B	436	ARG
1	B	466	ARG
1	B	473	ASP
1	B	474	TYR
1	B	478	SER
2	C	2	ARG
2	D	4	ARG
2	D	6	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	207	HIS
1	A	231	ASN
1	A	324	ASN
1	A	414	GLN
1	A	428	GLN
1	B	194	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	199	ASN
1	B	203	GLN
1	B	207	HIS
1	B	220	HIS
1	B	231	ASN
1	B	324	ASN
1	B	352	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	TPO	B	308	1	8,10,11	0.77	0	10,14,16	1.26	1 (10%)
1	TPO	A	308	1	8,10,11	0.95	0	10,14,16	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	B	308	1	-	0/9/11/13	-
1	TPO	A	308	1	-	1/9/11/13	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	308	TPO	O2P-P-O1P	2.08	118.84	110.68

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	308	TPO	CB-OG1-P-O1P

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SMY	B	2	-	31,34,34	0.99	3 (9%)	37,49,49	2.07	15 (40%)
3	SMY	A	1	-	31,34,34	1.02	3 (9%)	37,49,49	1.70	7 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SMY	B	2	-	-	1/15/35/35	0/5/5/5
3	SMY	A	1	-	-	2/15/35/35	0/5/5/5

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1	SMY	C22-N23	2.78	1.42	1.36
3	A	1	SMY	C27-C26	2.41	1.42	1.36
3	B	2	SMY	C22-N23	2.33	1.41	1.36
3	B	2	SMY	C27-C26	2.27	1.41	1.36
3	B	2	SMY	C28-C29	2.26	1.41	1.36
3	A	1	SMY	C28-C29	2.15	1.41	1.36

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	2	SMY	C4-C5-C9	4.65	116.25	110.36
3	A	1	SMY	C4-C5-C9	4.46	116.00	110.36
3	A	1	SMY	C5-C4-N3	-3.76	120.76	125.41
3	B	2	SMY	O18-C17-N14	-3.44	117.64	121.67
3	B	2	SMY	C5-C4-N3	-3.38	121.23	125.41
3	B	2	SMY	C20-C21-C25	3.26	131.32	126.25
3	B	2	SMY	C15-C16-N11	3.24	116.99	110.70
3	B	2	SMY	C20-C19-C17	3.17	116.86	109.27
3	A	1	SMY	C7-S8-C9	2.95	97.66	93.20
3	B	2	SMY	C28-C29-C25	-2.94	116.81	120.89
3	B	2	SMY	C29-C25-C21	-2.94	129.02	134.42
3	A	1	SMY	C29-C25-C21	-2.94	129.02	134.42
3	B	2	SMY	C7-S8-C9	2.90	97.59	93.20
3	B	2	SMY	C27-C26-C24	-2.79	116.07	120.08
3	A	1	SMY	O18-C17-C19	-2.75	114.65	119.66
3	B	2	SMY	C16-C15-N14	2.74	116.30	110.44
3	B	2	SMY	C19-C17-N14	2.73	123.00	118.87
3	A	1	SMY	C29-C25-C24	2.66	121.70	118.17
3	B	2	SMY	C2-N1-C6	2.60	117.88	111.75
3	B	2	SMY	C29-C25-C24	2.51	121.50	118.17
3	A	1	SMY	C2-N1-C6	2.35	117.29	111.75
3	B	2	SMY	N3-C2-N1	-2.13	125.28	128.60

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	2	SMY	C19-C17-N14-C13
3	A	1	SMY	O18-C17-N14-C15
3	A	1	SMY	C19-C17-N14-C13





There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	2	SMY	4	0
3	A	1	SMY	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	317/341 (92%)	-0.23	7 (2%) 62 56	20, 42, 72, 90	0
1	B	317/341 (92%)	-0.18	9 (2%) 53 46	21, 45, 78, 95	0
2	C	10/10 (100%)	0.22	0 100 100	59, 61, 65, 69	0
2	D	10/10 (100%)	1.41	2 (20%) 1 0	75, 78, 83, 85	0
All	All	654/702 (93%)	-0.17	18 (2%) 53 46	20, 45, 77, 95	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	220	HIS	4.9
1	A	444	ALA	4.4
1	A	478	SER	4.0
1	B	478	SER	4.0
1	B	446	MET	3.7
2	D	1	GLY	3.3
1	B	220	HIS	3.3
1	A	477	SER	3.1
1	A	304	ALA	3.0
1	B	184	GLU	2.8
1	A	144	ARG	2.8
1	B	440	GLU	2.5
1	B	444	ALA	2.3
1	B	221	ASP	2.3
1	A	446	MET	2.2
1	B	148	ASN	2.1
2	D	3	PRO	2.1
1	B	302	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	TPO	A	308	11/12	0.96	0.09	39,41,50,51	0
1	TPO	B	308	11/12	0.98	0.09	38,38,41,42	0

## 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SMY	B	2	30/30	0.93	0.18	41,47,52,54	0
3	SMY	A	1	30/30	0.93	0.18	38,46,49,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.