

# wwPDB X-ray Structure Validation Summary Report (i)

Oct 31, 2023 – 03:24 pm GMT

PDB ID : 8OW7

Title : Crystal structure of Tannerella forsythia sugar kinase K1058 in complex with

N-acetylmuramic acid (MurNAc)

Authors: Stasiak, A.C.; Gogler, K.; Fink, P.; Stehle, T.; Zocher, G.

Deposited on : 2023-04-27

Resolution : 3.06 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 3.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Percentile relative to X-ray structures of similar resolution

Metric	Whole archive	Similar resolution
Metric	$(\#  ext{Entries})$	$(\#  ext{Entries},  ext{ resolution range}( ext{Å}))$
$R_{free}$	130704	1754 (3.10-3.02)
Clashscore	141614	1864 (3.10-3.02)
Ramachandran outliers	138981	1794 (3.10-3.02)
Sidechain outliers	138945	1793 (3.10-3.02)
RSRZ outliers	127900	1713 (3.10-3.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	293	85%	10%	5%
1	В	293	83%	11%	5%
1	С	293	85%	10%	5%
1	D	293	86%	8%	5%
1	Е	293	84%	11%	5%



Continued from previous page...

Mol	Chain	Length	Quality of chain		
	1	202	3%		
1	F'	293	82%	12%	5%



## 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 13276 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called N-acetylglucosamine kinase.

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	Trace
1	A	277	Total	С	N	О	S	0	0	0
1	A	211	2159	1396	355	401	7	0	U	
1	В	277	Total	С	N	О	S	0	0	0
1	D	211	2184	1411	363	403	7		0	
1	С	277	Total	С	N	О	S	0	0	0
1		211	2190	1414	366	403	7		0	
1	D	277	Total	С	N	О	S	0	0	0
1	D	211	2190	1414	366	403	7	U	U	0
1	Е	277	Total	С	N	О	S	0	0	0
1	12	211	2184	1411	363	403	7		0	
1	F	277	Total	С	N	О	S	0	0	0
1	I.	211	2184	1411	363	403	7		U	

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual Comment		Reference
A	0	MET	-	initiating methionine	UNP G8UQH1
A	1	ALA	-	expression tag	UNP G8UQH1
A	285	LEU	-	expression tag	UNP G8UQH1
A	286	GLU	-	expression tag	UNP G8UQH1
A	287	HIS	-	expression tag	UNP G8UQH1
A	288	HIS	-	expression tag	UNP G8UQH1
A	289	HIS	-	expression tag	UNP G8UQH1
A	290	HIS	-	expression tag	UNP G8UQH1
A	291	HIS	-	expression tag	UNP G8UQH1
A	292	HIS	-	expression tag	UNP G8UQH1
В	0	MET	-	initiating methionine	UNP G8UQH1
В	1	ALA	-	expression tag	UNP G8UQH1
В	285	LEU	-	expression tag	UNP G8UQH1
В	286	GLU		expression tag	UNP G8UQH1
В	287	HIS	-	expression tag	UNP G8UQH1
В	288	HIS	-	expression tag	UNP G8UQH1
В	289	HIS	-	expression tag	UNP G8UQH1



 $Continued\ from\ previous\ page...$ 


Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
В	290	HIS	-	expression tag	UNP G8UQH1
В	291	HIS	-	expression tag	UNP G8UQH1
В	292	HIS	-	expression tag	UNP G8UQH1
С	0	MET	_	initiating methionine	UNP G8UQH1
С	1	ALA	-	expression tag	UNP G8UQH1
С	285	LEU	-	expression tag	UNP G8UQH1
С	286	GLU	-	expression tag	UNP G8UQH1
С	287	HIS	-	expression tag	UNP G8UQH1
С	288	HIS	_	expression tag	UNP G8UQH1
С	289	HIS	-	expression tag	UNP G8UQH1
С	290	HIS	_	expression tag	UNP G8UQH1
С	291	HIS	-	expression tag	UNP G8UQH1
С	292	HIS	-	expression tag	UNP G8UQH1
D	0	MET	_	initiating methionine	UNP G8UQH1
D	1	ALA	-	expression tag	UNP G8UQH1
D	285	LEU	_	expression tag	UNP G8UQH1
D	286	GLU	-	expression tag	UNP G8UQH1
D	287	HIS	-	expression tag	UNP G8UQH1
D	288	HIS	-	expression tag	UNP G8UQH1
D	289	HIS	-	expression tag	UNP G8UQH1
D	290	HIS	-	expression tag	UNP G8UQH1
D	291	HIS	-	expression tag	UNP G8UQH1
D	292	HIS	-	expression tag	UNP G8UQH1
Е	0	MET	-	initiating methionine	UNP G8UQH1
Е	1	ALA	-	expression tag	UNP G8UQH1
Е	285	LEU	-	expression tag	UNP G8UQH1
Е	286	GLU	-	expression tag	UNP G8UQH1
Е	287	HIS	-	expression tag	UNP G8UQH1
Е	288	HIS	-	expression tag	UNP G8UQH1
Е	289	HIS	_	expression tag	UNP G8UQH1
Е	290	HIS	-	expression tag	UNP G8UQH1
Е	291	HIS	_	expression tag	UNP G8UQH1
Е	292	HIS	-	expression tag	UNP G8UQH1
F	0	MET	_	initiating methionine	UNP G8UQH1
F	1	ALA	-	expression tag	UNP G8UQH1
F	285	LEU	-	expression tag	UNP G8UQH1
F	286	GLU	-	expression tag	UNP G8UQH1
F	287	HIS	-	expression tag	UNP G8UQH1
F	288	HIS	-	expression tag	UNP G8UQH1
F	289	HIS	-	expression tag	UNP G8UQH1
F	290	HIS	-	expression tag	UNP G8UQH1
F	291	HIS	-	expression tag	UNP G8UQH1



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference	
F	292	HIS	-	expression tag	UNP G8UQH1	


• Molecule 2 is N-acetyl-beta-muramic acid (three-letter code: AMU) (formula:  $C_{11}H_{19}NO_8$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	С	N	О	0	0
	A	1	20	11	1	8	U	U
2	В	1	Total	С	N	О	0	0
	Ъ	1	20	11	1	8	U	U
2	С	1	Total	С	N	О	0	0
		1	20	11	1	8	U	U
2	D	1	Total	С	N	О	0	0
	D	1	20	11	1	8	U	U
2	E	1	Total	С	N	О	0	0
	Ľ	1	20	11	1	8	U	U
2	F	1	Total	С	N	О	0	0
	I.	1	20	11	1	8	U	U

• Molecule 3 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).






Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	О	S	0	0
3	A	1	5	4	1	0	U
3	A	1	Total	О	S	0	0
	Λ	1	5	4	1	U	U
3	A	1	Total	Ο	S	0	0
	11	1	5	4	1	0	0
3	A	1	Total	Ο	S	0	0
	71	1	5	4	1	0	U
3	В	1	Total	Ο	S	0	0
	Ъ	1	5	4	1	· ·	<u> </u>
3	В	1	Total	Ο	S	0	0
	Б	1	5	4	1	· ·	<u> </u>
3	В	1	Total	Ο	S	0	0
		1	5	4	1	Ü	Ü
3	С	1	Total	Ο	S	0	0
			5	4	1	Ü	U
3	С	1	Total	Ο	S	0	0
		-	5	4	1	Ŭ	Ü
3	D	1	Total	Ο	S	0	0
		1	5	4	1	Ü	Ü
3	E	1	Total	O	S	0	0
		1	5	4	1		
3	F	1	Total	O	S	0	0
	*		5	4	1	Ŭ	
3	F	1	Total	O	S	0	0
	•	<u> </u>	5	4	1		




## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: N-acetylglucosamine kinase




• Molecule 1: N-acetylglucosamine kinase



• Molecule 1: N-acetylglucosamine kinase




• Molecule 1: N-acetylglucosamine kinase








 $\bullet$  Molecule 1: N-acetylglucosamine kinase



• Molecule 1: N-acetylglucosamine kinase





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants	143.75Å 143.75Å 210.95Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 120.00°	
Resolution (Å)	29.85 - 3.06	Depositor
. ,	29.85 - 3.06	EDS
% Data completeness	99.9 (29.85-3.06)	Depositor
(in resolution range)	100.0 (29.85-3.06)	EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.18 (at 3.06Å)	Xtriage
Refinement program	REFMAC 5.8.0352	Depositor
D D	0.219 , 0.245	Depositor
$R, R_{free}$	0.219 , $0.245$	DCC
$R_{free}$ test set	1444 reflections (3.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	74.3	Xtriage
Anisotropy	0.134	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.32, 34.0	EDS
L-test for twinning <sup>2</sup>	$< L > = 0.48, < L^2> = 0.31$	Xtriage
Estimated twinning fraction	0.022 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	13276	wwPDB-VP
Average B, all atoms $(\mathring{A}^2)$	87.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.33% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

#### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: AMU, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
IVIOI	Chain	$\mid \text{RMSZ} \mid \# Z  > 5$		RMSZ	# Z  > 5
1	A	0.26	0/2209	0.48	0/2993
1	В	0.26	0/2234	0.48	0/3020
1	С	0.26	0/2240	0.48	0/3027
1	D	0.26	0/2240	0.48	0/3027
1	Е	0.26	0/2234	0.48	0/3020
1	F	0.26	0/2234	0.48	0/3020
All	All	0.26	0/13391	0.48	0/18107

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintenain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	В	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	100	ARG	Sidechain
1	В	46	ARG	Sidechain



#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2159	0	2097	17	0
1	В	2184	0	2157	21	0
1	С	2190	0	2168	15	0
1	D	2190	0	2168	14	0
1	Ε	2184	0	2157	16	0
1	F	2184	0	2157	21	0
2	A	20	0	18	0	0
2	В	20	0	18	0	0
2	С	20	0	18	0	0
2	D	20	0	18	0	0
2	Ε	20	0	18	0	0
2	F	20	0	18	0	0
3	A	20	0	0	0	0
3	В	15	0	0	0	0
3	С	10	0	0	0	0
3	D	5	0	0	0	0
3	Е	5	0	0	0	0
3	F	10	0	0	0	0
All	All	13276	0	13012	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 97 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)	
1:A:95:LEU:HD21	1:A:120:CYS:HB3	1.74	0.69	
1:D:7:SER:HB2	1:D:12:THR:HG23	1.79	0.64	
1:C:95:LEU:HD21	1:C:120:CYS:HB3	1.81	0.63	
1:D:182:VAL:HA	1:D:186:ARG:HB3	1.80	0.63	
1:F:238:VAL:HG12	1:F:264:GLU:HB2	1.82	0.61	

There are no symmetry-related clashes.



#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	275/293 (94%)	267 (97%)	6 (2%)	2 (1%)	22	52
1	В	275/293 (94%)	266 (97%)	8 (3%)	1 (0%)	34	64
1	С	275/293 (94%)	265 (96%)	8 (3%)	2 (1%)	22	52
1	D	275/293 (94%)	260 (94%)	14 (5%)	1 (0%)	34	64
1	E	275/293 (94%)	261 (95%)	13 (5%)	1 (0%)	34	64
1	F	275/293 (94%)	262 (95%)	12 (4%)	1 (0%)	34	64
All	All	1650/1758 (94%)	1581 (96%)	61 (4%)	8 (0%)	29	60

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	9	SER
1	В	9	SER
1	Е	104	GLN
1	С	9	SER
1	С	104	GLN

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	$225/250\ (90\%)$	225 (100%)	0	100 100
1	В	232/250~(93%)	232 (100%)	0	100 100
1	С	233/250 (93%)	232 (100%)	1 (0%)	91 95



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	D	233/250 (93%)	232 (100%)	1 (0%)	91	95	
1	Е	232/250 (93%)	229 (99%)	3 (1%)	69	86	
1	F	232/250 (93%)	231 (100%)	1 (0%)	91	95	
All	All	1387/1500 (92%)	1381 (100%)	6 (0%)	91	95	

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Е	204	LEU
1	Е	206	LEU
1	F	38	ARG
1	D	268	MET
1	С	181	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	244	ASN
1	В	104	GLN
1	В	250	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

19 ligands are modelled in this entry.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Во	ond leng	ths	В	ond ang	les
IVIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	В	302	_	4,4,4	0.39	0	6,6,6	0.05	0
3	SO4	Е	302	_	4,4,4	0.39	0	6,6,6	0.06	0
3	SO4	A	303	_	4,4,4	0.39	0	6,6,6	0.05	0
3	SO4	D	302	-	4,4,4	0.38	0	6,6,6	0.05	0
3	SO4	A	305	_	4,4,4	0.39	0	6,6,6	0.05	0
3	SO4	В	304	-	4,4,4	0.38	0	6,6,6	0.05	0
3	SO4	A	302	-	4,4,4	0.39	0	6,6,6	0.05	0
2	AMU	F	301	_	20,20,20	0.53	0	24,28,28	0.77	1 (4%)
3	SO4	В	303	-	4,4,4	0.37	0	6,6,6	0.05	0
3	SO4	С	303	-	4,4,4	0.38	0	6,6,6	0.05	0
3	SO4	С	302	_	4,4,4	0.39	0	6,6,6	0.05	0
2	AMU	С	301	_	20,20,20	0.50	0	24,28,28	0.71	0
3	SO4	A	304	_	4,4,4	0.39	0	6,6,6	0.05	0
2	AMU	E	301	_	20,20,20	0.54	0	24,28,28	0.75	1 (4%)
2	AMU	A	301	-	20,20,20	0.50	0	24,28,28	0.64	0
3	SO4	F	302	-	4,4,4	0.38	0	6,6,6	0.05	0
2	AMU	D	301	-	20,20,20	0.50	0	24,28,28	0.77	1 (4%)
3	SO4	F	303	-	4,4,4	0.39	0	6,6,6	0.05	0
2	AMU	В	301	-	20,20,20	0.53	0	24,28,28	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AMU	Е	301	-	-	0/14/34/34	0/1/1/1
2	AMU	A	301	-	-	4/14/34/34	0/1/1/1
2	AMU	С	301	-	-	2/14/34/34	0/1/1/1
2	AMU	F	301	-	-	2/14/34/34	0/1/1/1
2	AMU	D	301	-	-	0/14/34/34	0/1/1/1
2	AMU	В	301	-	-	0/14/34/34	0/1/1/1

There are no bond length outliers.



All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
2	Ε	301	AMU	C1-C2-N2	2.17	113.24	110.73
2	F	301	AMU	C1-C2-N2	2.17	113.24	110.73
2	D	301	AMU	C1-C2-N2	2.11	113.17	110.73

There are no chirality outliers.

5 of 8 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	С	301	AMU	C8-C7-N2-C2
2	A	301	AMU	O11-C10-C9-O3
2	A	301	AMU	C8-C7-N2-C2
2	С	301	AMU	O7-C7-N2-C2
2	A	301	AMU	O7-C7-N2-C2

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





























## 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$\mathrm{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	277/293 (94%)	-0.31	0 100 100	51, 73, 99, 113	0
1	В	277/293 (94%)	-0.46	0 100 100	46, 66, 91, 104	0
1	С	277/293 (94%)	-0.30	0 100 100	48, 69, 102, 124	0
1	D	277/293 (94%)	0.30	13 (4%) 31 14	73, 109, 157, 171	0
1	E	277/293 (94%)	0.08	6 (2%) 62 38	65, 101, 142, 155	0
1	F	277/293 (94%)	0.01	9 (3%) 47 24	56, 89, 127, 151	0
All	All	$1662/1758 \ (94\%)$	-0.11	28 (1%) 70 46	46, 82, 136, 171	0

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	108	GLY	6.5
1	D	229	ASP	3.6
1	Е	229	ASP	3.5
1	Е	261	SER	3.2
1	D	125	HIS	3.2

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q<0.9
3	SO4	F	302	5/5	0.83	0.35	159,163,164,168	0
3	SO4	В	303	5/5	0.84	0.26	130,136,138,140	0
3	SO4	D	302	5/5	0.89	0.26	97,101,107,110	0
2	AMU	D	301	20/20	0.89	0.27	87,106,111,115	0
2	AMU	Е	301	20/20	0.90	0.29	72,77,83,87	0
3	SO4	A	305	5/5	0.91	0.38	125,126,128,129	0
3	SO4	A	304	5/5	0.92	0.28	141,144,145,146	0
3	SO4	Ε	302	5/5	0.93	0.16	83,87,88,89	0
2	AMU	F	301	20/20	0.93	0.28	62,78,82,88	0
3	SO4	A	303	5/5	0.94	0.17	109,111,114,115	0
2	AMU	A	301	20/20	0.94	0.23	52,62,65,71	0
3	SO4	С	303	5/5	0.94	0.13	102,102,108,108	0
3	SO4	F	303	5/5	0.94	0.28	89,93,96,99	0
2	AMU	В	301	20/20	0.95	0.22	54,56,59,64	0
2	AMU	С	301	20/20	0.95	0.20	51,56,65,69	0
3	SO4	В	302	5/5	0.96	0.42	102,106,106,107	0
3	SO4	В	304	5/5	0.97	0.16	55,58,60,60	0
3	SO4	A	302	5/5	0.98	0.17	72,73,75,78	0
3	SO4	С	302	5/5	0.99	0.20	72,74,78,80	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.
























### Electron density around AMU B 301:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)











### 6.5 Other polymers (i)

There are no such residues in this entry.

