



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 15, 2024 – 05:20 PM EDT

PDB ID : 4OYU  
Title : Crystal structure of the N-terminal domains of muskelin  
Authors : Delto, C.; Kuper, J.; Schindelin, H.  
Deposited on : 2014-02-13  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	196	 5% 75% 15% 9%
1	B	196	 6% 78% 14% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	EDO	A	301[B]	-	-	X	-
2	EDO	B	303	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6522 atoms, of which 3128 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Muskelin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	178	2992	975	1486	249	272	10	0	4	0
1	B	179	3009	979	1496	253	270	11	0	4	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	10	ALA	-	expression tag	UNP Q99PV3
A	11	MET	-	expression tag	UNP Q99PV3
B	10	ALA	-	expression tag	UNP Q99PV3
B	11	MET	-	expression tag	UNP Q99PV3

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	1
			10	2	6	2		
2	A	1	Total	C	H	O	0	1
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	1
			10	2	6	2		
2	B	1	Total	C	H	O	0	1
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	B	1	10	2	6	2	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	A	1	14	3	8	3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	145	Total	O	0	5
			150	150		
4	B	125	Total	O	0	2
			127	127		



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.06Å 65.30Å 101.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.80 – 1.80 33.80 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (33.80-1.80) 100.0 (33.80-1.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.56 (at 1.81Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE: 1.8_1069)	Depositor
R, $R_{free}$	0.156 , 0.193 0.164 , 0.198	Depositor DCC
$R_{free}$ test set	1979 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.2	Xtrriage
Anisotropy	0.031	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 56.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.028 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	6522	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.59	0/1562	0.67	0/2113
1	B	0.63	1/1569 (0.1%)	0.73	0/2122
All	All	0.61	1/3131 (0.0%)	0.70	0/4235

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	13	CYS	CB-SG	-7.38	1.69	1.82

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1506	1486	1487	36	0
1	B	1513	1496	1499	31	0
2	A	52	78	78	15	0
2	B	40	60	60	6	0
3	A	6	8	8	0	0
4	A	150	0	0	11	1
4	B	127	0	0	5	1
All	All	3394	3128	3132	67	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:50:GLU:OE1	4:B:401:HOH:O	1.97	0.82
1:B:127:PRO:HD3	1:B:164[A]:CYS:SG	2.30	0.72
1:A:81:VAL:HG21	1:A:143:PHE:CD2	2.27	0.70
1:B:140:GLY:N	1:B:141:PRO:HD3	2.07	0.69
1:A:18:TYR:CD2	2:A:301[B]:EDO:H22	2.29	0.68
1:B:139:TRP:N	1:B:140:GLY:HA3	2.07	0.67
1:B:81:VAL:CG1	1:B:140:GLY:HA2	2.26	0.64
1:B:140:GLY:N	1:B:141:PRO:CD	2.61	0.64
1:A:81:VAL:HG23	1:A:140:GLY:HA3	1.80	0.64
1:A:172:ARG:NH1	4:A:530[B]:HOH:O	2.28	0.63
1:A:132:LYS:HE2	2:A:304:EDO:H11	1.82	0.61
1:B:81:VAL:HG11	1:B:143:PHE:CD2	2.37	0.59
1:A:12:GLU:OE1	4:A:401:HOH:O	2.17	0.57
1:B:41:ASN:ND2	1:B:77:GLU:OE2	2.37	0.57
1:B:81:VAL:HG13	1:B:140:GLY:HA2	1.86	0.56
1:A:15:LEU:HD21	2:A:301[B]:EDO:H11	1.88	0.56
1:B:132:LYS:NZ	2:B:303:EDO:H12	2.22	0.55
1:A:18:TYR:CE2	2:A:301[B]:EDO:H22	2.42	0.54
1:B:142:SER:HB2	1:B:143:PHE:CD1	2.43	0.54
1:A:132:LYS:CE	2:A:304:EDO:H11	2.37	0.54
1:B:81:VAL:HG12	1:B:140:GLY:HA2	1.90	0.54
1:A:124:GLN:HG2	1:A:167:TRP:CD2	2.44	0.53
1:A:22:LYS:HD3	4:A:452:HOH:O	2.09	0.52
1:A:92:GLY:HA2	4:A:470:HOH:O	2.08	0.52
1:B:132:LYS:HZ1	2:B:303:EDO:H12	1.72	0.52
1:A:81:VAL:CG2	1:A:140:GLY:HA3	2.40	0.51
1:B:81:VAL:HB	1:B:143:PHE:HD2	1.76	0.50
1:B:115:PHE:CE2	2:B:310:EDO:H11	2.46	0.50
1:A:86:LYS:HG3	4:A:525:HOH:O	2.10	0.50
1:A:177:ILE:CD1	2:A:308:EDO:H21	2.42	0.50
1:B:140:GLY:O	1:B:141:PRO:C	2.50	0.50
1:B:187:HIS:O	1:B:188:ASN:C	2.51	0.49
1:B:81:VAL:CG1	1:B:143:PHE:CD2	2.96	0.47
1:A:81:VAL:CG2	1:A:143:PHE:CD2	2.97	0.47
1:A:51:SER:OG	2:A:307:EDO:H21	2.15	0.47
1:B:168:TYR:CE2	1:B:172:ARG:HD2	2.49	0.47
1:A:181:LEU:CD2	1:B:173:GLU:HG2	2.45	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:304:EDO:H21	4:B:450:HOH:O	2.16	0.46
1:A:168:TYR:CE2	1:A:172:ARG:HD3	2.50	0.46
1:A:81:VAL:HG21	1:A:143:PHE:CE2	2.51	0.46
1:A:14:ARG:HD2	4:A:439:HOH:O	2.15	0.46
1:A:123:GLU:O	4:A:402:HOH:O	2.21	0.45
1:A:55:PRO:HB2	2:A:313:EDO:H12	1.99	0.45
1:A:98[B]:MET:HB2	2:A:304:EDO:H12	1.98	0.45
1:B:72:THR:HB	1:B:151:GLU:HB3	1.99	0.45
1:B:140:GLY:O	1:B:142:SER:N	2.50	0.45
1:A:56:GLN:HE22	2:A:307:EDO:C2	2.30	0.44
1:A:165:LEU:HD23	4:A:512:HOH:O	2.17	0.44
1:B:43:GLN:OE1	1:B:46[A]:ARG:NH2	2.49	0.43
1:A:38:ASP:OD2	1:A:112:LYS:NZ	2.50	0.43
1:A:187:HIS:N	1:A:187:HIS:CD2	2.83	0.43
1:A:56:GLN:HE22	2:A:307:EDO:C1	2.31	0.43
2:A:313:EDO:H22	4:A:541:HOH:O	2.19	0.43
1:A:18:TYR:CE2	2:A:301[B]:EDO:C2	3.03	0.42
1:A:181:LEU:CD2	1:B:173:GLU:CG	2.97	0.42
1:B:138:SER:C	1:B:140:GLY:HA3	2.39	0.42
1:A:122:ASP:O	4:A:514:HOH:O	2.22	0.42
1:A:177:ILE:HD12	2:A:308:EDO:H21	2.00	0.42
1:B:132:LYS:NZ	2:B:303:EDO:C1	2.83	0.42
1:B:67:ILE:HD11	1:B:164[A]:CYS:SG	2.60	0.42
1:B:132:LYS:HZ1	2:B:303:EDO:C1	2.33	0.41
1:B:81:VAL:HG23	4:B:521:HOH:O	2.20	0.41
1:B:22:LYS:NZ	4:B:469:HOH:O	2.52	0.41
1:B:148:TRP:O	4:B:470:HOH:O	2.22	0.41
1:A:11:MET:N	2:A:314:EDO:C1	2.84	0.41
1:A:93:MET:CE	1:A:165:LEU:HD21	2.51	0.41
1:A:86:LYS:CE	4:A:525:HOH:O	2.68	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:435:HOH:O	4:B:407:HOH:O[2_454]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	180/196 (92%)	173 (96%)	7 (4%)	0	100	100
1	B	181/196 (92%)	173 (96%)	8 (4%)	0	100	100
All	All	361/392 (92%)	346 (96%)	15 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	173/184 (94%)	169 (98%)	4 (2%)	50	37
1	B	173/184 (94%)	173 (100%)	0	100	100
All	All	346/368 (94%)	342 (99%)	4 (1%)	71	65

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	MET
1	A	43	GLN
1	A	84	LEU
1	A	112	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	187	HIS
1	B	187	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

24 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	EDO	B	308	-	3,3,3	0.49	0	2,2,2	0.26	0
2	EDO	A	307	-	3,3,3	0.50	0	2,2,2	0.11	0
2	EDO	A	313	-	3,3,3	0.53	0	2,2,2	0.12	0
2	EDO	A	303	-	3,3,3	0.45	0	2,2,2	0.60	0
2	EDO	B	302[B]	-	3,3,3	0.50	0	2,2,2	0.32	0
2	EDO	A	302[B]	-	3,3,3	0.22	0	2,2,2	0.36	0
2	EDO	A	309	-	3,3,3	0.41	0	2,2,2	0.40	0
2	EDO	B	301[B]	-	3,3,3	0.49	0	2,2,2	0.19	0
3	GOL	A	311	-	5,5,5	0.35	0	5,5,5	0.28	0
2	EDO	B	305	-	3,3,3	0.56	0	2,2,2	0.15	0
2	EDO	A	306	-	3,3,3	0.52	0	2,2,2	0.22	0
2	EDO	A	314	-	3,3,3	0.41	0	2,2,2	0.93	0
2	EDO	A	312	-	3,3,3	0.59	0	2,2,2	0.21	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	EDO	B	304	-	3,3,3	0.56	0	2,2,2	0.19	0
2	EDO	A	304	-	3,3,3	0.50	0	2,2,2	0.10	0
2	EDO	A	301[B]	-	3,3,3	0.52	0	2,2,2	0.33	0
2	EDO	B	310	-	3,3,3	0.46	0	2,2,2	0.24	0
2	EDO	A	305	-	3,3,3	0.45	0	2,2,2	0.41	0
2	EDO	A	308	-	3,3,3	0.52	0	2,2,2	0.23	0
2	EDO	B	306	-	3,3,3	0.36	0	2,2,2	0.46	0
2	EDO	B	307	-	3,3,3	0.58	0	2,2,2	0.25	0
2	EDO	B	303	-	3,3,3	0.44	0	2,2,2	0.62	0
2	EDO	A	310	-	3,3,3	0.51	0	2,2,2	0.42	0
2	EDO	B	309	-	3,3,3	0.46	0	2,2,2	1.01	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	B	308	-	-	1/1/1/1	-
2	EDO	A	307	-	-	1/1/1/1	-
2	EDO	A	313	-	-	1/1/1/1	-
2	EDO	A	303	-	-	0/1/1/1	-
2	EDO	B	302[B]	-	-	0/1/1/1	-
2	EDO	A	302[B]	-	-	0/1/1/1	-
2	EDO	A	309	-	-	0/1/1/1	-
2	EDO	B	301[B]	-	-	0/1/1/1	-
3	GOL	A	311	-	-	4/4/4/4	-
2	EDO	B	305	-	-	0/1/1/1	-
2	EDO	A	306	-	-	0/1/1/1	-
2	EDO	A	314	-	-	1/1/1/1	-
2	EDO	A	312	-	-	1/1/1/1	-
2	EDO	B	304	-	-	1/1/1/1	-
2	EDO	A	304	-	-	1/1/1/1	-
2	EDO	A	301[B]	-	-	1/1/1/1	-
2	EDO	B	310	-	-	0/1/1/1	-
2	EDO	A	305	-	-	1/1/1/1	-
2	EDO	A	308	-	-	0/1/1/1	-
2	EDO	B	306	-	-	1/1/1/1	-
2	EDO	B	307	-	-	1/1/1/1	-
2	EDO	B	303	-	-	0/1/1/1	-
2	EDO	A	310	-	-	0/1/1/1	-
2	EDO	B	309	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	311	GOL	C1-C2-C3-O3
3	A	311	GOL	O2-C2-C3-O3
3	A	311	GOL	O1-C1-C2-C3
3	A	311	GOL	O1-C1-C2-O2
2	A	307	EDO	O1-C1-C2-O2
2	B	304	EDO	O1-C1-C2-O2
2	A	304	EDO	O1-C1-C2-O2
2	A	313	EDO	O1-C1-C2-O2
2	B	308	EDO	O1-C1-C2-O2
2	A	301[B]	EDO	O1-C1-C2-O2
2	B	306	EDO	O1-C1-C2-O2
2	B	307	EDO	O1-C1-C2-O2
2	A	312	EDO	O1-C1-C2-O2
2	A	314	EDO	O1-C1-C2-O2
2	A	305	EDO	O1-C1-C2-O2
2	B	309	EDO	O1-C1-C2-O2

There are no ring outliers.

9 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	307	EDO	3	0
2	A	313	EDO	2	0
2	A	314	EDO	1	0
2	B	304	EDO	1	0
2	A	304	EDO	3	0
2	A	301[B]	EDO	4	0
2	B	310	EDO	1	0
2	A	308	EDO	2	0
2	B	303	EDO	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	178/196 (90%)	0.11	9 (5%) 28 22	21, 34, 77, 109	0
1	B	179/196 (91%)	0.02	11 (6%) 21 16	20, 34, 74, 101	0
All	All	357/392 (91%)	0.06	20 (5%) 24 19	20, 34, 75, 109	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	187	HIS	6.5
1	B	143	PHE	5.8
1	B	187	HIS	5.1
1	A	186	GLN	4.9
1	A	143	PHE	4.8
1	B	110	TYR	4.8
1	A	41	ASN	4.6
1	B	183	HIS	4.1
1	A	11	MET	3.5
1	A	110	TYR	3.4
1	B	188	ASN	3.3
1	B	10	ALA	3.2
1	A	188	ASN	3.0
1	B	186	GLN	3.0
1	B	141	PRO	2.9
1	A	183	HIS	2.8
1	B	97	ASN	2.8
1	A	141	PRO	2.3
1	B	142	SER	2.3
1	B	140	GLY	2.2

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	EDO	B	304	4/4	0.54	0.33	55,69,83,83	0
3	GOL	A	311	6/6	0.57	0.21	60,77,88,92	0
2	EDO	B	308	4/4	0.59	0.39	65,78,79,86	0
2	EDO	B	301[B]	4/4	0.60	0.23	44,56,69,70	10
2	EDO	A	306	4/4	0.66	0.34	63,76,82,82	0
2	EDO	A	301[B]	4/4	0.73	0.25	22,28,36,43	10
2	EDO	A	308	4/4	0.76	0.24	58,70,80,88	0
2	EDO	B	307	4/4	0.77	0.39	32,56,68,82	0
2	EDO	A	312	4/4	0.77	0.28	52,63,80,80	0
2	EDO	A	313	4/4	0.77	0.22	44,56,67,68	0
2	EDO	B	303	4/4	0.82	0.15	43,54,65,65	0
2	EDO	B	305	4/4	0.84	0.29	52,66,80,80	0
2	EDO	B	309	4/4	0.86	0.25	38,46,56,66	0
2	EDO	A	310	4/4	0.87	0.16	42,61,73,76	0
2	EDO	A	307	4/4	0.88	0.19	34,44,56,68	0
2	EDO	A	305	4/4	0.89	0.15	53,70,83,85	0
2	EDO	B	310	4/4	0.89	0.21	51,76,82,94	0
2	EDO	B	306	4/4	0.89	0.17	38,55,68,68	0
2	EDO	A	309	4/4	0.90	0.19	51,71,73,87	0
2	EDO	A	304	4/4	0.91	0.16	42,55,67,67	0
2	EDO	A	303	4/4	0.94	0.14	35,52,68,81	0
2	EDO	A	314	4/4	0.94	0.15	38,53,73,82	0
2	EDO	B	302[B]	4/4	0.96	0.18	29,43,52,52	10
2	EDO	A	302[B]	4/4	0.96	0.20	27,32,33,35	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.