



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 16, 2023 – 04:20 PM EDT

PDB ID : 4P07  
Title : Bacterial aryl sulfotransferase (ASST) soaked with human urine  
Authors : Malojcic, G.; Owen, R.L.; Glockshuber, R.  
Deposited on : 2014-02-20  
Resolution : 2.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	571	 92% 7% .
1	B	571	 92% 7% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	601	-	-	X	-
2	SO4	B	601	-	-	X	-

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9219 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Arylsulfate sulfotransferase AssT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	570	Total	C	N	O	S	6	5	0
			4500	2860	770	860	10			
1	B	570	Total	C	N	O	S	5	5	0
			4497	2859	768	860	10			

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	119	Total 119	O 119	0	0
3	B	93	Total 93	O 93	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Arylsulfate sulfotransferase AssT

Chain A:  92% 7%



- Molecule 1: Arylsulfate sulfotransferase AssT

Chain B:  92% 7%



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 1 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	182.29Å 182.29Å 101.59Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.40 – 2.59 48.35 – 2.59	Depositor EDS
% Data completeness (in resolution range)	98.2 (48.40-2.59) 98.2 (48.35-2.59)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	0.13	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.47 (at 2.58Å)	Xtrriage
Refinement program	REFMAC 5.7.0029, PHENIX	Depositor
R, $R_{free}$	0.180 , 0.224 0.184 , 0.226	Depositor DCC
$R_{free}$ test set	2945 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	47.3	Xtrriage
Anisotropy	0.739	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 17.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	0.105 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9219	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.76	2/4616 (0.0%)	0.90	5/6265 (0.1%)
1	B	0.74	1/4613 (0.0%)	0.85	5/6262 (0.1%)
All	All	0.75	3/9229 (0.0%)	0.87	10/12527 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	323	VAL	CB-CG2	-6.55	1.39	1.52
1	B	136	ASN	CG-ND2	-6.08	1.17	1.32
1	A	21	GLY	C-O	-5.20	1.15	1.23

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	323	VAL	CG1-CB-CG2	-22.54	74.84	110.90
1	B	52	GLU	OE1-CD-OE2	-7.17	114.70	123.30
1	A	283	ARG	NE-CZ-NH2	6.62	123.61	120.30
1	A	52	GLU	OE1-CD-OE2	-6.17	115.89	123.30
1	A	191	ASP	CB-CG-OD2	-6.14	112.77	118.30
1	A	172	ASP	CB-CG-OD1	5.64	123.38	118.30
1	B	213	ARG	NE-CZ-NH2	-5.39	117.61	120.30
1	B	509	ASP	CB-CG-OD1	5.39	123.15	118.30

*Continued on next page...*



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	283	ARG	NE-CZ-NH1	-5.25	117.67	120.30
1	B	172	ASP	CB-CG-OD1	5.11	122.90	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	469	ARG	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4500	0	4348	52	0
1	B	4497	0	4336	36	0
2	A	5	0	0	5	0
2	B	5	0	0	6	0
3	A	119	0	0	4	0
3	B	93	0	0	0	0
All	All	9219	0	8684	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (86) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:436:HIS:HE1	2:B:601:SO4:O2	1.28	1.17
1:B:326:ASP:HA	1:B:327:LEU:HB2	1.29	1.09
1:A:319:GLY:O	1:A:326:ASP:OD2	1.69	1.08
1:A:436:HIS:HE1	2:A:601:SO4:S	1.78	1.06
1:A:323:VAL:O	1:A:324:ASN:O	1.74	1.03
1:B:436:HIS:CE1	2:B:601:SO4:O2	2.12	1.02
1:A:436:HIS:HE1	2:A:601:SO4:O1	1.42	1.02
1:A:319:GLY:HA2	1:A:326:ASP:CG	1.82	0.99

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:116:LEU:HB3	3:A:805:HOH:O	1.61	0.98
1:A:319:GLY:CA	1:A:326:ASP:OD2	2.15	0.95
1:A:319:GLY:HA2	1:A:326:ASP:OD2	1.67	0.94
1:B:436:HIS:HE1	2:B:601:SO4:S	1.91	0.93
1:A:326:ASP:CG	1:A:327:LEU:HA	1.89	0.92
1:A:436:HIS:CE1	2:A:601:SO4:S	2.63	0.92
1:A:436:HIS:CE1	2:A:601:SO4:O1	2.25	0.90
1:A:319:GLY:C	1:A:326:ASP:OD2	2.10	0.88
1:A:326:ASP:CB	1:A:327:LEU:HA	2.08	0.83
1:A:186:TYR:OH	3:A:805:HOH:O	1.95	0.82
1:B:436:HIS:CE1	2:B:601:SO4:S	2.70	0.82
1:A:455:ARG:HG2	1:A:456:HIS:HD2	1.44	0.82
1:B:326:ASP:CA	1:B:327:LEU:HB2	2.10	0.81
1:B:323:VAL:O	1:B:325:VAL:HG23	1.81	0.79
1:A:436:HIS:CE1	2:A:601:SO4:O2	2.36	0.78
1:A:326:ASP:OD1	1:A:327:LEU:HA	1.83	0.78
1:A:323:VAL:HG23	1:A:324:ASN:H	1.50	0.76
1:A:455:ARG:HG2	1:A:456:HIS:CD2	2.21	0.75
1:B:455:ARG:HG2	1:B:456:HIS:HD2	1.52	0.73
1:A:323:VAL:CG2	1:A:324:ASN:H	2.01	0.73
1:A:323:VAL:HG23	1:A:324:ASN:N	2.04	0.73
1:B:455:ARG:HG2	1:B:456:HIS:CD2	2.23	0.73
1:B:323:VAL:HG22	1:B:324:ASN:H	1.61	0.66
1:B:326:ASP:HA	1:B:327:LEU:CB	2.11	0.65
1:A:323:VAL:C	1:A:324:ASN:O	2.35	0.63
1:A:294:ARG:NH1	1:B:60:GLU:OE1	2.32	0.61
1:A:323:VAL:CG2	1:A:324:ASN:N	2.60	0.59
1:B:324:ASN:O	1:B:325:VAL:HB	2.01	0.59
1:B:384:ASP:OD1	1:B:386[B]:GLN:HB2	2.02	0.59
1:A:327:LEU:H	1:A:327:LEU:HD23	1.68	0.58
1:A:319:GLY:CA	1:A:326:ASP:CG	2.62	0.58
1:A:323:VAL:HA	1:A:500:ILE:HD12	1.85	0.58
1:A:169:LEU:N	1:A:170:PRO:CD	2.69	0.55
1:A:60:GLU:OE1	1:B:294:ARG:NH1	2.38	0.55
1:B:323:VAL:HG22	1:B:324:ASN:N	2.22	0.55
1:A:326:ASP:HB2	1:A:327:LEU:HA	1.88	0.54
1:A:117:GLN:N	3:A:805:HOH:O	2.41	0.53
1:A:320:ALA:C	1:A:321:VAL:HG23	2.30	0.52
1:A:326:ASP:CB	1:A:327:LEU:CA	2.85	0.52
1:A:327:LEU:N	1:A:327:LEU:CD2	2.73	0.52
1:B:323:VAL:CG2	1:B:324:ASN:N	2.72	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:325:VAL:CG1	1:B:326:ASP:N	2.73	0.51
1:B:326:ASP:OD2	1:B:327:LEU:HB3	2.10	0.51
1:A:327:LEU:H	1:A:327:LEU:CD2	2.24	0.50
1:A:470:PHE:CD1	1:A:470:PHE:N	2.79	0.50
1:A:436:HIS:CE1	1:A:501:THR:HA	2.48	0.49
1:B:325:VAL:HG12	1:B:326:ASP:N	2.26	0.49
1:B:169:LEU:N	1:B:170:PRO:CD	2.75	0.49
1:A:56:PRO:HD3	1:A:567:GLN:OE1	2.14	0.48
1:B:423:LEU:HD13	1:B:456:HIS:CE1	2.50	0.47
1:A:324:ASN:O	1:A:325:VAL:HG23	2.15	0.47
1:A:423:LEU:HD13	1:A:456:HIS:CE1	2.49	0.47
1:B:320:ALA:C	1:B:321:VAL:HG23	2.35	0.47
1:A:265:ARG:NE	1:A:265:ARG:C	2.69	0.46
1:B:319:GLY:HA3	1:B:330:ALA:HA	1.98	0.46
1:B:326:ASP:CG	1:B:327:LEU:CB	2.85	0.45
1:B:436:HIS:CE1	2:B:601:SO4:O1	2.69	0.45
1:B:46:LYS:HG3	1:B:77:PHE:CZ	2.52	0.45
1:A:326:ASP:HB2	1:A:327:LEU:CA	2.47	0.44
1:B:323:VAL:HA	1:B:500:ILE:HD12	2.00	0.43
1:B:326:ASP:CG	1:B:327:LEU:HB3	2.38	0.43
1:A:223:VAL:HG13	1:A:254:SER:HB3	2.00	0.43
1:A:326:ASP:OD2	1:A:326:ASP:N	2.46	0.43
1:A:48:GLU:H	1:A:48:GLU:CD	2.21	0.43
1:B:48:GLU:H	1:B:48:GLU:CD	2.22	0.42
1:A:470:PHE:N	1:A:470:PHE:HD1	2.18	0.42
1:A:320:ALA:O	1:A:321:VAL:HG23	2.19	0.42
1:B:19:PRO:HG3	1:B:28:LEU:HB2	2.03	0.41
1:B:356:HIS:O	1:B:356:HIS:CD2	2.72	0.41
1:A:117:GLN:O	3:A:805:HOH:O	2.22	0.41
1:B:126:PRO:HA	1:B:129:GLU:OE2	2.20	0.41
1:A:46:LYS:HG3	1:A:77:PHE:CZ	2.56	0.41
1:B:325:VAL:O	1:B:326:ASP:HB3	2.21	0.41
1:B:374:ARG:HD2	2:B:601:SO4:O1	2.20	0.41
1:A:456:HIS:CD2	1:A:456:HIS:N	2.90	0.40
1:A:169:LEU:HD12	1:A:169:LEU:HA	1.95	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	568/571 (100%)	536 (94%)	29 (5%)	3 (0%)	29	52
1	B	568/571 (100%)	530 (93%)	33 (6%)	5 (1%)	17	35
All	All	1136/1142 (100%)	1066 (94%)	62 (6%)	8 (1%)	22	43

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	324	ASN
1	A	326	ASP
1	B	325	VAL
1	B	326	ASP
1	B	327	LEU
1	A	502	SER
1	B	324	ASN
1	B	502	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	478/483 (99%)	464 (97%)	14 (3%)	42	68
1	B	476/483 (99%)	464 (98%)	12 (2%)	47	73
All	All	954/966 (99%)	928 (97%)	26 (3%)	47	71

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38[A]	ASP
1	A	38[B]	ASP
1	A	52	GLU
1	A	265	ARG
1	A	323	VAL
1	A	326	ASP
1	A	327	LEU
1	A	403	SER
1	A	423	LEU
1	A	425	GLU
1	A	427	SER
1	A	503	ILE
1	A	524	ASP
1	A	534	ASN
1	B	38[A]	ASP
1	B	38[B]	ASP
1	B	52	GLU
1	B	265	ARG
1	B	323	VAL
1	B	403	SER
1	B	423	LEU
1	B	425	GLU
1	B	427	SER
1	B	445	THR
1	B	503	ILE
1	B	534	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	436	HIS
1	A	456	HIS
1	B	436	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	B	601	-	4,4,4	0.67	0	6,6,6	1.18	0
2	SO4	A	601	-	4,4,4	0.38	0	6,6,6	1.19	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	SO4	6	0
2	A	601	SO4	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	570/571 (99%)	-0.14	1 (0%) 95 95	27, 43, 73, 140	0
1	B	570/571 (99%)	-0.13	1 (0%) 95 95	28, 44, 78, 127	0
All	All	1140/1142 (99%)	-0.13	2 (0%) 95 95	27, 44, 76, 140	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	155	LYS	3.0
1	B	323	VAL	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	B	601	5/5	0.94	0.19	42,49,53,54	0
2	SO4	A	601	5/5	0.97	0.16	42,45,50,51	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.