

# Full wwPDB X-ray Structure Validation Report (i)

#### Aug 21, 2023 – 06:32 PM EDT

PDB ID : 2P1O

Title: Mechanism of Auxin Perception by the TIR1 ubiquitin ligase

Authors: Tan, X.; Calderon-Villalobos, L.I.A.; Sharon, M.; Robinson, C.V.; Estelle, M.;

Zheng, N.

Deposited on : 2007-03-06

Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35

### 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{ resolution range}( ext{Å}))$
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
	Δ.	4.00	34%				
1	A	160	59%	19%	• 21%		
	D	504	2%				
2	В	594	82%		11% • •		
			15%				
3	С	13	77%		15% 8%		



## 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 6308 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called SKP1-like protein 1A.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	А	197	Total	С	N	О	S	0	0	0
1	11	121	1018	641	166	206	5			


• Molecule 2 is a protein called TRANSPORT INHIBITOR RESPONSE 1 protein.

Mol	Chain	Residues		Atoms			ZeroOcc	AltConf	Trace	
2	B	571	Total	С	N	О	S	0	0	0
	ט	371	4495	2871	759	828	37		0	

• Molecule 3 is a protein called Auxin-responsive protein IAA7.

Mol	Chain	Residues	Aton	ns	ZeroOcc	AltConf	Trace
3	С	13	Total C 114 74	N O 23 17	0	0	0


• Molecule 4 is INOSITOL HEXAKISPHOSPHATE (three-letter code: IHP) (formula:  $C_6H_{18}O_{24}P_6$ ).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
1	R	1	Total	С	О	Р	0	0
4	Ъ	1	36	6	24	6	0	

 $\bullet$  Molecule 5 is NAPHTHALEN-1-YL-ACETIC ACID (three-letter code: NLA) (formula:  $C_{12}H_{10}O_2).$ 



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	В	1	Total 14	C 12	O 2	0	0

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	85	Total O 85 85	0	0
6	В	537	Total O 537 537	0	0
6	С	9	Total O 9 9	0	0




## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: SKP1-like protein 1A



• Molecule 2: TRANSPORT INHIBITOR RESPONSE 1 protein



• Molecule 3: Auxin-responsive protein IAA7







## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	101.97Å 82.46Å 125.32Å	Donositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 103.84° 90.00°	Depositor
Resolution (Å)	50.00 - 1.90	Depositor
Resolution (A)	49.51 - 1.90	EDS
% Data completeness	93.8 (50.00-1.90)	Depositor
(in resolution range)	93.8 (49.51-1.90)	EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$< I/\sigma(I) > 1$	3.15 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
D D.	0.184 , 0.231	Depositor
$R, R_{free}$	0.219 , $0.221$	DCC
$R_{free}$ test set	3757 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.2	Xtriage
Anisotropy	0.058	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.37, 62.6	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6308	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.63% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

### 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NLA, IHP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI		RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.60	0/1030	0.65	0/1388	
2	В	0.72	0/4593	0.83	$12/6226 \ (0.2\%)$	
3	С	0.69	0/118	0.75	0/159	
All	All	0.70	0/5741	0.80	12/7773~(0.2%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	С	0	1

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	В	399	MET	CG-SD-CE	-8.11	87.22	100.20
2	В	541	ARG	NE-CZ-NH2	-7.04	116.78	120.30
2	В	439	LEU	CA-CB-CG	6.98	131.35	115.30
2	В	321	VAL	CB-CA-C	-6.98	98.15	111.40
2	В	27	ARG	NE-CZ-NH2	-6.89	116.86	120.30
2	В	399	MET	CB-CG-SD	6.59	132.18	112.40
2	В	344	ARG	NE-CZ-NH2	-6.20	117.20	120.30
2	В	541	ARG	NE-CZ-NH1	5.95	123.28	120.30
2	В	484	ARG	NE-CZ-NH1	5.87	123.23	120.30
2	В	483	LEU	CB-CG-CD2	5.29	119.99	111.00
2	В	321	VAL	CG1-CB-CG2	5.19	119.21	110.90
2	В	483	LEU	CA-CB-CG	5.00	126.81	115.30



There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	С	12	ARG	Peptide

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1018	0	1010	24	0
2	В	4495	0	4529	72	0
3	С	114	0	118	2	0
4	В	36	0	6	1	0
5	В	14	0	9	0	0
6	A	85	0	0	6	0
6	В	537	0	0	20	0
6	С	9	0	0	1	0
All	All	6308	0	5672	93	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (93) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
2:B:458:MET:CE	2:B:461:LEU:HB2	1.33	1.57
2:B:269:CYS:HB3	6:B:1007:HOH:O	1.23	1.26
2:B:438:SER:HB3	6:B:1188:HOH:O	1.08	1.21
2:B:458:MET:CE	2:B:461:LEU:CB	2.26	1.13
2:B:458:MET:HE3	2:B:461:LEU:HB2	1.34	1.09
2:B:458:MET:HE1	2:B:461:LEU:HB2	1.22	1.08
2:B:551:CYS:HB2	6:B:1036:HOH:O	1.54	1.04
2:B:42:ARG:NH1	6:B:1039:HOH:O	1.97	0.97
2:B:369:MET:HG3	2:B:395:ASN:ND2	1.80	0.95
2:B:458:MET:HE2	2:B:461:LEU:HB2	1.53	0.90
2:B:458:MET:HE3	2:B:461:LEU:CB	1.96	0.86
2:B:332:VAL:O	2:B:336:THR:HG23	1.76	0.86



 $Continued\ from\ previous\ page...$ 

Continued from prev		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ ({\rm \AA})$	$\text{overlap } (\mathring{\mathbf{A}})$
1:A:149:GLU:HG2	6:A:203:HOH:O	1.75	0.86
2:B:290:ARG:HD2	6:B:1043:HOH:O	1.80	0.81
2:B:42:ARG:CZ	6:B:1039:HOH:O	2.28	0.80
2:B:300:THR:HG22	2:B:300:THR:O	1.83	0.78
2:B:174:HIS:HD2	6:B:1027:HOH:O	1.69	0.76
2:B:162:ASP:OD1	2:B:164:ARG:HG3	1.86	0.76
2:B:364:LEU:HD13	2:B:392:ILE:HD13	1.71	0.73
1:A:153:ARG:HH11	2:B:532:LYS:HD2	1.56	0.71
1:A:48:THR:HG22	1:A:51:ILE:H	1.57	0.70
1:A:98:LEU:O	1:A:102:ILE:HG12	1.91	0.70
1:A:58:TYR:OH	1:A:112:ASN:OD1	2.10	0.69
2:B:458:MET:HE1	2:B:461:LEU:CB	2.07	0.69
4:B:701:IHP:O43	6:B:1047:HOH:O	2.13	0.67
2:B:255:SER:HB2	6:B:1173:HOH:O	1.97	0.65
2:B:59:ALA:O	2:B:63:ARG:HG3	1.97	0.64
2:B:575:ASN:HD22	2:B:577:ASP:H	1.44	0.64
2:B:270:LEU:HD12	2:B:291:LEU:HD11	1.79	0.64
1:A:143:PHE:O	6:A:209:HOH:O	2.14	0.64
2:B:300:THR:O	2:B:300:THR:CG2	2.46	0.64
2:B:18:PHE:O	2:B:27:ARG:NH2	2.31	0.63
1:A:153:ARG:HH11	2:B:532:LYS:CD	2.12	0.62
1:A:153:ARG:NH1	2:B:532:LYS:HD2	2.15	0.61
2:B:566:ARG:HG2	2:B:568:ASP:OD1	2.00	0.61
1:A:102:ILE:HD13	1:A:117:THR:HB	1.82	0.60
1:A:5:LYS:HD3	6:A:219:HOH:O	2.01	0.59
2:B:496:LYS:CE	6:B:1013:HOH:O	2.51	0.58
2:B:255:SER:CB	6:B:1173:HOH:O	2.52	0.58
2:B:369:MET:HG3	2:B:395:ASN:HD21	1.64	0.58
1:A:48:THR:HG21	6:A:189:HOH:O	2.03	0.57
2:B:465:PHE:HZ	3:C:2:VAL:HG11	1.68	0.57
2:B:43:TRP:O	2:B:67:LYS:HD2	2.05	0.56
2:B:226:LYS:NZ	6:B:965:HOH:O	2.40	0.55
2:B:520:PHE:CZ	2:B:524:LYS:HD3	2.42	0.54
2:B:206:LEU:C	2:B:206:LEU:HD23	2.29	0.53
2:B:539:ASP:OD2	2:B:541:ARG:CD	2.57	0.53
2:B:47:LYS:HD2	2:B:70:SER:OG	2.09	0.53
2:B:458:MET:HE3	2:B:461:LEU:CD1	2.39	0.53
2:B:539:ASP:OD2	2:B:541:ARG:HD3	2.09	0.53
2:B:575:ASN:ND2	2:B:577:ASP:H	2.08	0.52
2:B:458:MET:HE3	2:B:461:LEU:HD13	1.93	0.51
1:A:148:GLU:HG3	2:B:32:LEU:HD22	1.94	0.50



Continued from previous page...

A + 1	A 4 0	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ ({\rm \AA})$	overlap (Å)
2:B:551:CYS:CB	6:B:1036:HOH:O	2.33	0.49
2:B:225:GLU:H	2:B:225:GLU:CD	2.15	0.49
2:B:328:ALA:HB3	6:B:1048:HOH:O	2.13	0.48
2:B:458:MET:HE3	2:B:461:LEU:CA	2.43	0.48
3:C:12:ARG:HD3	6:C:334:HOH:O	2.13	0.48
2:B:27:ARG:NH2	6:B:1045:HOH:O	2.44	0.47
1:A:48:THR:HG23	1:A:50:LYS:H	1.80	0.47
1:A:133:ILE:HG21	2:B:33:VAL:HG11	1.97	0.46
2:B:42:ARG:HB2	2:B:64:ARG:O	2.15	0.46
2:B:307:VAL:HG13	2:B:332:VAL:HG11	1.98	0.46
1:A:41:GLY:O	1:A:43:PRO:HD3	2.16	0.46
1:A:112:ASN:ND2	6:A:239:HOH:O	2.49	0.46
2:B:206:LEU:HD23	2:B:206:LEU:O	2.15	0.45
2:B:432:LYS:HD3	6:B:898:HOH:O	2.16	0.45
2:B:186:LEU:CD2	2:B:206:LEU:HD21	2.47	0.45
1:A:5:LYS:HB2	1:A:20:GLU:OE1	2.16	0.45
1:A:31:HIS:HB3	6:A:178:HOH:O	2.15	0.45
1:A:134:ARG:HB2	1:A:139:ILE:O	2.17	0.45
2:B:206:LEU:C	2:B:206:LEU:CD2	2.85	0.45
2:B:236:GLN:HG3	6:B:1272:HOH:O	2.15	0.45
1:A:102:ILE:HG12	1:A:102:ILE:H	1.56	0.44
1:A:27:GLN:O	1:A:31:HIS:HD2	2.00	0.44
2:B:214:LYS:NZ	6:B:1272:HOH:O	2.45	0.44
2:B:186:LEU:HD23	2:B:206:LEU:HD21	1.99	0.43
2:B:438:SER:CB	6:B:1188:HOH:O	1.95	0.43
2:B:570:PRO:HG2	2:B:573:VAL:HG23	2.01	0.43
2:B:130:LYS:N	2:B:130:LYS:HD2	2.33	0.43
1:A:131:GLU:HA	1:A:134:ARG:HG2	2.01	0.42
2:B:575:ASN:HD21	2:B:577:ASP:HB2	1.84	0.42
2:B:169:ASP:HB2	6:B:1232:HOH:O	2.19	0.42
2:B:424:PHE:CE2	2:B:439:LEU:HB3	2.55	0.42
2:B:520:PHE:CE2	2:B:524:LYS:HD3	2.54	0.42
1:A:5:LYS:HB2	1:A:20:GLU:CD	2.40	0.41
2:B:294:LEU:HD21	2:B:309:LEU:CD2	2.50	0.41
2:B:446:LYS:HE3	2:B:449:GLU:OE1	2.21	0.41
2:B:458:MET:CE	2:B:461:LEU:HD13	2.50	0.41
1:A:47:VAL:HG23	1:A:107:TYR:CD2	2.56	0.41
2:B:54:TYR:CZ	2:B:566:ARG:HD2	2.56	0.40
2:B:444:THR:HB	2:B:468:ASP:OD2	2.20	0.40
2:B:505:LEU:HB2	2:B:533:LEU:HD21	2.03	0.40

There are no symmetry-related clashes.



### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	121/160 (76%)	119 (98%)	1 (1%)	1 (1%)	19 9
2	В	569/594 (96%)	558 (98%)	11 (2%)	0	100 100
3	С	11/13 (85%)	11 (100%)	0	0	100 100
All	All	701/767 (91%)	688 (98%)	12 (2%)	1 (0%)	51 42

All (1) Ramachandran outliers are listed below:

$\mathbf{Mol}$	Chain	Res	Type
1	A	34	GLU

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percei	ntiles
1	A	115/137 (84%)	106 (92%)	9 (8%)	12	5
2	В	504/525 (96%)	481 (95%)	23 (5%)	27	17
3	С	12/12 (100%)	11 (92%)	1 (8%)	11	4
All	All	631/674 (94%)	598 (95%)	33 (5%)	23	14

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	ASN
1	A	48	THR



Continued from previous page...

Mol	Chain	Res	Type
1	A	52	LEU
1	A	92	LYS
1	A A A A	95	GLN
1	A	101	LEU
1	A	102	ILE
1	A	114	LEU
1	A A	136	THR
2	В	33	VAL
2	В	35	LYS
2	В	124	LEU
2	В	130	LYS
2 2	В	138	SER
	В	227	LEU
2	В	257	LEU
2 2	В	321	VAL
2	В	336	THR
2	В	359	LEU
2	В	364	LEU
2	В	373	LYS
2	В	374	LEU
2	В	399	MET
2 2	В	439	LEU
2	В	446	LYS
2	В	471	LEU
2	В	483	LEU
2	В	524	LYS
2	В	525	LEU
2	В	533	LEU
2	В	575	ASN
2	В	578	GLN
3	С	1	GLN

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such side chains are listed below:

Mol	Chain	Res	Type
1	A	31	HIS
1	A	95	GLN
2	В	317	GLN
2	В	383	GLN
2	В	395	ASN
2	В	501	ASN
2	В	575	ASN



Continued from previous page...

Mol	Chain	Res	Type
3	С	1	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Das	Res Link	Bond lengths			Bond angles		
MIOI			nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	IHP	В	701	-	36,36,36	1.31	5 (13%)	54,60,60	1.04	2 (3%)
5	NLA	В	801	-	15,15,15	1.21	1 (6%)	20,20,20	1.24	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

$\mathbf{Mol}$	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	IHP	В	701	-	-	5/30/54/54	0/1/1/1
5	NLA	В	801	-	-	0/4/4/4	0/2/2/2



All	(6)	) bond l	ength	outliers	are	listed	below:
-----	-----	----------	-------	----------	-----	--------	--------

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
4	В	701	IHP	P3-O43	3.22	1.67	1.54
4	В	701	IHP	P2-O12	3.14	1.65	1.59
4	В	701	IHP	P3-O33	2.91	1.66	1.54
4	В	701	IHP	P3-O13	2.55	1.64	1.59
4	В	701	IHP	P2-O32	2.32	1.63	1.54
5	В	801	NLA	C9-C10	2.24	1.41	1.36

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
4	В	701	IHP	O14-C4-C5	-2.29	103.29	108.69
5	В	801	NLA	C7-C6-C5	-2.23	117.98	123.19
5	В	801	NLA	C9-C10-C1	-2.19	117.86	120.89
5	В	801	NLA	O1-C12-C11	-2.07	117.12	123.04
4	В	701	IHP	O41-P1-O21	2.02	118.59	110.68

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	В	701	IHP	C1-O11-P1-O21
4	В	701	IHP	C2-O12-P2-O22
4	В	701	IHP	C4-O14-P4-O24
4	В	701	IHP	C3-O13-P3-O23
4	В	701	IHP	C4-O14-P4-O34

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	В	701	IHP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the



average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



### 5.7 Other polymers (i)

There are no such residues in this entry.



## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



### 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2	$OWAB(A^2)$	Q<0.9
1	A	127/160 (79%)	1.93	55 (43%) 0 0	12, 20, 26, 27	0
2	В	571/594 (96%)	0.17	13 (2%) 60 63	8, 16, 24, 32	0
3	С	13/13 (100%)	0.84	2 (15%) 2 2	14, 18, 24, 28	0
All	All	711/767 (92%)	0.50	70 (9%) 7 8	8, 17, 25, 32	0

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	137	PHE	7.1
1	A	39	ASP	6.9
2	В	8	SER	6.8
1	A	34	GLU	5.6
1	A	31	HIS	5.4
1	A	138	ASN	4.8
1	A	99	PHE	4.7
1	A	6	ILE	4.7
3	С	13	LYS	4.6
1	A	139	ILE	4.5
1	A	63	VAL	4.4
1	A	91	MET	4.4
1	A	40	ASN	4.3
1	A	61	ARG	4.2
1	A	35	ASP	4.2
2	В	578	GLN	4.2
1	A	21	ALA	4.1
1	A	124	MET	4.0
1	A	144	THR	3.8
1	A	140	LYS	3.8
1	A	62	HIS	3.8
1	A	92	LYS	3.7
1	A	33	VAL	3.7



Continued from previous page...

Mol	nued fron Chain	Res	Type	RSRZ
1	A	131	GLU	3.7
1	A	58	TYR	3.6
1	A	96	ALA	3.6
1	A	102	ILE	3.4
1	A	64	GLU	3.4
1	A	30	ALA	3.4
1	A	95	GLN	3.3
1	A	97	THR	3.2
2	В	9	PHE	3.2
2	В	12	GLU	3.1
1	A	135	THR	2.9
1	A	94	ASP	2.9
1	A	112	ASN	2.8
3	С	1	GLN	2.8
1	A	146	GLU	2.7
1	A	136	THR	2.7
1	A	125	ILE	2.7
1	A	19	GLU	2.7
1	A	24	LEU	2.7
2	В	106	TRP	2.7
1	A	27	GLN	2.7
1	A	107	TYR	2.6
1	A	41	GLY	2.6
1	A	127	GLY	2.6
1	A	100	GLU	2.5
1	A	142	ASP	2.5
1	A	134	ARG	2.4
1	A	111	LYS	2.4
2	В	320	TRP	2.4
1	A	20	GLU	2.4
1	A	59	CYS	2.4
1	A	42	VAL	2.4
1	A	103	LEU	2.3
1	A	143	PHE	2.3
1	A	130	PRO	2.2
1	A	121	VAL	2.2
1	A	147	GLU	2.2
2	В	174	HIS	2.2
2	В	345	VAL	2.2
2	В	319	LEU	2.1
2	В	33	VAL	2.0
2	В	346	PHE	2.0



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	В	290	ARG	2.0
1	A	98	LEU	2.0
2	В	378	LEU	2.0
1	A	22	VAL	2.0
1	A	32	MET	2.0

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
4	IHP	В	701	36/36	0.95	0.20	19,30,46,47	0
5	NLA	В	801	14/14	0.96	0.17	18,20,21,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





### 6.5 Other polymers (i)

There are no such residues in this entry.

