



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 16, 2023 – 04:23 PM EDT

PDB ID : 4P3W
Title : Crystal structure of the human filamin A Ig-like domains 20-21 in complex with migfilin peptide
Authors : Seppala, J.; Pentikainen, U.; Ylanne, J.
Deposited on : 2014-03-10
Resolution : 2.00 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	182	 4% 84% 10% • 5%
1	B	182	 4% 82% 12% • 6%
1	C	182	 7% 81% 14% 5%
1	D	182	 11% 85% • 11%
1	E	182	 2% 82% 13% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	182	<p>7% 79% 7% 14%</p>
2	G	24	<p>4% 58% 38%</p>
2	H	24	<p>4% 58% 38%</p>
2	I	24	<p>4% 50% 8% 42%</p>
2	J	24	<p>54% 46%</p>
2	K	24	<p>4% 54% 38%</p>
2	L	24	<p>4% 46% 50%</p>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8100 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Filamin-A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	172	1269	796	214	254	5	0	0	0
1	B	171	1264	794	215	250	5	0	0	0
1	E	172	1259	792	212	250	5	0	0	0
1	F	157	1120	703	192	220	5	0	0	0
1	D	162	1153	724	194	230	5	0	0	0
1	C	172	1262	794	213	250	5	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2148	GLY	-	expression tag	UNP P21333
A	2149	ALA	-	expression tag	UNP P21333
A	2150	MET	-	expression tag	UNP P21333
A	2151	GLY	-	expression tag	UNP P21333
B	2148	GLY	-	expression tag	UNP P21333
B	2149	ALA	-	expression tag	UNP P21333
B	2150	MET	-	expression tag	UNP P21333
B	2151	GLY	-	expression tag	UNP P21333
E	2148	GLY	-	expression tag	UNP P21333
E	2149	ALA	-	expression tag	UNP P21333
E	2150	MET	-	expression tag	UNP P21333
E	2151	GLY	-	expression tag	UNP P21333
F	2148	GLY	-	expression tag	UNP P21333
F	2149	ALA	-	expression tag	UNP P21333
F	2150	MET	-	expression tag	UNP P21333
F	2151	GLY	-	expression tag	UNP P21333
D	2148	GLY	-	expression tag	UNP P21333

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	2149	ALA	-	expression tag	UNP P21333
D	2150	MET	-	expression tag	UNP P21333
D	2151	GLY	-	expression tag	UNP P21333
C	2148	GLY	-	expression tag	UNP P21333
C	2149	ALA	-	expression tag	UNP P21333
C	2150	MET	-	expression tag	UNP P21333
C	2151	GLY	-	expression tag	UNP P21333

- Molecule 2 is a protein called Filamin-binding LIM protein 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	G	15	Total 113	C 74	N 19	O 20	0	0	0
2	H	15	Total 113	C 74	N 19	O 20	0	0	0
2	K	15	Total 113	C 74	N 19	O 20	0	0	0
2	L	12	Total 90	C 59	N 16	O 15	0	0	0
2	J	13	Total 97	C 64	N 17	O 16	0	0	0
2	I	14	Total 104	C 69	N 18	O 17	0	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	E	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0

- Molecule 4 is PRASEODYMIUM ION (three-letter code: PR) (formula: Pr).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Pr 1 1	0	0
4	B	1	Total Pr 1 1	0	0
4	E	1	Total Pr 1 1	0	0
4	D	1	Total Pr 1 1	0	0

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	E	1	Total Na 1 1	0	0
5	C	1	Total Na 1 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	31	Total O 31 31	0	0
6	B	32	Total O 32 32	0	0
6	H	4	Total O 4 4	0	0
6	E	20	Total O 20 20	0	0
6	F	5	Total O 5 5	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	D	7	Total O 7 7	0	0
6	C	17	Total O 17 17	0	0
6	I	1	Total O 1 1	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Filamin-A





- Molecule 1: Filamin-A




- Molecule 1: Filamin-A




- Molecule 1: Filamin-A





- Molecule 1: Filamin-A







• Molecule 1: Filamin-A




• Molecule 2: Filamin-binding LIM protein 1




• Molecule 2: Filamin-binding LIM protein 1



• Molecule 2: Filamin-binding LIM protein 1




• Molecule 2: Filamin-binding LIM protein 1




• Molecule 2: Filamin-binding LIM protein 1

Chain J:  54% 46%



● Molecule 2: Filamin-binding LIM protein 1

Chain I:  4% 50% 8% 42%



4 Data and refinement statistics i

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	88.15Å 88.15Å 394.71Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.86 – 2.00 43.86 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.6 (43.86-2.00) 100.0 (43.86-2.00)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.36 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.200 , 0.226 0.211 , 0.230	Depositor DCC
R_{free} test set	3874 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	40.6	Xtrriage
Anisotropy	0.360	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 45.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtrriage
Estimated twinning fraction	0.296 for -h-k,k,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8100	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, PR, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.09	1/1301 (0.1%)	1.03	3/1770 (0.2%)
1	B	1.11	1/1297 (0.1%)	1.05	6/1765 (0.3%)
1	C	0.92	2/1295 (0.2%)	0.95	2/1762 (0.1%)
1	D	0.85	0/1181	0.88	0/1609
1	E	1.10	4/1292 (0.3%)	0.99	1/1759 (0.1%)
1	F	1.00	1/1146 (0.1%)	0.99	2/1558 (0.1%)
2	G	1.03	0/115	1.05	0/156
2	H	1.13	0/115	1.18	0/156
2	I	0.91	0/106	1.02	1/144 (0.7%)
2	J	0.97	0/98	1.17	0/132
2	K	1.14	0/115	1.18	1/156 (0.6%)
2	L	0.88	0/90	1.14	0/120
All	All	1.02	9/8151 (0.1%)	0.99	16/11087 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	F	0	1
2	K	0	1
All	All	0	2

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	2189	GLU	CD-OE2	-7.92	1.17	1.25
1	E	2306	GLU	CD-OE1	7.77	1.34	1.25
1	E	2189	GLU	CD-OE2	-7.46	1.17	1.25
1	B	2189	GLU	CD-OE2	-6.49	1.18	1.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	2276	GLU	CD-OE1	-6.00	1.19	1.25

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	2288	ARG	NE-CZ-NH2	-7.98	116.31	120.30
1	B	2288	ARG	NE-CZ-NH1	7.28	123.94	120.30
1	A	2288	ARG	NE-CZ-NH2	-6.43	117.08	120.30
1	B	2250	ARG	NE-CZ-NH1	6.15	123.37	120.30
1	C	2288	ARG	NE-CZ-NH1	5.74	123.17	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	F	2327	SER	Peptide
2	K	6	GLU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1269	0	1205	10	0
1	B	1264	0	1201	8	0
1	C	1262	0	1198	11	0
1	D	1153	0	1064	4	0
1	E	1259	0	1189	10	0
1	F	1120	0	1035	6	0
2	G	113	0	121	0	0
2	H	113	0	121	0	0
2	I	104	0	115	1	0
2	J	97	0	108	0	0
2	K	113	0	121	1	0
2	L	90	0	101	1	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	C	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	5	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
5	C	1	0	0	1	0
5	E	1	0	0	0	0
6	A	31	0	0	0	0
6	B	32	0	0	0	0
6	C	17	0	0	1	0
6	D	7	0	0	0	0
6	E	20	0	0	0	0
6	F	5	0	0	0	0
6	H	4	0	0	0	0
6	I	1	0	0	0	0
All	All	8100	0	7579	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 45 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2235:GLU:HG2	1:A:2235:GLU:O	1.46	1.10
5:C:2402:NA:NA	6:C:2502:HOH:O	1.54	0.79
1:F:2178:VAL:HG22	1:F:2212:VAL:HG22	1.67	0.75
1:A:2235:GLU:O	1:A:2235:GLU:CG	2.30	0.73
1:C:2182:SER:OG	1:C:2184:LYS:HD3	1.95	0.66

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	170/182 (93%)	164 (96%)	4 (2%)	2 (1%)	13	7
1	B	169/182 (93%)	166 (98%)	2 (1%)	1 (1%)	25	19
1	C	170/182 (93%)	168 (99%)	2 (1%)	0	100	100
1	D	156/182 (86%)	151 (97%)	5 (3%)	0	100	100
1	E	170/182 (93%)	168 (99%)	2 (1%)	0	100	100
1	F	149/182 (82%)	143 (96%)	5 (3%)	1 (1%)	22	16
2	G	13/24 (54%)	13 (100%)	0	0	100	100
2	H	13/24 (54%)	13 (100%)	0	0	100	100
2	I	12/24 (50%)	12 (100%)	0	0	100	100
2	J	11/24 (46%)	10 (91%)	1 (9%)	0	100	100
2	K	13/24 (54%)	12 (92%)	0	1 (8%)	1	0
2	L	10/24 (42%)	10 (100%)	0	0	100	100
All	All	1056/1236 (85%)	1030 (98%)	21 (2%)	5 (0%)	29	23

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	2327	SER
2	K	7	LYS
1	A	2237	GLY
1	B	2195	ASN
1	A	2195	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	135/142 (95%)	131 (97%)	4 (3%)	41	41
1	B	134/142 (94%)	131 (98%)	3 (2%)	52	55
1	C	133/142 (94%)	129 (97%)	4 (3%)	41	41
1	D	117/142 (82%)	117 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	132/142 (93%)	130 (98%)	2 (2%)	65	69
1	F	112/142 (79%)	110 (98%)	2 (2%)	59	63
2	G	13/20 (65%)	12 (92%)	1 (8%)	13	8
2	H	13/20 (65%)	12 (92%)	1 (8%)	13	8
2	I	12/20 (60%)	12 (100%)	0	100	100
2	J	11/20 (55%)	11 (100%)	0	100	100
2	K	13/20 (65%)	12 (92%)	1 (8%)	13	8
2	L	10/20 (50%)	9 (90%)	1 (10%)	7	4
All	All	835/972 (86%)	816 (98%)	19 (2%)	50	53

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	L	8	ARG
1	C	2322	VAL
1	C	2327	SER
1	C	2219	GLN
2	H	7	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	2177	GLN
1	C	2239	HIS
1	E	2177	GLN
1	E	2220	HIS
1	E	2239	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 6 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	2401	-	4,4,4	0.52	0	6,6,6	0.72	0
3	SO4	C	2401	-	4,4,4	0.45	0	6,6,6	0.14	0
3	SO4	E	2401	-	4,4,4	0.56	0	6,6,6	0.24	0
3	SO4	B	2401	-	4,4,4	0.46	0	6,6,6	0.80	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	172/182 (94%)	0.22	7 (4%) 37 36	27, 38, 61, 87	0
1	B	171/182 (93%)	0.26	8 (4%) 31 30	26, 36, 61, 96	0
1	C	172/182 (94%)	0.44	13 (7%) 13 13	31, 47, 65, 80	0
1	D	162/182 (89%)	0.68	20 (12%) 4 3	35, 51, 77, 85	0
1	E	172/182 (94%)	0.23	4 (2%) 60 59	26, 40, 54, 65	0
1	F	157/182 (86%)	0.55	13 (8%) 11 10	28, 48, 76, 95	0
2	G	15/24 (62%)	0.40	1 (6%) 17 17	28, 34, 61, 68	0
2	H	15/24 (62%)	0.39	1 (6%) 17 17	28, 34, 54, 60	0
2	I	14/24 (58%)	0.58	1 (7%) 16 15	36, 42, 65, 65	0
2	J	13/24 (54%)	0.53	0 100 100	35, 41, 58, 67	0
2	K	15/24 (62%)	0.80	1 (6%) 17 17	33, 41, 67, 69	0
2	L	12/24 (50%)	0.88	1 (8%) 11 10	34, 36, 53, 60	0
All	All	1090/1236 (88%)	0.41	70 (6%) 19 18	26, 43, 71, 96	0

The worst 5 of 70 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	2167	PRO	5.5
1	D	2216	TYR	5.4
1	B	2196	HIS	5.1
1	D	2217	LYS	5.1
1	F	2216	TYR	5.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	NA	E	2402	1/1	0.68	0.21	50,50,50,50	1
3	SO4	C	2401	5/5	0.92	0.30	57,69,71,72	0
3	SO4	A	2401	5/5	0.95	0.17	43,50,53,56	0
3	SO4	B	2401	5/5	0.95	0.16	42,46,50,57	0
4	PR	B	2402	1/1	0.97	0.08	53,53,53,53	0
3	SO4	E	2401	5/5	0.97	0.25	47,56,60,60	0
4	PR	E	2403	1/1	0.98	0.08	45,45,45,45	0
4	PR	D	2401	1/1	0.98	0.04	57,57,57,57	0
4	PR	A	2402	1/1	0.98	0.08	53,53,53,53	0
5	NA	C	2402	1/1	0.99	0.35	66,66,66,66	1

6.5 Other polymers [i](#)

There are no such residues in this entry.