



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 24, 2023 – 10:30 PM EST

PDB ID : 3P6K  
Title : Crystal structure of a PLP-dependent aminotransferase (ZP\_03625122.1) from Streptococcus suis 89-1591 at 2.07 Å resolution  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2010-10-11  
Resolution : 2.07 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.31.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2684 (2.08-2.04)
Clashscore	141614	2801 (2.08-2.04)
Ramachandran outliers	138981	2768 (2.08-2.04)
Sidechain outliers	138945	2768 (2.08-2.04)
RSRZ outliers	127900	2646 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	375	 9% 90% 8% ..
1	B	375	 2% 91% 8% .

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6384 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aminotransferase class I and II.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S	Se			
1	A	369	2983	1916	484	573	1	4	5	0	6	0
1	B	374	3049	1956	495	587	1	4	6	0	9	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP B9WVA1
B	0	GLY	-	expression tag	UNP B9WVA1

- Molecule 2 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0

- Molecule 4 is UNKNOWN LIGAND (three-letter code: UNL) (formula: ).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	2	Total O 24 24	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	119	Total O 122 122	0	3
5	B	186	Total O 187 187	0	1


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aminotransferase class I and II



- Molecule 1: Aminotransferase class I and II



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	83.98Å 83.98Å 176.39Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.40 – 2.07 29.40 – 2.07	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.40-2.07) 99.8 (29.40-2.07)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.45 (at 2.06Å)	Xtrriage
Refinement program	REFMAC 5.5.0110	Depositor
R, $R_{free}$	0.178 , 0.208 0.184 , 0.213	Depositor DCC
$R_{free}$ test set	2249 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.0	Xtrriage
Anisotropy	0.244	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 49.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.033 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6384	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: LLP, UNL, EDO, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.78	0/3035	0.87	4/4121 (0.1%)
1	B	0.80	0/3110	0.89	6/4219 (0.1%)
All	All	0.79	0/6145	0.88	10/8340 (0.1%)

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	369	ARG	NE-CZ-NH1	6.85	123.73	120.30
1	B	275	ARG	NE-CZ-NH1	6.58	123.59	120.30
1	A	275	ARG	NE-CZ-NH1	5.99	123.30	120.30
1	A	229	ARG	NE-CZ-NH1	5.63	123.12	120.30
1	B	301	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	B	229	ARG	NE-CZ-NH2	-5.38	117.61	120.30
1	B	142	ASP	CB-CG-OD1	5.27	123.05	118.30
1	B	229	ARG	NE-CZ-NH1	5.23	122.92	120.30
1	A	369	ARG	NE-CZ-NH1	5.11	122.85	120.30
1	A	196	ARG	NE-CZ-NH2	-5.04	117.78	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2983	0	2940	24	0
1	B	3049	0	3017	20	0
2	A	5	0	0	0	0
2	B	10	0	0	0	0
3	A	4	0	6	0	0
4	B	24	0	0	0	0
5	A	122	0	0	0	0
5	B	187	0	0	0	0
All	All	6384	0	5963	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:147[B]:ARG:HH11	1:B:147[B]:ARG:HG2	1.36	0.88
1:B:147[B]:ARG:HG2	1:B:147[B]:ARG:NH1	1.91	0.81
1:A:98:TYR:CD1	1:B:3:LEU:HD22	2.30	0.67
1:B:306:SER:HB2	1:B:350:ALA:HB2	1.76	0.66
1:A:94:LEU:HD22	1:B:250:THR:HA	1.87	0.56
1:A:98:TYR:CE1	1:B:3:LEU:HD22	2.40	0.56
1:B:280:LEU:HD22	1:B:304:VAL:HG12	1.87	0.55
1:A:160:ASN:HD21	1:A:221:LLP:H2'2	1.72	0.55
1:A:3:LEU:HD23	1:B:98:TYR:HE2	1.73	0.53
1:A:20:ILE:HG23	1:A:21:TYR:CD2	2.44	0.52
1:A:39:LEU:C	1:A:39:LEU:HD12	2.30	0.51
1:A:98:TYR:OH	1:B:120:ILE:HG23	2.10	0.51
1:A:98:TYR:HD1	1:B:3:LEU:HD22	1.72	0.50
1:A:45[A]:GLU:HA	1:A:45[A]:GLU:OE1	2.11	0.49
1:B:160:ASN:HD21	1:B:221:LLP:H2'2	1.78	0.48
1:B:28:ILE:HD11	1:B:226:PRO:HG3	1.95	0.48
1:A:20:ILE:HG22	1:A:330:GLY:HA3	1.96	0.47
1:A:280:LEU:HD21	1:A:306:SER:HB3	1.96	0.47
1:A:297:VAL:HG12	1:A:312:ILE:HD13	1.97	0.47
1:A:169:VAL:HG22	1:A:303:ALA:CB	2.46	0.46
1:B:16[B]:GLU:OE2	1:B:26:VAL:HG21	2.16	0.45
1:B:306:SER:CB	1:B:350:ALA:HB2	2.46	0.44
1:B:147[B]:ARG:HH11	1:B:147[B]:ARG:CG	2.13	0.44
1:A:197:SER:HA	1:A:305:VAL:HG23	2.00	0.44
1:A:240:THR:HG22	1:A:244:ARG:NH1	2.33	0.43
1:A:133:ILE:HD12	1:A:166:THR:HG21	2.00	0.43

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3:LEU:HD12	1:A:4:PRO:HD3	2.01	0.43
1:A:103:PRO:HG2	1:B:1:MSE:HG3	2.00	0.42
1:A:3:LEU:HD11	1:A:119:ASP:HB3	2.02	0.42
1:A:162:ALA:HB2	1:A:169:VAL:HG13	2.01	0.42
1:A:3:LEU:HD11	1:A:119:ASP:O	2.19	0.42
1:B:294:GLU:CB	1:B:297:VAL:HG22	2.50	0.42
1:B:92:ALA:HB1	1:B:233:VAL:HG22	2.03	0.41
1:A:94:LEU:HD21	1:B:94:LEU:CD2	2.51	0.41
1:A:114:TYR:HB2	1:A:221:LLP:H2'3	2.03	0.41
1:B:86:THR:HA	1:B:251:MSE:SE	2.72	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	372/375 (99%)	361 (97%)	10 (3%)	1 (0%)	41	32
1	B	380/375 (101%)	373 (98%)	7 (2%)	0	100	100
All	All	752/750 (100%)	734 (98%)	17 (2%)	1 (0%)	51	45

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	29	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	325/322 (101%)	321 (99%)	4 (1%)	71	69
1	B	334/322 (104%)	330 (99%)	4 (1%)	71	69
All	All	659/644 (102%)	651 (99%)	8 (1%)	71	69

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	LEU
1	A	98	TYR
1	A	299	TYR
1	A	369	ARG
1	B	45	GLU
1	B	134	GLU
1	B	280	LEU
1	B	299	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	160	ASN
1	B	160	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	LLP	B	221	1	23,24,25	1.74	6 (26%)	25,32,34	1.49	6 (24%)
1	LLP	A	221	1	23,24,25	1.88	6 (26%)	25,32,34	1.30	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	B	221	1	-	7/16/17/19	0/1/1/1
1	LLP	A	221	1	-	9/16/17/19	0/1/1/1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	221	LLP	O3-C3	-5.18	1.25	1.37
1	A	221	LLP	O3-C3	-4.76	1.25	1.37
1	A	221	LLP	C2-N1	3.83	1.41	1.33
1	A	221	LLP	C4-C4'	3.46	1.53	1.46
1	B	221	LLP	C2-N1	3.39	1.40	1.33
1	A	221	LLP	C4'-NZ	2.90	1.37	1.27
1	B	221	LLP	C4-C4'	2.44	1.51	1.46
1	B	221	LLP	C4'-NZ	2.40	1.35	1.27
1	A	221	LLP	C6-N1	2.21	1.39	1.34
1	B	221	LLP	C6-N1	2.13	1.38	1.34
1	B	221	LLP	P-OP3	-2.11	1.46	1.54
1	A	221	LLP	CE-NZ	2.05	1.51	1.46

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	221	LLP	C4-C4'-NZ	-3.14	109.88	124.31
1	B	221	LLP	C3-C4-C5	2.89	120.48	118.26
1	A	221	LLP	C4-C4'-NZ	-2.88	111.08	124.31
1	A	221	LLP	CD-CE-NZ	2.65	117.43	110.93
1	A	221	LLP	C5-C6-N1	-2.40	119.82	123.82
1	B	221	LLP	O3-C3-C2	2.10	122.06	117.49
1	B	221	LLP	OP4-P-OP1	-2.08	100.65	106.47
1	B	221	LLP	C3-C4-C4'	-2.06	116.58	120.41
1	B	221	LLP	C5-C6-N1	-2.01	120.47	123.82

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	221	LLP	C4-C4'-NZ-CE
1	B	221	LLP	C4-C4'-NZ-CE
1	A	221	LLP	C3-C4-C4'-NZ
1	B	221	LLP	C3-C4-C4'-NZ
1	A	221	LLP	C5-C4-C4'-NZ
1	A	221	LLP	C6-C5-C5'-OP4
1	A	221	LLP	C5'-OP4-P-OP3
1	A	221	LLP	CD-CE-NZ-C4'
1	A	221	LLP	C4-C5-C5'-OP4
1	B	221	LLP	C4-C5-C5'-OP4
1	B	221	LLP	CD-CE-NZ-C4'
1	B	221	LLP	CE-CD-CG-CB
1	B	221	LLP	C6-C5-C5'-OP4
1	A	221	LLP	C5'-OP4-P-OP2
1	B	221	LLP	C5'-OP4-P-OP2
1	A	221	LLP	C5'-OP4-P-OP1

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	221	LLP	1	0
1	A	221	LLP	2	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are unknown - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	B	402	-	4,4,4	0.18	0	6,6,6	0.28	0
2	SO4	B	400	-	4,4,4	0.39	0	6,6,6	0.33	0
2	SO4	A	401	-	4,4,4	0.25	0	6,6,6	0.27	0
3	EDO	A	403	-	3,3,3	0.52	0	2,2,2	0.73	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	403	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	363/375 (96%)	0.39	34 (9%) <b>8</b> <b>9</b>	30, 53, 96, 132	0
1	B	367/375 (97%)	-0.00	7 (1%) 66 68	30, 44, 60, 93	0
All	All	730/750 (97%)	0.19	41 (5%) <b>24</b> <b>25</b>	30, 46, 84, 132	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	296	LEU	5.3
1	A	371	ASP	4.8
1	A	3	LEU	4.5
1	A	6	PHE	4.4
1	A	313	ALA	4.4
1	A	368	ARG	4.3
1	A	14	VAL	4.2
1	A	10	GLU	4.0
1	A	152	PRO	3.7
1	A	153	THR	3.3
1	A	17[A]	ASN	3.3
1	A	13	ASN	3.0
1	A	295	PRO	2.9
1	B	237	HIS	2.8
1	A	292	GLU	2.7
1	A	199	SER	2.6
1	A	20	ILE	2.6
1	A	370	PHE	2.6
1	A	4	PRO	2.6
1	A	98	TYR	2.6
1	A	315	ASP	2.4
1	B	374	ASN	2.4
1	A	111	TYR	2.4
1	B	249	TYR	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	19	ALA	2.4
1	A	270	GLN	2.3
1	A	289	GLN	2.3
1	B	43	ASN	2.3
1	B	373	GLU	2.3
1	A	293	GLU	2.2
1	B	45	GLU	2.2
1	A	5	ARG	2.2
1	A	358	LYS	2.2
1	A	7	GLY	2.2
1	A	328	GLU	2.1
1	B	46	ASP	2.1
1	A	291	ILE	2.1
1	A	21	TYR	2.1
1	A	117	LEU	2.0
1	A	148	GLN	2.0
1	A	136	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	LLP	A	221	24/25	0.97	0.14	33,46,54,55	0
1	LLP	B	221	24/25	0.98	0.15	31,40,45,48	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	UNL	B	390	12/-	0.68	0.30	45,48,49,49	12
4	UNL	B	375	12/-	0.92	0.11	32,40,42,43	12
3	EDO	A	403	4/4	0.93	0.10	54,56,56,59	0
2	SO4	B	402	5/5	0.98	0.10	65,66,67,67	0
2	SO4	A	401	5/5	0.98	0.08	63,63,65,66	0
2	SO4	B	400	5/5	0.99	0.08	47,49,52,53	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.