



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 7, 2023 – 12:37 PM JST

PDB ID : 4P76
Title : Cellular response to a crystal-forming protein
Authors : Tsutsui, H.; Jinno, Y.; Shoda, K.; Tomita, A.; Matsuda, M.; Yamashita, E.;
Katayama, H.; Nakagawa, A.; Miyawaki, A.
Deposited on : 2014-03-26
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	226	 81% 17% ..
1	B	226	 83% 11% . .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3609 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Photoconvertible fluorescent protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	224	1798	1147	305	335	11	0	0	0
1	B	220	1767	1127	298	331	11	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	cloning artifact	UNP Q53UG8
A	2	VAL	-	cloning artifact	UNP Q53UG8
A	65	CR8	HIS	chromophore	UNP Q53UG8
A	65	CR8	TYR	chromophore	UNP Q53UG8
A	65	CR8	GLY	chromophore	UNP Q53UG8
A	198	SER	GLN	engineered mutation	UNP Q53UG8
B	1	MET	-	cloning artifact	UNP Q53UG8
B	2	VAL	-	cloning artifact	UNP Q53UG8
B	65	CR8	HIS	chromophore	UNP Q53UG8
B	65	CR8	TYR	chromophore	UNP Q53UG8
B	65	CR8	GLY	chromophore	UNP Q53UG8
B	198	SER	GLN	engineered mutation	UNP Q53UG8

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Na	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	19	Total	O	0	0
			19	19		

Continued on next page...


Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	24	Total	O	0	0
			24	24		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Photoconvertible fluorescent protein

Chain A:  81% 17% ..



- Molecule 1: Photoconvertible fluorescent protein

Chain B:  83% 11% . .



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	49.86Å 84.43Å 118.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.35 – 2.90 40.35 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (40.35-2.90) 99.9 (40.35-2.90)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.45 (at 2.90Å)	Xtrriage
Refinement program	REFMAC 5.8.0069	Depositor
R, R_{free}	0.174 , 0.250 0.175 , 0.250	Depositor DCC
R_{free} test set	539 reflections (4.65%)	wwPDB-VP
Wilson B-factor (Å ²)	25.9	Xtrriage
Anisotropy	0.161	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 22.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3609	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 23.62 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.4943e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CR8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/1815	0.95	11/2448 (0.4%)
1	B	0.65	0/1784	0.94	12/2406 (0.5%)
All	All	0.65	0/3599	0.95	23/4854 (0.5%)

There are no bond length outliers.

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	153	ARG	NE-CZ-NH1	11.37	125.99	120.30
1	A	178	ARG	NE-CZ-NH1	11.28	125.94	120.30
1	A	153	ARG	NE-CZ-NH2	-11.04	114.78	120.30
1	B	138	ARG	NE-CZ-NH1	10.23	125.41	120.30
1	A	178	ARG	NE-CZ-NH2	-10.13	115.23	120.30
1	B	123	ARG	NE-CZ-NH2	9.88	125.24	120.30
1	B	92	ARG	NE-CZ-NH1	-9.18	115.71	120.30
1	B	138	ARG	NE-CZ-NH2	-8.57	116.02	120.30
1	B	123	ARG	NE-CZ-NH1	-8.32	116.14	120.30
1	B	178	ARG	NE-CZ-NH1	-8.11	116.24	120.30
1	B	153	ARG	NE-CZ-NH1	-7.47	116.56	120.30
1	A	138	ARG	NE-CZ-NH2	7.28	123.94	120.30
1	B	178	ARG	NE-CZ-NH2	7.23	123.92	120.30
1	A	138	ARG	NE-CZ-NH1	-7.07	116.76	120.30
1	B	153	ARG	NE-CZ-NH2	6.82	123.71	120.30
1	A	123	ARG	NE-CZ-NH2	-6.46	117.07	120.30
1	A	92	ARG	NE-CZ-NH1	6.34	123.47	120.30
1	B	92	ARG	NE-CZ-NH2	6.33	123.46	120.30
1	A	178	ARG	CD-NE-CZ	5.92	131.89	123.60
1	A	123	ARG	NE-CZ-NH1	5.91	123.25	120.30
1	A	92	ARG	NE-CZ-NH2	-5.35	117.63	120.30
1	B	153	ARG	CG-CD-NE	-5.32	100.62	111.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	138	ARG	CD-NE-CZ	5.10	130.73	123.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1798	0	1733	22	0
1	B	1767	0	1695	21	0
2	B	1	0	0	0	0
3	A	19	0	0	3	0
3	B	24	0	0	4	0
All	All	3609	0	3428	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:92:ARG:HD3	1:B:179:THR:OG1	1.73	0.88
1:B:138:ARG:HD3	3:B:420:HOH:O	1.78	0.83
1:A:2:VAL:N	3:A:301:HOH:O	2.12	0.83
1:A:106:ASN:HD21	1:A:120:ASN:HD22	1.31	0.77
1:B:106:ASN:HD21	1:B:120:ASN:HD22	1.34	0.73
1:B:132:ASN:ND2	3:B:401:HOH:O	2.21	0.72
1:B:138:ARG:CD	3:B:420:HOH:O	2.40	0.62
1:B:42:ASP:OD1	1:B:212:LYS:HG2	2.04	0.57
1:A:153:ARG:HG2	3:A:305:HOH:O	2.04	0.57
1:A:148:GLU:HA	1:A:161:VAL:HB	1.86	0.57
1:A:91:GLU:HB3	1:B:127:VAL:HB	1.88	0.55
1:A:101:ILE:HD11	1:B:101:ILE:CD1	2.38	0.54
1:A:203:SER:HB3	1:A:212:LYS:HD3	1.89	0.54
1:B:148:GLU:HA	1:B:161:VAL:HB	1.90	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:41:VAL:HG12	1:B:213:LEU:HB3	1.92	0.51
1:A:149:LYS:O	1:A:159:GLY:HA2	2.11	0.51
1:A:42:ASP:OD1	1:A:212:LYS:HG2	2.11	0.51
1:A:101:ILE:CD1	1:B:101:ILE:HD11	2.41	0.51
1:B:116:ASN:HD22	1:B:116:ASN:H	1.59	0.51
1:A:41:VAL:HG12	1:A:213:LEU:HB3	1.94	0.50
1:A:127:VAL:HB	1:B:91:GLU:HB3	1.93	0.50
1:B:151:TYR:HB3	1:B:192:TYR:CD1	2.49	0.48
1:A:101:ILE:HD11	1:B:101:ILE:HD11	1.96	0.47
1:A:175:CYS:HG	1:A:177:PHE:HE1	1.63	0.47
1:B:106:ASN:ND2	1:B:120:ASN:HD22	2.06	0.47
1:A:151:TYR:HB3	1:A:192:TYR:CD1	2.51	0.46
1:A:103:LEU:HG	1:B:103:LEU:HG	1.98	0.45
1:A:16:GLU:HG3	1:A:25:VAL:HG22	1.99	0.44
1:A:111:LYS:HB2	1:A:117:CYS:HB3	2.00	0.44
1:B:116:ASN:HD22	1:B:116:ASN:N	2.15	0.44
1:B:175:CYS:HG	1:B:177:PHE:HE1	1.64	0.44
1:A:157:LEU:HB3	1:A:181:TYR:HB2	2.03	0.41
1:B:67:ARG:NH1	3:B:411:HOH:O	2.53	0.40
1:B:140:THR:HG21	1:B:165:LEU:HD13	2.02	0.40
1:A:13:LEU:HB3	1:A:120:ASN:HB2	2.01	0.40
1:A:204:HIS:CE1	3:A:318:HOH:O	2.74	0.40
1:A:215:GLU:HG2	1:A:216:HIS:N	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	219/226 (97%)	214 (98%)	5 (2%)	0	100	100
1	B	215/226 (95%)	211 (98%)	4 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	434/452 (96%)	425 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/195 (99%)	188 (97%)	5 (3%)	46	77
1	B	190/195 (97%)	180 (95%)	10 (5%)	22	54
All	All	383/390 (98%)	368 (96%)	15 (4%)	32	66

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60	THR
1	A	75	GLU
1	A	132	ASN
1	A	178	ARG
1	A	225	ARG
1	B	29	LYS
1	B	41	VAL
1	B	60	THR
1	B	67	ARG
1	B	75	GLU
1	B	116	ASN
1	B	123	ARG
1	B	132	ASN
1	B	153	ARG
1	B	189	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	GLN
1	A	120	ASN
1	A	128	ASN
1	A	162	ASN
1	B	33	GLN
1	B	116	ASN
1	B	120	ASN
1	B	128	ASN
1	B	132	ASN
1	B	162	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CR8	B	65	1	20,27,28	1.67	4 (20%)	17,37,39	1.89	7 (41%)
1	CR8	A	65	1	20,27,28	1.65	5 (25%)	17,37,39	2.26	7 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CR8	B	65	1	-	2/8/25/26	0/3/3/3
1	CR8	A	65	1	-	3/8/25/26	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	65	CR8	C8-C7	4.72	1.47	1.36
1	A	65	CR8	C8-C7	4.15	1.46	1.36
1	B	65	CR8	C12-C11	-2.83	1.39	1.45
1	A	65	CR8	C12-C11	-2.75	1.39	1.45
1	A	65	CR8	CA3-N3	-2.45	1.44	1.49
1	B	65	CR8	C4-C11	-2.38	1.40	1.45
1	A	65	CR8	C1-N2	2.25	1.38	1.34
1	B	65	CR8	C1-N2	2.15	1.38	1.34
1	A	65	CR8	C4-C11	-2.12	1.41	1.45

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	65	CR8	C4-C5-C7	-5.12	117.60	121.95
1	B	65	CR8	C4-C5-C7	-3.55	118.93	121.95
1	A	65	CR8	C4-C11-C12	3.22	122.28	116.62
1	A	65	CR8	C6-C7-C5	3.15	122.81	116.20
1	A	65	CR8	C12-C6-C7	-2.78	119.58	121.95
1	B	65	CR8	O3-C3-CA3	-2.70	118.63	126.32
1	B	65	CR8	O13-C11-C12	-2.57	117.42	121.56
1	B	65	CR8	C4-C11-C12	2.56	121.14	116.62
1	B	65	CR8	C6-C7-C5	2.53	121.53	116.20
1	B	65	CR8	C12-C6-C7	-2.46	119.86	121.95
1	A	65	CR8	O3-C3-CA3	-2.43	119.42	126.32
1	A	65	CR8	C23-N11-C10	2.33	109.42	105.78
1	A	65	CR8	C6-C12-C11	-2.32	118.03	121.23
1	B	65	CR8	CA3-N3-C2	2.10	127.74	124.32

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	65	CR8	CA1-C20-C21-N22
1	A	65	CR8	CA1-C20-C21-C23
1	B	65	CR8	CA1-C20-C21-N22
1	B	65	CR8	CA1-C20-C21-C23
1	A	65	CR8	C21-C20-CA1-N1

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	223/226 (98%)	-0.53	0 100 100	10, 17, 35, 41	0
1	B	219/226 (96%)	-0.50	1 (0%) 91 91	9, 16, 32, 49	0
All	All	442/452 (97%)	-0.52	1 (0%) 95 95	9, 17, 34, 49	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	169	GLY	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CR8	B	65	25/26	0.97	0.14	16,16,17,18	0
1	CR8	A	65	25/26	0.98	0.12	15,15,18,20	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NA	B	301	1/1	0.94	0.17	19,19,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.