W O R L D W I D E
wwPDB
P R O T E I N   D A T A   B A N K

Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2022 – 01:17 pm BST

PDB ID	:	7P9I
Title	:	Structure of E.coli RlmJ in complex with an RNA conjugate (GAA-SAM)
Authors	:	Meynier, V.; Catala, M.; Oerum, S.; Barraud, P.; Tisne, C.
Deposited on	:	2021-07-27
Resolution	:	1.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see references ⓘ) were used in the production of this report:

|		
MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.28.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0267
CCP4	:	7.1.010 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.28.1

1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.

Metric	Percentile Ranks	Value
Rfree		0.213
Clashscore		4
Ramachandran outliers		0
Sidechain outliers		0.2%
RSRZ outliers		7.7%
RNA backbone		0.01

Worse   *Better*

■ Percentile relative to all X-ray structures

▯ Percentile relative to X-ray structures of similar resolution

Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R$_{free}$	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)
RNA backbone	3102	1015 (2.36-0.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	280	6% / 91% / 5% ·
1	B	280	8% / 89% / 8% ·
2	D	3	33% / 33% / 67%
2	E	3	33% / 33% / 33% / 33%

2 Entry composition ⓘ

There are 4 unique types of molecules in this entry. The entry contains 9610 atoms, of which 4488 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribosomal RNA large subunit methyltransferase J.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	271	Total	C	H	N	O	S	0	0	0
			4355	1394	2178	384	394	5			
1	B	273	Total	C	H	N	O	S	0	0	0
			4386	1402	2194	389	396	5			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	106	GLN	LEU	conflict	UNP P37634
A	191	THR	ILE	conflict	UNP P37634
B	106	GLN	LEU	conflict	UNP P37634
B	191	THR	ILE	conflict	UNP P37634

- Molecule 2 is a RNA chain called RNA conjugate (GAA-SAM).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	D	3	Total	C	H	N	O	P	0	0	0
			97	30	33	15	17	2			
2	E	3	Total	C	H	N	O	P	0	0	0
			97	30	33	15	17	2			

- Molecule 3 is 5'-{[(3S)-3-amino-3-carboxypropyl](3-aminopropyl)amino}-5'-deoxyadenosine (three-letter code: 6D6) (formula: $C_{17}H_{28}N_8O_5$) (labeled as "Ligand of Interest" by depositor).

6D6

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	D	1	Total	C	H	N	O	0	0
			54	17	25	7	5		
3	E	1	Total	C	H	N	O	0	0
			54	17	25	7	5		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	267	Total	O	0	0
			267	267		
4	B	274	Total	O	0	0
			274	274		
4	D	11	Total	O	0	0
			11	11		
4	E	15	Total	O	0	0
			15	15		

3 Residue-property plots ⓘ

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Ribosomal RNA large subunit methyltransferase J

• Molecule 1: Ribosomal RNA large subunit methyltransferase J

• Molecule 2: RNA conjugate (GAA-SAM)

• Molecule 2: RNA conjugate (GAA-SAM)

4 Data and refinement statistics (i)

| Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	62.48Å 57.59Å 79.44Å 90.00° 98.14° 90.00°	Depositor
Resolution (Å)	42.15 – 1.59 42.15 – 1.59	Depositor EDS
% Data completeness (in resolution range)	99.3 (42.15-1.59) 99.3 (42.15-1.59)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.36 (at 1.59Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.188 , 0.223 0.188 , 0.213	Depositor DCC
R_{free} test set	3705 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å^2)	19.7	Xtriage
Anisotropy	0.100	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	(Not available) , (Not available)	EDS
L-test for twinning[2]	$< |L| > = 0.50, < L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	9610	wwPDB-VP
Average B, all atoms (Å^2)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 74.29 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.5715e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

---

[1]Intensities estimated from amplitudes.
[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: 6D6, GMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#|Z| >5	RMSZ	#|Z| >5
1	A	0.37	0/2231	0.56	0/3028
1	B	0.40	0/2246	0.58	0/3047
2	D	7.72	19/49 (38.8%)	5.93	24/74 (32.4%)
2	E	7.70	18/49 (36.7%)	6.01	23/74 (31.1%)
All	All	1.19	37/4575 (0.8%)	1.08	47/6223 (0.8%)

All (37) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	4	A	C3'-C2'	-22.80	1.27	1.52
2	D	4	A	C3'-C2'	-22.57	1.27	1.52
2	E	3	A	C3'-C2'	-22.28	1.28	1.52
2	D	3	A	C3'-C2'	-21.89	1.28	1.52
2	D	3	A	C6-N6	21.61	1.51	1.33
2	E	3	A	C6-N6	21.32	1.51	1.33
2	D	4	A	C6-N6	16.54	1.47	1.33
2	E	4	A	C6-N6	16.43	1.47	1.33
2	D	4	A	C4'-O4'	-14.93	1.26	1.45
2	E	4	A	C4'-O4'	-14.84	1.26	1.45
2	E	3	A	O4'-C1'	-11.26	1.27	1.41
2	D	3	A	O4'-C1'	-11.06	1.27	1.41
2	E	3	A	N7-C5	-8.97	1.33	1.39
2	D	4	A	O4'-C1'	8.74	1.53	1.41
2	D	4	A	P-O5'	8.59	1.68	1.59
2	D	4	A	N7-C5	-8.42	1.34	1.39
2	D	3	A	N7-C5	-8.35	1.34	1.39
2	E	4	A	N7-C5	-8.19	1.34	1.39
2	E	4	A	P-O5'	8.10	1.67	1.59
2	E	4	A	O4'-C1'	7.86	1.51	1.41
2	E	3	A	N9-C8	-7.81	1.31	1.37
2	D	4	A	N9-C8	-7.80	1.31	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	3	A	C2'-C1'	7.78	1.61	1.53
2	E	3	A	C5'-C4'	-7.71	1.42	1.51
2	E	3	A	C2'-C1'	7.67	1.61	1.53
2	D	3	A	N9-C8	-7.51	1.31	1.37
2	E	4	A	N9-C8	-7.33	1.31	1.37
2	E	4	A	C4'-C3'	7.30	1.61	1.53
2	D	3	A	C5'-C4'	-7.26	1.42	1.51
2	D	3	A	P-O5'	7.18	1.67	1.59
2	D	4	A	C4'-C3'	6.97	1.60	1.53
2	D	4	A	C1'-N9	-6.88	1.37	1.46
2	E	4	A	C1'-N9	-6.70	1.37	1.46
2	E	3	A	P-O5'	6.44	1.66	1.59
2	D	3	A	O3'-P	5.92	1.68	1.61
2	D	4	A	C5'-C4'	5.65	1.58	1.51
2	E	3	A	O3'-P	5.15	1.67	1.61

All (47) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	3	A	C2-N3-C4	19.83	120.52	110.60
2	D	3	A	C2-N3-C4	19.44	120.32	110.60
2	E	4	A	C2-N3-C4	19.27	120.24	110.60
2	D	4	A	C2-N3-C4	19.12	120.16	110.60
2	E	3	A	N1-C2-N3	-14.26	122.17	129.30
2	D	3	A	N1-C2-N3	-13.80	122.40	129.30
2	E	4	A	N1-C2-N3	-13.46	122.57	129.30
2	D	4	A	N1-C2-N3	-12.20	123.20	129.30
2	D	4	A	N3-C4-C5	-11.61	118.67	126.80
2	E	3	A	N3-C4-C5	-11.01	119.09	126.80
2	E	4	A	N3-C4-C5	-10.73	119.29	126.80
2	D	3	A	N3-C4-C5	-10.62	119.37	126.80
2	E	3	A	N3-C4-N9	9.83	135.26	127.40
2	D	4	A	N3-C4-N9	9.39	134.91	127.40
2	D	3	A	N3-C4-N9	9.18	134.75	127.40
2	D	3	A	N7-C8-N9	-9.05	109.27	113.80
2	D	4	A	C1'-O4'-C4'	-9.04	102.67	109.90
2	E	3	A	N7-C8-N9	-8.74	109.43	113.80
2	D	3	A	C5-N7-C8	8.66	108.23	103.90
2	E	4	A	C5-N7-C8	8.64	108.22	103.90
2	E	3	A	C5-N7-C8	8.60	108.20	103.90
2	E	4	A	N3-C4-N9	8.54	134.23	127.40
2	D	4	A	C5-N7-C8	8.43	108.12	103.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	4	A	N7-C8-N9	-8.41	109.60	113.80
2	D	3	A	C8-N9-C4	8.22	109.09	105.80
2	E	3	A	C3'-C2'-C1'	8.16	108.03	101.50
2	E	3	A	C8-N9-C4	7.99	109.00	105.80
2	D	4	A	N7-C8-N9	-7.83	109.88	113.80
2	D	3	A	C3'-C2'-C1'	7.74	107.69	101.50
2	D	4	A	OP1-P-OP2	-7.19	108.82	119.60
2	E	4	A	C4-C5-N7	-7.09	107.16	110.70
2	D	4	A	C4-C5-N7	-6.99	107.20	110.70
2	E	4	A	C8-N9-C4	6.85	108.54	105.80
2	E	4	A	OP1-P-OP2	-6.72	109.52	119.60
2	E	3	A	C5'-C4'-C3'	-6.51	105.58	116.00
2	D	3	A	C5-C6-N6	-6.50	118.50	123.70
2	D	4	A	C8-N9-C4	6.44	108.38	105.80
2	D	3	A	C4-C5-N7	-6.38	107.51	110.70
2	E	3	A	C5-C6-N6	-6.32	118.64	123.70
2	E	4	A	C1'-O4'-C4'	-6.31	104.85	109.90
2	D	4	A	C5-C6-N6	-6.25	118.70	123.70
2	E	3	A	C4-C5-N7	-5.96	107.72	110.70
2	D	3	A	N1-C6-N6	5.41	121.85	118.60
2	D	3	A	OP1-P-OP2	-5.33	111.61	119.60
2	D	4	A	N1-C6-N6	5.25	121.75	118.60
2	E	3	A	OP1-P-OP2	-5.19	111.81	119.60
2	E	3	A	N1-C6-N6	5.09	121.65	118.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2177	2178	2176	12	0
1	B	2192	2194	2192	21	0
2	D	64	33	34	0	0
2	E	64	33	34	0	0
3	D	29	25	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	29	25	0	0	0
4	A	267	0	0	2	0
4	B	274	0	0	2	0
4	D	11	0	0	0	0
4	E	15	0	0	0	0
All	All	5122	4488	4436	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:77:GLU:HB2	1:B:256:VAL:HG11	1.59	0.84
1:B:232:ARG:HD2	1:B:234:GLY:H	1.47	0.79
1:A:77:GLU:HB2	1:A:256:VAL:HG11	1.72	0.71
1:B:77:GLU:HB2	1:B:256:VAL:CG1	2.22	0.68
1:A:77:GLU:HB2	1:A:256:VAL:CG1	2.24	0.67
1:A:87:HIS:HD2	4:A:433:HOH:O	1.82	0.61
1:B:232:ARG:CZ	1:B:234:GLY:HA3	2.37	0.55
1:B:18:LYS:HE3	1:B:19:HIS:NE2	2.24	0.53
1:B:232:ARG:HD2	1:B:234:GLY:N	2.21	0.53
1:B:257:LEU:HD12	1:B:274:VAL:HG23	1.91	0.53
1:B:277:ILE:HG22	1:B:278:VAL:HG23	1.92	0.52
1:A:42:HIS:CD2	1:A:165:PRO:HB3	2.46	0.51
1:A:176:VAL:HG11	1:A:194:LEU:HD11	1.91	0.50
1:A:176:VAL:HG11	1:A:194:LEU:CD1	2.41	0.50
1:B:199:VAL:HG12	1:B:200:LEU:HD22	1.94	0.50
1:A:170:LYS:H	1:A:170:LYS:HZ3	1.58	0.50
1:B:19:HIS:HB3	1:B:103:ILE:HD13	1.92	0.49
1:B:87:HIS:HD2	4:B:433:HOH:O	1.95	0.49
1:B:232:ARG:HD2	1:B:233:ARG:N	2.27	0.49
1:B:124:TYR:CZ	1:B:128:ARG:HD2	2.49	0.48
1:B:232:ARG:CD	1:B:234:GLY:H	2.21	0.48
1:B:225:ALA:HA	1:B:237:ALA:CB	2.45	0.47
1:A:157:ARG:HG2	4:B:550:HOH:O	2.15	0.46
1:B:257:LEU:CD1	1:B:274:VAL:HG23	2.46	0.45
1:B:75:PRO:HD2	1:B:78:LEU:HD12	1.99	0.44
1:A:274:VAL:HG13	4:A:349:HOH:O	2.18	0.43
1:B:77:GLU:CB	1:B:256:VAL:HG11	2.38	0.43
1:A:277:ILE:HG22	1:A:278:VAL:HG23	2.00	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:206:ARG:NH2	1:A:206:ARG:HB2	2.34	0.42
1:B:7:SER:HB3	1:B:10:ALA:HB2	2.02	0.42
1:B:263:LYS:HD3	1:B:263:LYS:HA	1.79	0.41
1:B:225:ALA:HA	1:B:237:ALA:HB2	2.00	0.41
1:A:38:TYR:CE2	1:A:40:ASP:HB2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	267/280 (95%)	261 (98%)	6 (2%)	0	100	100
1	B	269/280 (96%)	261 (97%)	8 (3%)	0	100	100
All	All	536/560 (96%)	522 (97%)	14 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/241 (96%)	231 (100%)	0	100	100
1	B	232/241 (96%)	231 (100%)	1 (0%)	91	84
All	All	463/482 (96%)	462 (100%)	1 (0%)	93	88

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	232	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	HIS
1	A	87	HIS
1	B	87	HIS

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	D	1/3 (33%)	0	0
2	E	1/3 (33%)	1 (100%)	0
All	All	2/6 (33%)	1 (50%)	0

All (1) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	E	4	A

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	#$|Z| > 2$	Counts	RMSZ	#$|Z| > 2$
3	6D6	E	101	2	24,31,32	3.48	8 (33%)	23,44,45	1.47	3 (13%)
3	6D6	D	101	2	24,31,32	3.56	8 (33%)	23,44,45	1.39	3 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	6D6	E	101	2	-	2/12/36/37	0/3/3/3
3	6D6	D	101	2	-	3/12/36/37	0/3/3/3

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	101	6D6	O3-C13	-9.26	1.28	1.41
3	E	101	6D6	O3-C13	-9.12	1.28	1.41
3	D	101	6D6	C14-C7	-7.90	1.31	1.53
3	D	101	6D6	C14-C13	7.77	1.65	1.53
3	E	101	6D6	C14-C7	-7.70	1.32	1.53
3	E	101	6D6	C14-C13	7.28	1.64	1.53
3	D	101	6D6	O3-C12	5.96	1.58	1.45
3	E	101	6D6	O3-C12	5.85	1.58	1.45
3	E	101	6D6	C6-C12	-3.76	1.40	1.51
3	D	101	6D6	C6-C12	-3.76	1.40	1.51
3	E	101	6D6	C15-N6	3.64	1.47	1.34
3	D	101	6D6	C15-N6	3.64	1.47	1.34
3	D	101	6D6	C10-C4	2.71	1.58	1.52
3	E	101	6D6	C10-C4	2.65	1.58	1.52
3	D	101	6D6	C4-N2	2.58	1.53	1.47
3	E	101	6D6	C4-N2	2.55	1.53	1.47

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	E	101	6D6	N8-C16-N7	-4.30	121.95	128.68
3	D	101	6D6	N8-C16-N7	-4.26	122.03	128.68
3	E	101	6D6	C7-C14-C13	3.30	105.95	100.98
3	D	101	6D6	C17-C9-N5	-3.09	106.18	109.40
3	D	101	6D6	C7-C14-C13	2.78	105.16	100.98
3	E	101	6D6	C17-C9-N5	-2.00	107.31	109.40

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	101	6D6	C7-C12-C6-N2
3	E	101	6D6	C7-C12-C6-N2
3	D	101	6D6	C1-C2-C3-N2
3	E	101	6D6	C1-C2-C3-N2
3	D	101	6D6	C12-C6-N2-C3

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 6D6 E 101

Ligand 6D6 D 101

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	A	271/280 (96%)	0.54	18 (6%)	18	17	12, 21, 34, 47	0
1	B	273/280 (97%)	0.60	22 (8%)	12	10	12, 19, 33, 46	0
2	D	2/3 (66%)	5.76	1 (50%)	0	0	20, 20, 20, 20	0
2	E	2/3 (66%)	2.79	1 (50%)	0	0	20, 20, 20, 20	0
All	All	548/566 (96%)	0.60	42 (7%)	13	11	12, 20, 34, 47	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	4	A	10.8
1	B	155	VAL	4.7
1	A	233	ARG	4.6
1	B	58	THR	4.5
1	B	200	LEU	4.5
2	E	4	A	4.0
1	A	200	LEU	4.0
1	B	232	ARG	3.9
1	A	170	LYS	3.9
1	B	57	ARG	3.7
1	B	233	ARG	3.7
1	B	132	GLN	3.4
1	A	155	VAL	3.3
1	B	231	ASP	3.2
1	A	58	THR	3.1
1	B	33	ASP	3.1
1	A	133	LYS	3.0
1	A	231	ASP	3.0
1	A	213	ALA	2.8
1	B	3	SER	2.8
1	B	74	LEU	2.7

Continued on next page...

WORLDWIDE
wwPDB
PROTEIN DATA BANK

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	8	PHE	2.7
1	B	72	ASP	2.6
1	B	171	THR	2.5
1	A	230	SER	2.5
1	A	8	PHE	2.5
1	B	170	LYS	2.4
1	A	138	ARG	2.4
1	B	267	ALA	2.4
1	A	202	GLN	2.4
1	B	206	ARG	2.3
1	A	247	TRP	2.3
1	A	171	THR	2.3
1	A	33	ASP	2.3
1	A	206	ARG	2.3
1	A	211	LEU	2.3
1	B	110	GLU	2.3
1	B	7	SER	2.3
1	B	230	SER	2.1
1	A	232	ARG	2.1
1	B	87	HIS	2.1
1	B	34	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å²)	Q<0.9
3	6D6	E	101	29/30	0.83	0.15	20,20,20,20	0
3	6D6	D	101	29/30	0.88	0.12	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 6D6 E 101:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative)
and green (positive)

Electron density around 6D6 D 101:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

6.5 Other polymers ⓘ

There are no such residues in this entry.