

Full wwPDB X-ray Structure Validation Report (i)

Dec 17, 2023 - 09:45 AM EST

PDB ID	:	6PAX
Title	:	CRYSTAL STRUCTURE OF THE HUMAN PAX-6 PAIRED DOMAIN-DNA
		COMPLEX REVEALS A GENERAL MODEL FOR PAX PROTEIN-DNA
		INTERACTIONS
Authors	:	Xu, H.E.; Rould, M.A.; Xu, W.; Epstein, J.A.; Maas, R.L.; Pabo, C.O.
Deposited on	:	1999-04-22
Resolution	:	2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motria	Whole archive	Similar resolution
	$(\# {\rm Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R _{free}	130704	$4661 \ (2.50-2.50)$
Clashscore	141614	$5346 \ (2.50-2.50)$
Ramachandran outliers	138981	$5231 \ (2.50-2.50)$
Sidechain outliers	138945	5233 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length		Quality of chain		
1	В	26	27%	65%		8%
2	С	26	15%	77%		8%
3	А	133		75%	21%	•



6PAX

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 2528 atoms, of which 367 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a DNA chain called 26 NUCLEOTIDE DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	В	26	Total 585	C 254	Н 54	N 100	O 152	Р 25	0	0	0

• Molecule 2 is a DNA chain called 26 NUCLEOTIDE DNA.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace		
2	С	26	Total 581	C 254	Н 52	N 94	O 156	Р 25	0	0	0

• Molecule 3 is a protein called HOMEOBOX PROTEIN PAX-6.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace		
3	А	133	Total 1278	C 627	Н 261	N 196	O 190	${S \atop 4}$	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	25	ARG	LYS	conflict	UNP P26367
А	59	ALA	GLU	conflict	UNP P26367
А	89	GLN	ARG	conflict	UNP P26367

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	В	15	Total O 15 15	0	0
4	С	29	Total O 29 29	0	0
4	А	40	Total O 40 40	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain B:
 27%
 65%
 8%

 Image: State of the state of th
- Molecule 1: 26 NUCLEOTIDE DNA



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	33.84Å 61.69Å 171.11Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Bosolution (Å)	20.00 - 2.50	Depositor
Resolution (A)	19.82 - 2.00	EDS
% Data completeness	96.3 (20.00-2.50)	Depositor
(in resolution range)	76.6(19.82-2.00)	EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$< I/\sigma(I) >$	-	Xtriage
Refinement program	X-PLOR 3.1	Depositor
B B.	0.233 , 0.256	Depositor
$\mathbf{n}, \mathbf{n}_{free}$	0.274 , 0.284	DCC
R_{free} test set	2020 reflections $(9.84%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	19.9	Xtriage
Anisotropy	0.035	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.34 , 78.7	EDS
L-test for twinning ¹	$ L > = 0.29, < L^2 > = 0.13$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	2528	wwPDB-VP
Average B, all atoms $(Å^2)$	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.29% of the height of the origin peak. No significant pseudotranslation is detected.

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
	Ullaili	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	В	0.84	0/596	1.13	1/918~(0.1%)	
2	С	0.80	0/592	0.96	0/912	
3	А	0.57	0/1033	0.66	0/1397	
All	All	0.71	0/2221	0.90	1/3227~(0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	1
2	С	0	3
All	All	0	4

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^{o})$	$Ideal(^{o})$
1	В	1002	DA	C5'-C4'-C3'	-8.91	98.06	114.10

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	В	1005	DA	Sidechain
2	С	2003	DC	Sidechain
2	С	2009	DA	Sidechain
2	С	2022	DA	Sidechain



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	В	531	54	294	40	1
2	С	529	52	296	54	1
3	А	1017	261	1040	30	0
4	А	40	0	0	2	0
4	В	15	0	0	3	0
4	С	29	0	0	5	0
All	All	2161	367	1630	111	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 31.

All (111) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
2:C:2002:DT:H2"	2:C:2003:DC:C5'	1.61	1.30
2:C:2001:DT:H2"	2:C:2002:DT:C5'	1.66	1.25
2:C:2023:DA:H2"	2:C:2024:DT:C5'	1.78	1.14
2:C:2002:DT:H2"	2:C:2003:DC:H5"	1.23	1.10
2:C:2023:DA:H2"	2:C:2024:DT:H5"	1.18	1.09
2:C:2001:DT:C2'	2:C:2002:DT:H5"	1.82	1.07
2:C:2002:DT:H2"	2:C:2003:DC:H5'	1.38	1.02
2:C:2023:DA:C2'	2:C:2024:DT:H5"	1.92	0.99
2:C:2002:DT:C2'	2:C:2003:DC:H5"	1.97	0.95
1:B:1002:DA:H2"	1:B:1003:DG:H5"	1.46	0.95
1:B:1011:DA:H2"	1:B:1012:DC:H5'	1.49	0.92
2:C:2010:DC:H2"	2:C:2011:DT:H5'	1.49	0.92
1:B:1011:DA:H2"	1:B:1012:DC:C5'	2.02	0.89
2:C:2012:DC:H2"	2:C:2013:DA:N7	1.87	0.89
2:C:2002:DT:H5'	2:C:2002:DT:H6	1.44	0.83
1:B:1002:DA:C2'	1:B:1003:DG:H5"	2.08	0.82
1:B:1008:DT:H2"	1:B:1009:DT:H5'	1.58	0.82
2:C:2001:DT:H5'	2:C:2001:DT:H6	1.43	0.80
2:C:2001:DT:H2"	2:C:2002:DT:H5"	0.86	0.78
2:C:2002:DT:C2'	2:C:2003:DC:C5'	2.52	0.77
2:C:2010:DC:H2"	2:C:2011:DT:C5'	2.16	0.76

Continued on next page...



	ti a a	Interatomic Clash		
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:B:1002:DA:H2"	1:B:1003:DG:C5'	2.15	0.75	
2:C:2012:DC:H2"	2:C:2013:DA:C8	2.22	0.74	
1:B:1024:DC:H2"	1:B:1025:DA:OP2	1.86	0.74	
2:C:2023:DA:H2"	2:C:2024:DT:H5'	1.69	0.70	
1:B:1001:DA:H2"	1:B:1002:DA:OP2	1.90	0.70	
3:A:87:TYR:HE1	3:A:103:LEU:HD22	1.57	0.69	
3:A:41:ARG:NH1	4:A:738:HOH:O	2.26	0.68	
1:B:1001:DA:H8	4:B:789:HOH:O	1.78	0.67	
2:C:2002:DT:H5'	2:C:2002:DT:C6	2.30	0.66	
3:A:95:PHE:HB2	3:A:98:GLU:HG3	1.78	0.66	
2:C:2008:DC:H5'	2:C:2008:DC:H6	1.61	0.65	
1:B:1005:DA:H2"	1:B:1006:DT:H5'	1.79	0.64	
2:C:2001:DT:C2'	2:C:2002:DT:C5'	2.59	0.62	
1:B:1011:DA:C2'	1:B:1012:DC:C5'	2.78	0.61	
1:B:1008:DT:H2"	1:B:1009:DT:C5'	2.30	0.60	
3:A:4:GLY:O	3:A:11:VAL:HA	2.02	0.60	
1:B:1002:DA:C3'	1:B:1003:DG:H5"	2.26	0.60	
2:C:2015:DG:H1'	2:C:2016:DC:H5'	1.82	0.60	
1:B:1001:DA:H2'	4:B:789:HOH:O	2.02	0.59	
1:B:1016:DT:O2	3:A:69:GLY:HA3	2.03	0.59	
3:A:86:GLN:O	3:A:90:GLU:HG2	2.03	0.59	
2:C:2010:DC:H1'	2:C:2011:DT:H5"	1.84	0.58	
1:B:1016:DT:O4	2:C:2013:DA:N1	2.35	0.58	
2:C:2010:DC:H2"	2:C:2011:DT:OP2	2.03	0.58	
2:C:2003:DC:H5'	4:C:772:HOH:O	2.06	0.55	
2:C:2005:DG:H2"	2:C:2006:DT:H5'	1.87	0.55	
4:C:814:HOH:O	3:A:72:LYS:HG3	2.06	0.55	
2:C:2003:DC:H5'	2:C:2003:DC:H6	1.73	0.54	
2:C:2008:DC:H5'	2:C:2008:DC:C6	2.43	0.54	
2:C:2010:DC:C2'	2:C:2011:DT:C5'	2.86	0.54	
1:B:1022:DC:H2'	1:B:1023:DA:C8	2.43	0.54	
1:B:1001:DA:C8	4:B:789:HOH:O	2.54	0.52	
1:B:1005:DA:H2"	1:B:1006:DT:C5'	2.39	0.52	
1:B:1003:DG:H2"	1:B:1004:DC:C6	2.45	0.52	
2:C:2014:DT:H2'	2:C:2015:DG:C8	2.45	0.51	
2:C:2017:DG:H5"	3:A:18:LEU:HD13	1.93	0.50	
1:B:1011:DA:C2'	1:B:1012:DC:H5"	2.42	0.50	
1:B:1011:DA:H1'	1:B:1012:DC:H5"	1.93	0.50	
1:B:1013:DG:H2"	1:B:1014:DC:H5'	1.94	0.50	
2:C:2007:DG:H2"	2:C:2008:DC:H5'	1.94	0.50	
2:C:2023:DA:C2'	2:C:2024:DT:C5'	2.65	0.50	

Continued from previous page...

Continued on next page...



		Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:B:1014:DC:H5'	1:B:1014:DC:H6	1.77	0.49	
2:C:2019:DG:H3'	3:A:7:GLN:NE2	2.27	0.49	
2:C:2001:DT:H5'	2:C:2001:DT:C6	2.35	0.48	
2:C:2019:DG:OP2	3:A:6:ASN:HB2	2.14	0.47	
3:A:88:LYS:HG3	3:A:124:LEU:CD2	2.45	0.47	
2:C:2015:DG:C4	2:C:2016:DC:C5	3.02	0.47	
1:B:1016:DT:O4	2:C:2013:DA:C6	2.68	0.47	
1:B:1005:DA:H1'	1:B:1006:DT:H5"	1.95	0.47	
3:A:1:SER:HA	4:A:702:HOH:O	2.14	0.47	
3:A:18:LEU:HD22	3:A:49:CYS:SG	2.55	0.46	
2:C:2003:DC:H2"	2:C:2004:DT:C6	2.51	0.46	
2:C:2009:DA:H2"	2:C:2010:DC:H5'	1.96	0.46	
2:C:2024:DT:H1'	2:C:2025:DG:H5'	1.97	0.46	
2:C:2010:DC:H1'	2:C:2011:DT:C5'	2.46	0.46	
3:A:18:LEU:HD12	3:A:18:LEU:HA	1.72	0.46	
1:B:1016:DT:H2'	1:B:1017:DG:C8	2.51	0.45	
1:B:1012:DC:H2"	1:B:1013:DG:C8	2.52	0.45	
1:B:1019:DG:C1'	3:A:74:ARG:NH2	2.80	0.45	
1:B:1025:DA:O5'	1:B:1025:DA:H2'	2.17	0.45	
1:B:1013:DG:H1'	1:B:1014:DC:H5"	1.99	0.45	
3:A:25:ARG:HD2	3:A:25:ARG:HA	1.84	0.45	
1:B:1019:DG:O4'	3:A:74:ARG:NH2	2.50	0.45	
1:B:1023:DA:H1'	1:B:1024:DC:H5'	1.99	0.44	
2:C:2003:DC:H2'	2:C:2004:DT:H71	1.99	0.44	
2:C:2010:DC:C2'	2:C:2011:DT:OP2	2.64	0.44	
1:B:1022:DC:C2'	1:B:1023:DA:C8	3.01	0.44	
2:C:2007:DG:H1'	2:C:2008:DC:H5"	2.00	0.44	
2:C:2008:DC:H2"	2:C:2009:DA:C8	2.53	0.44	
4:C:722:HOH:O	3:A:52:LYS:HE2	2.18	0.44	
2:C:2003:DC:H3'	4:C:770:HOH:O	2.16	0.43	
3:A:103:LEU:HD12	3:A:108:VAL:HB	2.01	0.43	
1:B:1003:DG:H2"	1:B:1004:DC:C5	2.54	0.43	
3:A:23:ARG:O	3:A:27:VAL:HG12	2.19	0.43	
2:C:2010:DC:C2'	2:C:2011:DT:H5'	2.33	0.43	
2:C:2013:DA:H2"	2:C:2014:DT:H5"	2.00	0.43	
4:C:770:HOH:O	3:A:97:TRP:HD1	2.02	0.43	
2:C:2012:DC:H2"	2:C:2013:DA:C5	2.53	0.43	
2:C:2017:DG:OP2	3:A:52:LYS:NZ	2.40	0.42	
3:A:54:LEU:HD23	3:A:54:LEU:HA	1.85	0.42	
3:A:83:LYS:HD3	3:A:83:LYS:HA	1.89	0.42	
1:B:1019:DG:C4'	3:A:74:ARG:HH21	2.33	0.42	

Continued from previous page...

Continued on next page...



6PAX

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:88:LYS:HG3	3:A:124:LEU:HD22	2.02	0.42
2:C:2010:DC:C1'	2:C:2011:DT:H5"	2.48	0.42
1:B:1012:DC:H5'	1:B:1012:DC:H6	1.85	0.41
1:B:1019:DG:H1'	3:A:74:ARG:NH2	2.35	0.41
2:C:2007:DG:H1'	2:C:2008:DC:C5'	2.51	0.41
1:B:1017:DG:H5'	3:A:68:ILE:O	2.20	0.40
1:B:1019:DG:H4'	3:A:74:ARG:HH21	1.86	0.40
3:A:104:LEU:HG	3:A:114:ILE:HD13	2.03	0.40

Continued from previous page...

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1002:DA:N1	2:C:2001:DT:H3[2_555]	1.49	0.11

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
3	А	131/133~(98%)	128 (98%)	3~(2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Perce	entiles
3	А	114/114~(100%)	106~(93%)	8 (7%)	15	29

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	А	18	LEU
3	А	27	VAL
3	А	52	LYS
3	А	86	GLN
3	А	89	GLN
3	А	90	GLU
3	А	102	ARG
3	А	133	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

