

# Full wwPDB X-ray Structure Validation Report (i)

Oct 3, 2023 – 08:53 AM EDT

PDB ID : 6PCA

Title : Crystal structure of beta-ketoadipyl-CoA thiolase

Authors: Sukritee, B.; Panjikar, S.

Deposited on : 2019-06-17

Resolution : 1.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : FAILED

Mogul : 1.8.5 (274361), CSD as 541be (2020)

Xtriage (Phenix) : 1.13

EDS : FAILED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.81 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



## 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 12855 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Beta-ketoadipyl-CoA thiolase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Λ	397	Total	С	N	О	S	3	3	0
1	A	391	2932	1830	527	556	19	3	3	
1	В	403	Total	С	N	О	S	4	0	0
1	Б	400	2956	1842	537	559	18	4	0	
1	С	402	Total	С	N	О	S	0	2	0
1		C 403	2973	1852	539	563	19	0	2	
1	D	206	Total	С	N	О	S	0	5	0
1		396	2934	1832	528	555	19	U	3	

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MET	-	initiating methionine	UNP Q88N39
A	-20	HIS	-	expression tag	UNP Q88N39
A	-19	HIS	-	expression tag	UNP Q88N39
A	-18	HIS	-	expression tag	UNP Q88N39
A	-17	HIS	-	expression tag	UNP Q88N39
A	-16	HIS	-	expression tag	UNP Q88N39
A	-15	HIS	-	expression tag	UNP Q88N39
A	-14	SER	-	expression tag	UNP Q88N39
A	-13	SER	-	expression tag	UNP Q88N39
A	-12	GLY	-	expression tag	UNP Q88N39
A	-11	VAL	-	expression tag	UNP Q88N39
A	-10	ASP	-	expression tag	UNP Q88N39
A	-9	LEU	-	expression tag	UNP Q88N39
A	-8	GLY	-	expression tag	UNP Q88N39
A	-7	THR	-	expression tag	UNP Q88N39
A	-6	GLU	-	expression tag	UNP Q88N39
A	-5	ASN	-	expression tag	UNP Q88N39
A	-4	LEU	-	expression tag	UNP Q88N39
A	-3	TYR	-	expression tag	UNP Q88N39
A	-2	HIS	-	expression tag	UNP Q88N39
A	-1	GLN	-	expression tag	UNP Q88N39

Continued on next page...



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q88N39
В	-21	MET	-	initiating methionine	UNP Q88N39
В	-20	HIS	-	expression tag	UNP Q88N39
В	-19	HIS	-	expression tag	UNP Q88N39
В	-18	HIS	-	expression tag	UNP Q88N39
В	-17	HIS	-	expression tag	UNP Q88N39
В	-16	HIS	-	expression tag	UNP Q88N39
В	-15	HIS	-	expression tag	UNP Q88N39
В	-14	SER	-	expression tag	UNP Q88N39
В	-13	SER	-	expression tag	UNP Q88N39
В	-12	GLY	-	expression tag	UNP Q88N39
В	-11	VAL	-	expression tag	UNP Q88N39
В	-10	ASP	-	expression tag	UNP Q88N39
В	-9	LEU	-	expression tag	UNP Q88N39
В	-8	GLY	-	expression tag	UNP Q88N39
В	-7	THR	-	expression tag	UNP Q88N39
В	-6	GLU	-	expression tag	UNP Q88N39
В	-5	ASN	-	expression tag	UNP Q88N39
В	-4	LEU	-	expression tag	UNP Q88N39
В	-3	TYR	-	expression tag	UNP Q88N39
В	-2	HIS	-	expression tag	UNP Q88N39
В	-1	GLN	-	expression tag	UNP Q88N39
В	0	SER	-	expression tag	UNP Q88N39
С	-21	MET	-	initiating methionine	UNP Q88N39
С	-20	HIS	-	expression tag	UNP Q88N39
С	-19	HIS	-	expression tag	UNP Q88N39
С	-18	HIS	-	expression tag	UNP Q88N39
С	-17	HIS	-	expression tag	UNP Q88N39
С	-16	HIS	-	expression tag	UNP Q88N39
С	-15	HIS	-	expression tag	UNP Q88N39
С	-14	SER	-	expression tag	UNP Q88N39
С	-13	SER	-	expression tag	UNP Q88N39
С	-12	GLY	-	expression tag	UNP Q88N39
С	-11	VAL	-	expression tag	UNP Q88N39
С	-10	ASP	-	expression tag	UNP Q88N39
С	-9	LEU	-	expression tag	UNP Q88N39
С	-8	GLY	-	expression tag	UNP Q88N39
С	-7	THR	-	expression tag	UNP Q88N39
С	-6	GLU	-	expression tag	UNP Q88N39
С	-5	ASN	-	expression tag	UNP Q88N39
С	-4	LEU	-	expression tag	UNP Q88N39
С	-3	TYR	-	expression tag	UNP Q88N39

Continued on next page...




 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
С	-2	HIS	-	expression tag	UNP Q88N39
С	-1	GLN	-	expression tag	UNP Q88N39
С	0	SER	-	expression tag	UNP Q88N39
D	-21	MET	-	initiating methionine	UNP Q88N39
D	-20	HIS	-	expression tag	UNP Q88N39
D	-19	HIS	-	expression tag	UNP Q88N39
D	-18	HIS	-	expression tag	UNP Q88N39
D	-17	HIS	-	expression tag	UNP Q88N39
D	-16	HIS	-	expression tag	UNP Q88N39
D	-15	HIS	-	expression tag	UNP Q88N39
D	-14	SER	-	expression tag	UNP Q88N39
D	-13	SER	-	expression tag	UNP Q88N39
D	-12	GLY	-	expression tag	UNP Q88N39
D	-11	VAL	_	expression tag	UNP Q88N39
D	-10	ASP	-	expression tag	UNP Q88N39
D	-9	LEU	-	expression tag	UNP Q88N39
D	-8	GLY	-	expression tag	UNP Q88N39
D	-7	THR	-	expression tag	UNP Q88N39
D	-6	GLU	-	expression tag	UNP Q88N39
D	-5	ASN	-	expression tag	UNP Q88N39
D	-4	LEU		expression tag	UNP Q88N39
D	-3	TYR	-	expression tag	UNP Q88N39
D	-2	HIS	-	expression tag	UNP Q88N39
D	-1	GLN	-	expression tag	UNP Q88N39
D	0	SER	-	expression tag	UNP Q88N39

• Molecule 2 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	В	1	Total C O 6 3 3	0	0
2	В	1	Total C O 6 3 3	0	0
2	С	1	Total C O 6 3 3	0	0
2	С	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0

• Molecule 3 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	С	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0

• Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total Cl 2 2	0	0
4	В	3	Total Cl 3 3	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	256	Total O 256 256	0	0
5	В	252	Total O 252 252	0	0
5	С	247	Total O 247 247	0	0
5	D	240	Total O 240 240	0	0



MolProbity and EDS failed to run properly - this section is therefore empty.



## 3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	110.08Å 113.84Å 127.42Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 - 1.81	Depositor
% Data completeness	99.4 (20.00-1.81)	Depositor
(in resolution range)	,	
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.08 (at 1.81Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
$R, R_{free}$	0.156 , $0.196$	Depositor
Wilson B-factor $(Å^2)$	19.2	Xtriage
Anisotropy	0.136	Xtriage
L-test for twinning <sup>2</sup>	$< L > = 0.50, < L^2> = 0.34$	Xtriage
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
Total number of atoms	12855	wwPDB-VP
Average B, all atoms $(\mathring{A}^2)$	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.71% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $< L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 4 Model quality (i)

### 4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3 Torsion angles (i)

#### 4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

### 4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 4.6 Ligand geometry (i)

Of 16 ligands modelled in this entry, 5 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	В	ond leng	$_{ m gths}$	В	ond ang	gles
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	ACT	D	503	-	3,3,3	1.33	0	3,3,3	0.57	0
2	GOL	A	501	-	5,5,5	0.14	0	5,5,5	0.29	0
3	ACT	A	503	-	3,3,3	1.09	0	3,3,3	1.00	0
2	GOL	D	502	-	5,5,5	0.31	0	5,5,5	0.64	0
2	GOL	D	501	-	5,5,5	0.09	0	5,5,5	0.44	0
2	GOL	A	502	-	5,5,5	0.30	0	5,5,5	0.63	0
3	ACT	С	503	-	3,3,3	0.78	0	3,3,3	1.08	0
2	GOL	С	502	-	5,5,5	0.22	0	5,5,5	0.59	0
2	GOL	В	502	-	5,5,5	0.18	0	5,5,5	0.72	0
2	GOL	С	501	-	5,5,5	0.18	0	5,5,5	0.32	0
2	GOL	В	501	-	5,5,5	0.09	0	5,5,5	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	501	-	-	0/4/4/4	-
2	GOL	D	502	-	-	2/4/4/4	-
2	GOL	D	501	-	-	0/4/4/4	-
2	GOL	A	502	-	-	2/4/4/4	_
2	GOL	С	502	-	-	2/4/4/4	-
2	GOL	В	502	-	-	2/4/4/4	-
2	GOL	С	501	-	-	0/4/4/4	-
2	GOL	В	501	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	502	GOL	O1-C1-C2-C3
2	В	502	GOL	C1-C2-C3-O3
2	С	502	GOL	C1-C2-C3-O3

Continued on next page...



Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	D	502	GOL	O1-C1-C2-O2
2	D	502	GOL	O1-C1-C2-C3
2	A	502	GOL	O1-C1-C2-O2
2	В	502	GOL	O2-C2-C3-O3
2	С	502	GOL	O2-C2-C3-O3

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


















































### 4.7 Other polymers (i)

There are no such residues in this entry.

### 4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



### 5 Fit of model and data (i)

#### 5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

#### 5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

#### 5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

#### 5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

### 5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

