

Full wwPDB X-ray Structure Validation Report (i)

Oct 3, 2023 – 03:33 PM EDT

PDB ID : 6PEY

Title: MTHFR with mutation Asp120Ala

Authors: Gallagher, E.L.; Gurney, L.A.; Frasco, F.G.; Trimmer, E.; Bolen, R.L.; Collins,

K.; Garland, E.; Halloran, J.; Handley-Pendelton, J.; Hernandez, V.; Leffler,

S.; Perez, A.; Soares, A.; Stojanoff, V.; Williams, D.

Deposited on : 2019-06-21

Resolution : 2.88 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# ext{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	2691 (2.90-2.86)
Clashscore	141614	2947 (2.90-2.86)
Ramachandran outliers	138981	2868 (2.90-2.86)
Sidechain outliers	138945	2871 (2.90-2.86)
RSRZ outliers	127900	2629 (2.90-2.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	304	79%	15%	• 6%
1	В	304	75%	19%	• 5%
1	С	304	71%	17% •	11%



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 6631 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Methylenetetrahydrofolate reductase.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	٨	287	Total	С	N	О	S	0	0	0
1	A	201	2153	1366	369	407	11	0	0	
1	D	290	Total	С	N	О	S	0	0	0
1	Б	290	2229	1413	387	418	11	0	0	0
1	С	271	Total	С	N	О	S	0	0	0
1		211	2087	1328	361	387	11	U	U	U

There are 27 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	120	ALA	ASP	engineered mutation	UNP C3SIT0
A	297	LEU	-	expression tag	UNP C3SIT0
A	298	GLU	-	expression tag	UNP C3SIT0
A	299	HIS	-	expression tag	UNP C3SIT0
A	300	HIS	-	expression tag	UNP C3SIT0
A	301	HIS	-	expression tag	UNP C3SIT0
A	302	HIS	-	expression tag	UNP C3SIT0
A	303	HIS	-	expression tag	UNP C3SIT0
A	304	HIS	-	expression tag	UNP C3SIT0
В	120	ALA	ASP	engineered mutation	UNP C3SIT0
В	297	LEU	-	expression tag	UNP C3SIT0
В	298	GLU	-	expression tag	UNP C3SIT0
В	299	HIS	-	expression tag	UNP C3SIT0
В	300	HIS	-	expression tag	UNP C3SIT0
В	301	HIS	-	expression tag	UNP C3SIT0
В	302	HIS	-	expression tag	UNP C3SIT0
В	303	HIS	-	expression tag	UNP C3SIT0
В	304	HIS	-	expression tag	UNP C3SIT0
С	120	ALA	ASP	engineered mutation	UNP C3SIT0
С	297	LEU	-	expression tag	UNP C3SIT0
С	298	GLU	-	expression tag	UNP C3SIT0
С	299	HIS	-	expression tag	UNP C3SIT0
С	300	HIS	_	expression tag	UNP C3SIT0



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
С	301	HIS	-	expression tag	UNP C3SIT0
С	302	HIS	-	expression tag	UNP C3SIT0
С	303	HIS	-	expression tag	UNP C3SIT0
С	304	HIS	-	expression tag	UNP C3SIT0

• Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	Λ	1	Total	С	N	О	Р	0	0
2	Λ	1	53	27	9	15	2	U	0
2	D	1	Total	С	N	О	Р	0	0
2	Б	1	53	27	9	15	2	U	0
2	C	1	Total	С	N	О	Р	0	0
		1	53	27	9	15	2	U	0

• Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O 1 1	0	0
3	В	1	Total O 1 1	0	0
3	С	1	Total O 1 1	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Methylenetetrahydrofolate reductase



• Molecule 1: Methylenetetrahydrofolate reductase



• Molecule 1: Methylenetetrahydrofolate reductase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	102.75Å 127.74Å 95.46Å	Domogiton
a, b, c, α , β , γ	90.00° 120.90° 90.00°	Depositor
Resolution (Å)	28.97 - 2.88	Depositor
Resolution (A)	28.95 - 2.88	EDS
% Data completeness	98.2 (28.97-2.88)	Depositor
(in resolution range)	98.4 (28.95-2.88)	EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.40 (at 2.90Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
D D	0.188 , 0.236	Depositor
R, R_{free}	0.191 , 0.237	DCC
R_{free} test set	1131 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	73.3	Xtriage
Anisotropy	0.040	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.32,45.4	EDS
L-test for twinning ²	$< L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	0.001 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6631	wwPDB-VP
Average B, all atoms (Å ²)	82.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.03% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.69	0/2197	0.85	0/2991	
1	В	0.69	0/2276	0.88	1/3092 (0.0%)	
1	С	0.69	0/2132	0.86	0/2894	
All	All	0.69	0/6605	0.86	1/8977 (0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^o)$
1	В	178	ASN	CB-CA-C	-5.99	98.42	110.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	В	182	THR	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2153	0	2039	25	0
1	В	2229	0	2151	32	0
1	С	2087	0	2037	31	0
2	A	53	0	31	1	0
2	В	53	0	31	0	0
2	С	53	0	31	2	0
3	A	1	0	0	0	0
3	В	1	0	0	0	0
3	C	1	0	0	0	0
All	All	6631	0	6320	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A 4 1	A 4 0	Interatomic	Clash	
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)	
1:A:122:PRO:O	1:A:124:GLY:N	2.20	0.75	
1:B:60:TYR:HE1	1:B:67:ARG:CB	2.01	0.73	
1:C:65:GLY:O	1:C:67:ARG:N	2.27	0.68	
1:C:122:PRO:O	1:C:124:GLY:N	2.32	0.62	
1:A:193:ARG:HH11	1:A:193:ARG:HG3	1.65	0.61	
1:A:118:ARG:NH1	1:A:119:GLY:O	2.32	0.61	
1:A:181:ILE:HG21	1:A:273:HIS:CD2	2.37	0.60	
1:A:193:ARG:HH11	1:A:193:ARG:CG	2.15	0.59	
1:B:132:ALA:O	1:B:136:VAL:HG23	2.04	0.58	
1:C:29:PHE:O	1:C:58:VAL:HA	2.04	0.57	
1:C:281:GLU:HA	1:C:281:GLU:OE2	2.04	0.57	
1:C:57:SER:HA	1:C:86:ALA:O	2.04	0.57	
1:B:93:ASP:N	1:B:93:ASP:OD1	2.36	0.56	
1:C:88:HIS:ND1	2:C:401:FAD:O4	2.36	0.55	
1:C:173:VAL:HG21	1:C:203:ILE:HG23	1.89	0.55	
1:B:285:ALA:O	1:B:288:HIS:HB3	2.05	0.54	
1:B:16:LEU:O	1:B:19:VAL:HG22	2.08	0.53	
1:C:121:LEU:HD12	1:C:122:PRO:HD2	1.90	0.53	
1:C:148:SER:OG	1:C:179:ARG:HD2	2.08	0.53	
1:A:188:VAL:CG2	1:A:260:MET:HG3	2.39	0.53	
1:B:57:SER:HB2	1:B:88:HIS:NE2	2.26	0.51	
1:C:148:SER:HA	1:C:179:ARG:O	2.11	0.51	



 $Continued\ from\ previous\ page...$

Continued from prev		Interatomic	Clash	
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)	
1:A:121:LEU:N	1:A:122:PRO:HD3	2.26	0.50	
1:B:29:PHE:O	1:B:58:VAL:HA	2.11	0.50	
1:C:187:ASP:O	1:C:190:SER:HB2	2.11	0.50	
1:A:281:GLU:HA	1:A:281:GLU:OE2	2.12	0.49	
1:B:32:PRO:HB3	1:B:37:MET:HG3	1.95	0.49	
1:B:151:ALA:O	1:B:182:THR:HA	2.13	0.49	
1:A:111:ILE:HD12	1:A:111:ILE:N	2.27	0.48	
1:C:195:ARG:NH2	1:C:267:GLU:O	2.45	0.48	
1:C:183:GLN:HG3	1:C:184:PHE:N	2.28	0.48	
1:A:163:GLN:HA	1:A:163:GLN:OE1	2.13	0.48	
1:A:107:TRP:CZ3	1:A:112:ARG:HG3	2.48	0.47	
1:B:149:VAL:HB	1:B:172:LYS:CE	2.44	0.47	
1:B:60:TYR:CE1	1:B:67:ARG:CB	2.90	0.47	
1:C:185:PHE:CZ	1:C:191:TYR:HB2	2.50	0.47	
1:A:26:SER:OG	1:A:273:HIS:ND1	2.34	0.46	
1:C:179:ARG:HG3	1:C:208:ILE:HD12	1.95	0.46	
1:C:285:ALA:O	1:C:288:HIS:HB3	2.15	0.46	
1:B:216:ASN:ND2	1:B:219:GLN:HB2	2.30	0.46	
1:C:261:VAL:O	1:C:265:SER:HB2	2.17	0.45	
1:B:121:LEU:HD21	1:B:128:PRO:HB3	1.99	0.45	
1:A:187:ASP:O	1:A:190:SER:HB2	2.17	0.45	
1:B:199:VAL:HG23	1:C:163:GLN:HE22	1.81	0.44	
1:C:218:LYS:HE2	1:C:243:ASP:OD1	2.17	0.44	
1:A:44:SER:HA	1:A:280:ALA:HB2	2.00	0.44	
1:B:202:GLY:HA2	1:C:167:LEU:HD21	1.99	0.44	
1:C:41:LEU:O	1:C:45:ILE:HG13	2.18	0.43	
1:A:60:TYR:HE1	1:A:67:ARG:CB	2.31	0.43	
1:A:251:LEU:HD23	1:C:251:LEU:HD23	1.99	0.43	
1:C:185:PHE:CE2	1:C:191:TYR:HB2	2.54	0.43	
1:B:266:ARG:NH2	1:C:187:ASP:OD2	2.52	0.43	
1:C:132:ALA:O	1:C:136:VAL:HG23	2.18	0.43	
1:C:238:MET:HG3	1:C:252:VAL:HG11	2.01	0.43	
1:A:182:THR:O	1:A:275:TYR:OH	2.34	0.42	
1:B:181:ILE:HG21	1:B:273:HIS:CD2	2.54	0.42	
1:C:222:LYS:O	1:C:226:MET:HG3	2.19	0.42	
1:A:196:ASP:O	1:A:199:VAL:HG12	2.19	0.42	
1:B:69:ARG:O	1:B:73:ILE:HG12	2.19	0.42	
1:B:122:PRO:O	1:B:124:GLY:N	2.53	0.42	
1:B:128:PRO:HG2	1:B:131:TYR:CE1	2.55	0.42	
1:C:44:SER:HA	1:C:280:ALA:HB2	2.01	0.42	
1:C:88:HIS:ND1	2:C:401:FAD:N5	2.67	0.42	



Continued from previous page...

A 4 1	A 4 0	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
1:A:62:ALA:HB1	1:A:92:ILE:HG21	2.02	0.42
1:B:111:ILE:N	1:B:111:ILE:HD13	2.34	0.42
1:B:69:ARG:N	1:B:69:ARG:HD2	2.35	0.42
1:B:266:ARG:HH11	1:B:266:ARG:HA	1.82	0.41
1:C:59:THR:HA	1:C:88:HIS:ND1	2.36	0.41
1:B:30:PHE:HA	1:B:59:THR:OG1	2.20	0.41
1:A:238:MET:HG2	1:A:256:ILE:HD11	2.02	0.41
1:B:31:PRO:HA	1:B:32:PRO:HD3	1.83	0.41
1:B:182:THR:HG23	1:B:208:ILE:O	2.21	0.41
1:B:235:MET:O	1:B:238:MET:HB2	2.20	0.41
1:A:156:HIS:CE1	1:A:157:PRO:HD2	2.55	0.41
1:A:96:PRO:O	1:A:100:ARG:NH1	2.53	0.41
1:B:60:TYR:HE2	1:B:90:THR:HG1	1.65	0.41
1:B:238:MET:HA	1:B:238:MET:HE3	2.03	0.41
1:A:113:HIS:HA	1:A:146:ASP:O	2.21	0.41
1:A:238:MET:HG3	1:A:252:VAL:CG1	2.51	0.40
1:B:214:VAL:O	1:B:253:GLY:HA3	2.21	0.40
1:C:31:PRO:HA	1:C:32:PRO:HD3	1.81	0.40
1:A:88:HIS:CE1	2:A:401:FAD:H6	2.57	0.40
1:B:44:SER:HA	1:B:280:ALA:HB2	2.04	0.40
1:B:185:PHE:CZ	1:B:191:TYR:HB2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	P	erce	entiles
1	A	283/304 (93%)	257 (91%)	23 (8%)	3 (1%)		14	40
1	В	286/304 (94%)	262 (92%)	19 (7%)	5 (2%)		9	28
1	С	267/304 (88%)	242 (91%)	21 (8%)	4 (2%)		10	32
All	All	836/912 (92%)	761 (91%)	63 (8%)	12 (1%)		11	34



All (12) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	123	PRO
1	В	62	ALA
1	С	66	GLU
1	A	19	VAL
1	В	155	VAL
1	С	123	PRO
1	A	155	VAL
1	В	61	GLY
1	С	155	VAL
1	В	63	ASN
1	В	123	PRO
1	С	111	ILE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	215/257~(84%)	207 (96%)	8 (4%)	34 66		
1	В	230/257 (90%)	220 (96%)	10 (4%)	29 60		
1	С	217/257 (84%)	210 (97%)	7 (3%)	39 71		
All	All	662/771 (86%)	637 (96%)	25 (4%)	33 65		

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	GLU
1	A	117	LEU
1	A	133	SER
1	A	146	ASP
1	A	193	ARG
1	A	197	ARG
1	A	247	GLU
1	A	279	ARG
1	В	7	SER



Continued from previous page...

Mol	Chain	Res	Type
1	В	69	ARG
1	В	75	LYS
1	В	101	THR
1	В	105	ASP
1	В	129	GLU
1	В	130	MET
1	В	189	GLU
1	В	238	MET
1	В	279	ARG
1	С	38	GLU
1	C C C	101	THR
1	С	112	ARG
1	С	179	ARG
1		190	SER
1	С	195	ARG
1	С	279	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	168	ASN
1	В	109	ASN
1	С	39	GLN
1	С	108	ASN
1	С	228	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type		Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Res	Link	Bond lengths			Bond angles		
Moi Type	Type	nes		LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2															
2	FAD	С	401	-	53,58,58	0.63	0	68,89,89	0.83	2 (2%)															
2	FAD	A	401	-	53,58,58	0.64	0	68,89,89	0.81	2 (2%)															
2	FAD	В	401	-	53,58,58	0.60	0	68,89,89	0.78	1 (1%)															

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	С	401	-	-	2/30/50/50	0/6/6/6
2	FAD	A	401	-	-	9/30/50/50	0/6/6/6
2	FAD	В	401	-	-	2/30/50/50	0/6/6/6

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\mathrm{Ideal}(^{o})$
2	A	401	FAD	C4-N3-C2	-2.17	121.63	125.64
2	A	401	FAD	C5A-C6A-N6A	2.13	123.59	120.35
2	В	401	FAD	C4-N3-C2	-2.09	121.78	125.64
2	С	401	FAD	C4-N3-C2	-2.05	121.85	125.64
2	С	401	FAD	O5'-C5'-C4'	2.03	114.77	109.36

There are no chirality outliers.

All (13) torsion outliers are listed below:

\mathbf{Mol}	Chain	Res	Type	${f Atoms}$
2	A	401	FAD	C5B-O5B-PA-O1A



Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	401	FAD	C3'-C4'-C5'-O5'
2	A	401	FAD	O4'-C4'-C5'-O5'
2	A	401	FAD	C5'-O5'-P-O2P
2	С	401	FAD	O4B-C4B-C5B-O5B
2	В	401	FAD	C3'-C4'-C5'-O5'
2	С	401	FAD	C3B-C4B-C5B-O5B
2	A	401	FAD	C5B-O5B-PA-O3P
2	A	401	FAD	C5'-O5'-P-O3P
2	A	401	FAD	C5'-O5'-P-O1P
2	A	401	FAD	N10-C1'-C2'-O2'
2	A	401	FAD	C2'-C3'-C4'-C5'
2	В	401	FAD	C5B-O5B-PA-O1A


There are no ring outliers.

2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	401	FAD	2	0
2	A	401	FAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q<0.9
1	A	287/304~(94%)	0.10	11 (3%) 40 36	52, 85, 131, 189	0
1	В	290/304 (95%)	-0.08	6 (2%) 63 62	41, 69, 129, 190	0
1	С	271/304 (89%)	0.08	14 (5%) 27 23	43, 77, 137, 194	0
All	All	848/912 (92%)	0.03	31 (3%) 41 37	41, 78, 136, 194	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	124	GLY	6.6
1	В	123	PRO	6.5
1	В	62	ALA	6.4
1	A	62	ALA	5.2
1	A	122	PRO	4.8
1	С	122	PRO	4.1
1	С	128	PRO	3.9
1	A	121	LEU	3.7
1	С	21	GLY	3.7
1	A	123	PRO	3.5
1	В	122	PRO	3.3
1	С	124	GLY	3.1
1	С	82	GLY	3.0
1	A	5	HIS	2.9
1	С	109	ASN	2.8
1	В	63	ASN	2.7
1	A	120	ALA	2.7
1	С	123	PRO	2.7
1	С	79	ASP	2.6
1	A	79	ASP	2.5
1	С	34	THR	2.5
1	A	124	GLY	2.5
1	A	63	ASN	2.4



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	С	33	ARG	2.4
1	С	144	ASP	2.4
1	A	108	ASN	2.3
1	С	68	ASP	2.2
1	С	26	SER	2.2
1	A	110	GLY	2.1
1	В	131	TYR	2.1
1	С	131	TYR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	FAD	A	401	53/53	0.77	0.32	87,136,163,173	0
2	FAD	В	401	53/53	0.81	0.26	79,110,126,135	0
2	FAD	С	401	53/53	0.83	0.26	81,106,125,129	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FAD A 401:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around FAD B 401:


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











6.5 Other polymers (i)

There are no such residues in this entry.

