

wwPDB EM Validation Summary Report (i)

Oct 11, 2023 - 11:41 pm BST

PDB ID	:	7PFO
EMDB ID	:	EMD-13375
Title	:	Core human replisome
Authors	:	Jones, M.J.; Yeeles, J.T.P.
Deposited on	:	2021-08-11
Resolution	:	3.20 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev50
Mogul	:	1.8.4, CSD as541be (2020)
MolProbity	:	4.02b-467
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.9
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motric	Whole archive	EM structures
IVIEUTIC	$(\# {\rm Entries})$	$(\# { m Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	2	904	• 76%	• 22%
2	3	808	75%	• 23%
3	4	863	70% •	28%
4	5	734	81%	• 18%
5	6	821	76%	• 23%
6	7	719	82%	• 17%
7	А	527	97%	•
8	В	2286	7% 34% • 65%	



Mol	Chain	Length	Quality of chain	
9	С	569	93%	• 6%
10	D	196	99%	••
11	Е	185	95%	• 5%
12	F	216	87%	• 10%
13	G	262	75% •	23%
14	Н	1161	• 34% 65%	
14	Ι	1161	• 34% • 65%	
14	J	1161	7% 34% • 65%	
15	K	1209	• 47%	
16	L	301	28% • 71%	
17	М	85	• 35% • 62%	
18	Ν	54	· 59%	
19	Q	1371	6% 94%	



2 Entry composition (i)

There are 23 unique types of molecules in this entry. The entry contains 136660 atoms, of which 68143 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called DNA replication licensing factor MCM2.

Mol	Chain	Residues			AltConf	Trace				
1	2	703	Total 11177	C 3511	Н 5597	N 994	0 1043	S 32	0	0

• Molecule 2 is a protein called DNA replication licensing factor MCM3.

Mol	Chain	Residues			AltConf	Trace				
2	3	620	Total 9770	C 3037	Н 4912	N 859	O 936	S 26	0	0

• Molecule 3 is a protein called DNA replication licensing factor MCM4.

Mol	Chain	Residues			Atom	s			AltConf	Trace
3	4	619	Total 9922	C 3103	Н 4993	N 873	O 926	S 27	0	0

• Molecule 4 is a protein called DNA replication licensing factor MCM5.

Mol	Chain	Residues			AltConf	Trace				
4	5	605	Total 9544	C 2968	H 4809	N 841	0 891	S 35	0	0

• Molecule 5 is a protein called DNA replication licensing factor MCM6.

Mol	Chain	Residues			Atom	S			AltConf	Trace
5	6	632	Total 10065	C 3150	Н 5051	N 890	0 948	S 26	0	0

• Molecule 6 is a protein called DNA replication licensing factor MCM7.

Mol	Chain	Residues			Atom	s			AltConf	Trace
6	7	598	Total 9552	C 2983	H 4802	N 840	O 897	S 30	0	0



• Molecule 7 is a protein called DNA polymerase epsilon subunit 2.

Mol	Chain	Residues	Atoms						AltConf	Trace
7	А	527	Total 8359	C 2702	Н 4160	N 697	O 780	S 20	0	0

• Molecule 8 is a protein called DNA polymerase epsilon catalytic subunit A.

Mol	Chain	Residues			Aton	ns			AltConf	Trace
8	В	804	Total	C 4162	H 6478	N 1116	0 1176	S 42	0	0
			12970	4102	0478	1110	1170	43		

• Molecule 9 is a protein called Cell division control protein 45 homolog.

Mol	Chain	Residues			Atom	.S			AltConf	Trace
9	С	534	Total 8647	C 2764	Н 4300	N 745	O 807	S 31	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	135Z	ASP	-	linker	UNP 075419
С	136A	TYR	-	linker	UNP 075419
С	136B	LYS	-	linker	UNP 075419
С	136C	ASP	-	linker	UNP 075419
С	136D	ASP	-	linker	UNP 075419
С	136E	ASP	-	linker	UNP 075419

• Molecule 10 is a protein called DNA replication complex GINS protein PSF1.

Mol	Chain	Residues			Atom	s			AltConf	Trace
10	D	195	Total 3207	C 1013	Н 1601	N 289	O 292	S 12	0	0

• Molecule 11 is a protein called DNA replication complex GINS protein PSF2.

Mol	Chain	Residues			Atom	S			AltConf	Trace
11	Е	176	Total 2887	C 916	H 1456	N 242	0 264	S 9	0	0

• Molecule 12 is a protein called DNA replication complex GINS protein PSF3.



Mol	Chain	Residues			Atom	IS			AltConf	Trace
12	F	194	Total 3048	C 979	Н 1502	N 268	O 293	S 6	0	0

• Molecule 13 is a protein called DNA replication complex GINS protein SLD5.

Mol	Chain	Residues			Atom	.s			AltConf	Trace
13	G	203	Total 3381	C 1065	Н 1702	N 290	O 314	S 10	0	0

There are 39 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	-38	MET	-	initiating methionine	UNP Q9BRT9
G	-37	TRP	-	expression tag	UNP Q9BRT9
G	-36	SER	-	expression tag	UNP Q9BRT9
G	-35	HIS	-	expression tag	UNP Q9BRT9
G	-34	PRO	-	expression tag	UNP Q9BRT9
G	-33	GLN	-	expression tag	UNP Q9BRT9
G	-32	PHE	-	expression tag	UNP Q9BRT9
G	-31	GLU	-	expression tag	UNP Q9BRT9
G	-30	LYS	-	expression tag	UNP Q9BRT9
G	-29	GLY	-	expression tag	UNP Q9BRT9
G	-28	GLY	-	expression tag	UNP Q9BRT9
G	-27	GLY	-	expression tag	UNP Q9BRT9
G	-26	SER	-	expression tag	UNP Q9BRT9
G	-25	GLY	-	expression tag	UNP Q9BRT9
G	-24	GLY	-	expression tag	UNP Q9BRT9
G	-23	GLY	-	expression tag	UNP Q9BRT9
G	-22	SER	-	expression tag	UNP Q9BRT9
G	-21	GLY	-	expression tag	UNP Q9BRT9
G	-20	GLY	-	expression tag	UNP Q9BRT9
G	-19	SER	-	expression tag	UNP Q9BRT9
G	-18	ALA	-	expression tag	UNP Q9BRT9
G	-17	TRP	-	expression tag	UNP Q9BRT9
G	-16	SER	-	expression tag	UNP Q9BRT9
G	-15	HIS	-	expression tag	UNP Q9BRT9
G	-14	PRO	-	expression tag	UNP Q9BRT9
G	-13	GLN	-	expression tag	UNP Q9BRT9
G	-12	PHE	-	expression tag	UNP Q9BRT9
G	-11	GLU	-	expression tag	UNP Q9BRT9
G	-10	LYS	-	expression tag	UNP Q9BRT9
G	-9	SER	-	expression tag	UNP Q9BRT9
G	-8	GLY	-	expression tag	UNP Q9BRT9



Chain	Residue	Modelled	Actual	Comment	Reference
G	-7	LEU	-	expression tag	UNP Q9BRT9
G	-6	GLU	-	expression tag	UNP Q9BRT9
G	-5	VAL	-	expression tag	UNP Q9BRT9
G	-4	LEU	-	expression tag	UNP Q9BRT9
G	-3	PHE	-	expression tag	UNP Q9BRT9
G	-2	GLN	-	expression tag	UNP Q9BRT9
G	-1	GLY	-	expression tag	UNP Q9BRT9
G	0	PRO	-	expression tag	UNP Q9BRT9

• Molecule 14 is a protein called WD repeat and HMG-box DNA-binding protein 1.

Mol	Chain	Residues			Atom	.s			AltConf	Trace
14	Ц	401	Total	С	Η	Ν	0	\mathbf{S}	0	0
14	11	401	6309	2011	3140	552	585	21	0	0
14	т	401	Total	С	Η	Ν	0	S	0	0
14	1	401	6309	2011	3140	552	585	21	0	0
14	т	401	Total	С	Η	Ν	0	S	0	0
14	J	401	6308	2011	3139	552	585	21	0	U

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Н	-31	MET	-	initiating methionine	UNP 075717
Н	-30	ASP	-	expression tag	UNP 075717
Н	-29	TYR	-	expression tag	UNP 075717
Н	-28	LYS	-	expression tag	UNP 075717
H	-27	ASP	-	expression tag	UNP 075717
Н	-26	ASP	-	expression tag	UNP 075717
Н	-25	ASP	-	expression tag	UNP 075717
Н	-24	ASP	-	expression tag	UNP 075717
Н	-23	LYS	-	expression tag	UNP 075717
H	-22	ASP	-	expression tag	UNP 075717
Н	-21	TYR	-	expression tag	UNP 075717
Н	-20	LYS	-	expression tag	UNP 075717
Н	-19	ASP	-	expression tag	UNP 075717
Н	-18	ASP	-	expression tag	UNP 075717
Н	-17	ASP	-	expression tag	UNP 075717
Н	-16	ASP	-	expression tag	UNP 075717
Н	-15	LYS	-	expression tag	UNP 075717
Н	-14	ASP	-	expression tag	UNP 075717
Н	-13	TYR	-	expression tag	UNP 075717
Н	-12	LYS	_	expression tag	UNP 075717



Chain	Residue	Modelled	Actual	Comment	Reference
Н	-11	ASP	-	expression tag	UNP 075717
Н	-10	ASP	_	expression tag	UNP 075717
Н	-9	ASP	-	expression tag	UNP 075717
Н	-8	ASP	-	expression tag	UNP 075717
Н	-7	LYS	-	expression tag	UNP 075717
Н	-6	GLU	-	expression tag	UNP 075717
Н	-5	ASN	-	expression tag	UNP 075717
Н	-4	LEU	-	expression tag	UNP 075717
Н	-3	TYR	-	expression tag	UNP 075717
Н	-2	PHE	-	expression tag	UNP 075717
Н	-1	GLN	-	expression tag	UNP 075717
Н	0	GLY	-	expression tag	UNP 075717
Ι	-31	MET	-	initiating methionine	UNP 075717
Ι	-30	ASP	-	expression tag	UNP 075717
Ι	-29	TYR	-	expression tag	UNP 075717
Ι	-28	LYS	-	expression tag	UNP 075717
Ι	-27	ASP	-	expression tag	UNP 075717
Ι	-26	ASP	-	expression tag	UNP 075717
Ι	-25	ASP	-	expression tag	UNP 075717
Ι	-24	ASP	-	expression tag	UNP 075717
Ι	-23	LYS	-	expression tag	UNP 075717
Ι	-22	ASP	-	expression tag	UNP 075717
Ι	-21	TYR	-	expression tag	UNP 075717
Ι	-20	LYS	-	expression tag	UNP 075717
Ι	-19	ASP	-	expression tag	UNP 075717
Ι	-18	ASP	-	expression tag	UNP 075717
Ι	-17	ASP	-	expression tag	UNP 075717
Ι	-16	ASP	-	expression tag	UNP 075717
Ι	-15	LYS	-	expression tag	UNP 075717
Ι	-14	ASP	-	expression tag	UNP 075717
Ι	-13	TYR	-	expression tag	UNP 075717
Ι	-12	LYS	-	expression tag	UNP 075717
Ι	-11	ASP	-	expression tag	UNP 075717
Ι	-10	ASP	-	expression tag	UNP 075717
Ι	-9	ASP	-	expression tag	UNP 075717
Ι	-8	ASP	-	expression tag	UNP 075717
Ι	-7	LYS	-	expression tag	UNP 075717
Ι	-6	GLU	-	expression tag	UNP 075717
Ι	-5	ASN	-	expression tag	UNP 075717
Ι	-4	LEU	-	expression tag	UNP 075717
Ι	-3	TYR	-	expression tag	UNP 075717
Ι	-2	PHE	-	expression tag	UNP 075717



Chain	Residue	Modelled	Actual	Comment	Reference
Ι	-1	GLN	-	expression tag	UNP 075717
Ι	0	GLY	-	expression tag	UNP 075717
J	-31	MET	-	initiating methionine	UNP 075717
J	-30	ASP	-	expression tag	UNP 075717
J	-29	TYR	-	expression tag	UNP 075717
J	-28	LYS	-	expression tag	UNP 075717
J	-27	ASP	-	expression tag	UNP 075717
J	-26	ASP	-	expression tag	UNP 075717
J	-25	ASP	-	expression tag	UNP 075717
J	-24	ASP	-	expression tag	UNP 075717
J	-23	LYS	-	expression tag	UNP 075717
J	-22	ASP	-	expression tag	UNP 075717
J	-21	TYR	-	expression tag	UNP 075717
J	-20	LYS	-	expression tag	UNP 075717
J	-19	ASP	-	expression tag	UNP 075717
J	-18	ASP	-	expression tag	UNP 075717
J	-17	ASP	-	expression tag	UNP 075717
J	-16	ASP	-	expression tag	UNP 075717
J	-15	LYS	-	expression tag	UNP 075717
J	-14	ASP	-	expression tag	UNP 075717
J	-13	TYR	-	expression tag	UNP 075717
J	-12	LYS	-	expression tag	UNP 075717
J	-11	ASP	-	expression tag	UNP 075717
J	-10	ASP	-	expression tag	UNP 075717
J	-9	ASP	-	expression tag	UNP 075717
J	-8	ASP	-	expression tag	UNP 075717
J	-7	LYS	-	expression tag	UNP 075717
J	-6	GLU	-	expression tag	UNP 075717
J	-5	ASN	-	expression tag	UNP 075717
J	-4	LEU	-	expression tag	UNP 075717
J	-3	TYR	-	expression tag	UNP 075717
J	-2	PHE	-	expression tag	UNP 075717
J	-1	GLN	-	expression tag	UNP 075717
J	0	GLY	-	expression tag	UNP 075717

• Molecule 15 is a protein called Protein timeless homolog.

Mol	Chain	Residues	Atoms					AltConf	Trace	
15	K	636	Total	С	Н	N	0	S	0	0
			10448	3324	5233	929	936	26		

There is a discrepancy between the modelled and reference sequences:



Chain	Residue	Modelled	Actual	Comment	Reference
K	0	GLY	-	expression tag	UNP Q9UNS1

• Molecule 16 is a protein called TIMELESS-interacting protein.

Mol	Chain	Residues	Atoms				AltConf	Trace		
16	L	87	Total 1502	C 471	Н 767	N 140	0 121	${ m S} { m 3}$	0	0

• Molecule 17 is a DNA chain called Leading strand DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace	
17	М	32	Total	C	H	N 100	0	Р	0	0
			1032	320	370	106	204	32		

• Molecule 18 is a DNA chain called Lagging Strand DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace	
18	Ν	22	Total 686	C 211	Н 248	N 74	0 131	Р 22	0	0

• Molecule 19 is a protein called Claspin.

Mol	Chain	Residues	Atoms				AltConf	Trace		
19	Q	79	Total 1387	C 431	Н 704	N 133	0 117	${ m S} { m 2}$	0	0

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Q	1340	LEU	-	expression tag	UNP Q9HAW4
Q	1341	GLU	-	expression tag	UNP Q9HAW4
Q	1342	VAL	-	expression tag	UNP Q9HAW4
Q	1343	LEU	-	expression tag	UNP Q9HAW4
Q	1344	PHE	-	expression tag	UNP Q9HAW4
Q	1345	GLN	-	expression tag	UNP Q9HAW4
Q	1346	GLY	-	expression tag	UNP Q9HAW4
Q	1347	PRO	-	expression tag	UNP Q9HAW4
Q	1348	ASP	-	expression tag	UNP Q9HAW4
Q	1349	TYR	-	expression tag	UNP Q9HAW4
Q	1350	LYS	-	expression tag	UNP Q9HAW4
Q	1351	ASP	-	expression tag	UNP Q9HAW4
Q	1352	ASP	-	expression tag	UNP Q9HAW4



Chain	Residue	Modelled	Actual	Comment	Reference
Q	1353	ASP	-	expression tag	UNP Q9HAW4
Q	1354	ASP	-	expression tag	UNP Q9HAW4
Q	1355	LYS	-	expression tag	UNP Q9HAW4
Q	1356	ASP	-	expression tag	UNP Q9HAW4
Q	1357	TYR	-	expression tag	UNP Q9HAW4
Q	1358	LYS	-	expression tag	UNP Q9HAW4
Q	1359	ASP	-	expression tag	UNP Q9HAW4
Q	1360	ASP	-	expression tag	UNP Q9HAW4
Q	1361	ASP	-	expression tag	UNP Q9HAW4
Q	1362	ASP	-	expression tag	UNP Q9HAW4
Q	1363	LYS	-	expression tag	UNP Q9HAW4
Q	1364	ASP	-	expression tag	UNP Q9HAW4
Q	1365	TYR	-	expression tag	UNP Q9HAW4
Q	1366	LYS	-	expression tag	UNP Q9HAW4
Q	1367	ASP	-	expression tag	UNP Q9HAW4
Q	1368	ASP	-	expression tag	UNP Q9HAW4
Q	1369	ASP	-	expression tag	UNP Q9HAW4
Q	1370	ASP	-	expression tag	UNP Q9HAW4
Q	1371	LYS	-	expression tag	UNP Q9HAW4

• Molecule 20 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	AltConf
20	2	1	Total Zn 1 1	0
20	4	1	Total Zn 1 1	0
20	5	1	Total Zn 1 1	0
20	6	1	Total Zn 1 1	0
20	7	1	Total Zn 1 1	0

• Molecule 21 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	AltConf
21	2	1	Total Mg 1 1	0



Mol	Chain	Residues	Atoms	AltConf
21	3	1	Total Mg 1 1	0
21	5	1	Total Mg 1 1	0

• Molecule 22 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (threeletter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						AltConf
22	2	1	Total	С	Η	Ν	Ο	Р	0
	2	1	44	10	13	6	12	3	0
22	<u></u>	1	Total	С	Η	Ν	Ο	Р	0
	5	1	44	10	13	6	12	3	0
22	5	1	Total	С	Η	Ν	Ο	Р	0
	5		44	10	13	6	12	3	U

• Molecule 23 is SULFATE ION (three-letter code: SO4) (formula: O₄S) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	AltConf
23	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA replication licensing factor MCM2



Chain 4:







GLU ARGEN VAL ARGE CYSS CYSS CYSS CYSS ARGEN PHE PHE CITRE C

• Molecule 7: DNA polymerase epsilon subunit 2



























• Molecule 16: TIMELESS-interacting protein







4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	110000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	39.8	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	3500	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.062	Depositor
Minimum map value	-0.017	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.0113	Depositor
Map size (Å)	470.80002, 470.80002, 470.80002	wwPDB
Map dimensions	440, 440, 440	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.07, 1.07, 1.07	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ANP, ZN, MG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bo	nd lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	2	0.32	0/5683	0.63	0/7675	
2	3	0.30	0/4931	0.64	0/6653	
3	4	0.29	0/5013	0.61	0/6776	
4	5	0.29	0/4807	0.64	0/6468	
5	6	0.30	0/5095	0.62	0/6876	
6	7	0.33	0/4822	0.67	1/6507~(0.0%)	
7	А	0.32	0/4310	0.62	0/5853	
8	В	0.36	0/6647	0.73	1/9011~(0.0%)	
9	С	0.28	0/4439	0.57	0/5992	
10	D	0.31	0/1638	0.65	0/2202	
11	Е	0.29	0/1462	0.57	0/1981	
12	F	0.29	0/1580	0.58	0/2133	
13	G	0.29	0/1711	0.61	0/2305	
14	Н	0.30	0/3244	0.61	0/4395	
14	Ι	0.30	0/3244	0.60	0/4395	
14	J	0.28	0/3244	0.60	0/4395	
15	Κ	0.30	0/5317	0.63	0/7162	
16	L	0.28	0/750	0.67	0/999	
17	М	0.86	2/738~(0.3%)	1.13	0/1138	
18	N	0.90	1/488~(0.2%)	1.04	0/747	
19	Q	0.29	0/693	0.65	0/919	
All	All	0.33	3/69856~(0.0%)	0.65	2/94582~(0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
6	7	0	1



Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
17	М	55	DT	C1'-N1	5.21	1.56	1.49
18	Ν	25	DC	C1'-N1	5.18	1.55	1.49
17	М	58	DT	C1'-N1	5.17	1.55	1.49

All (3) bond length outliers are listed below:

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
6	7	385	VAL	CA-CB-CG1	6.74	121.00	110.90
8	В	1366	PHE	CB-CG-CD1	5.35	124.54	120.80

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
6	7	407	ARG	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	2	5580	5597	5590	10	0
2	3	4858	4912	4908	11	0
3	4	4929	4993	4985	6	0
4	5	4735	4809	4800	4	0
5	6	5014	5051	5043	4	0
6	7	4750	4802	4793	6	0
7	А	4199	4160	4160	5	0
8	В	6497	6478	6471	13	0
9	С	4347	4300	4296	5	0
10	D	1606	1601	1601	1	0
11	Е	1431	1456	1456	1	0
12	F	1546	1502	1500	2	0
13	G	1679	1702	1700	3	0
14	Н	3169	3140	3138	2	0
14	Ι	3169	3140	3138	3	0
14	J	3169	3139	3138	2	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
15	K	5215	5233	5230	3	0
16	L	735	767	766	0	0
17	М	662	370	372	0	0
18	N	438	248	249	0	0
19	Q	683	704	701	1	0
20	2	1	0	0	0	0
20	4	1	0	0	0	0
20	5	1	0	0	0	0
20	6	1	0	0	0	0
20	7	1	0	0	0	0
21	2	1	0	0	0	0
21	3	1	0	0	0	0
21	5	1	0	0	0	0
22	2	31	13	13	1	0
22	3	31	13	13	3	0
22	5	31	13	13	1	0
23	A	5	0	0	0	0
All	All	68517	68143	68074	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 75 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:2:290:LEU:HD12	1:2:290:LEU:O	1.59	1.02
3:4:674:GLU:OE2	3:4:675:GLU:HG3	1.76	0.85
8:B:1813:TYR:CE1	8:B:1817:GLN:HG3	2.19	0.78
6:7:385:VAL:CG2	6:7:523:ASP:HB2	2.16	0.76
6:7:385:VAL:HG23	6:7:523:ASP:HB2	1.77	0.65

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	\mathbf{ntiles}
1	2	697/904~(77%)	671~(96%)	25~(4%)	1 (0%)	51	83
2	3	614/808~(76%)	594 (97%)	20 (3%)	0	100	100
3	4	611/863~(71%)	594 (97%)	17 (3%)	0	100	100
4	5	597/734~(81%)	580~(97%)	17 (3%)	0	100	100
5	6	624/821~(76%)	607~(97%)	17 (3%)	0	100	100
6	7	586/719~(82%)	572 (98%)	14 (2%)	0	100	100
7	А	525/527~(100%)	503~(96%)	22~(4%)	0	100	100
8	В	792/2286~(35%)	706 (89%)	82 (10%)	4 (0%)	29	67
9	С	530/569~(93%)	522 (98%)	8 (2%)	0	100	100
10	D	193/196~(98%)	191 (99%)	2(1%)	0	100	100
11	Е	174/185~(94%)	172 (99%)	2 (1%)	0	100	100
12	F	190/216~(88%)	185 (97%)	5 (3%)	0	100	100
13	G	201/262~(77%)	200 (100%)	1 (0%)	0	100	100
14	Н	399/1161~(34%)	385~(96%)	14 (4%)	0	100	100
14	Ι	399/1161~(34%)	392 (98%)	7 (2%)	0	100	100
14	J	399/1161~(34%)	390~(98%)	9(2%)	0	100	100
15	Κ	630/1209~(52%)	615~(98%)	15 (2%)	0	100	100
16	L	85/301~(28%)	81 (95%)	4 (5%)	0	100	100
19	Q	73/1371~(5%)	73 (100%)	0	0	100	100
All	All	8319/15454 (54%)	8033 (97%)	281 (3%)	5 (0%)	54	83

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
8	В	1431	VAL
8	В	1580	ARG
8	В	1656	PRO
1	2	713	GLU
8	В	1918	ASP



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	2	618/781~(79%)	613~(99%)	5 (1%)	81	93
2	3	534/707~(76%)	531~(99%)	3~(1%)	86	94
3	4	549/753~(73%)	545~(99%)	4 (1%)	84	94
4	5	515/625~(82%)	513 (100%)	2~(0%)	91	95
5	6	556/724~(77%)	552~(99%)	4 (1%)	84	94
6	7	518/619~(84%)	513~(99%)	5 (1%)	76	90
7	А	471/471~(100%)	466 (99%)	5(1%)	73	88
8	В	725/2012~(36%)	709~(98%)	16~(2%)	52	79
9	\mathbf{C}	487/520~(94%)	487 (100%)	0	100	100
10	D	173/174~(99%)	173~(100%)	0	100	100
11	Ε	160/169~(95%)	160 (100%)	0	100	100
12	\mathbf{F}	167/186~(90%)	163~(98%)	4(2%)	49	77
13	G	188/233~(81%)	188 (100%)	0	100	100
14	Η	345/1018~(34%)	343~(99%)	2(1%)	86	94
14	Ι	345/1018~(34%)	341~(99%)	4 (1%)	71	88
14	J	345/1018~(34%)	341~(99%)	4 (1%)	71	88
15	Κ	561/1055~(53%)	557~(99%)	4 (1%)	84	94
16	L	78/274~(28%)	76~(97%)	2(3%)	46	76
19	Q	72/1230~(6%)	72 (100%)	0	100	100
All	All	7407/13587 (54%)	7343~(99%)	64 (1%)	79	91

5 of 64 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
14	J	649	MET
15	Κ	181	ASP
7	А	165	LEU
7	А	151	VAL



 $Continued \ from \ previous \ page...$

Mol	Chain	\mathbf{Res}	Type
15	Κ	480	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
14	Н	736	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	Dec	Tink	Bo	ond leng	$_{\rm sths}$	B	ond ang	les
INIOI	туре	Unain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
22	ANP	3	1500	21	29,33,33	1.01	2 (6%)	31,52,52	1.09	3 (9%)
22	ANP	2	1003	21	29,33,33	1.12	4 (13%)	31,52,52	0.99	2 (6%)
23	SO4	А	601	-	4,4,4	0.14	0	6,6,6	0.05	0
22	ANP	5	802	21	29,33,33	1.10	4 (13%)	31,52,52	1.09	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral



centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
22	ANP	3	1500	21	-	3/14/38/38	0/3/3/3
22	ANP	2	1003	21	-	3/14/38/38	0/3/3/3
22	ANP	5	802	21	-	5/14/38/38	0/3/3/3

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\operatorname{Ideal}(\operatorname{\AA})$
22	3	1500	ANP	PB-O3A	-3.40	1.54	1.59
22	2	1003	ANP	PB-O3A	-3.25	1.55	1.59
22	5	802	ANP	PB-O3A	-2.90	1.55	1.59
22	5	802	ANP	PG-01G	2.39	1.49	1.46
22	5	802	ANP	PG-N3B	2.31	1.69	1.63

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms		$Observed(^{o})$	$Ideal(^{o})$
22	5	802	ANP	PB-O3A-PA	-4.06	118.30	132.62
22	3	1500	ANP	PB-O3A-PA	-3.58	120.00	132.62
22	2	1003	ANP	PB-O3A-PA	-3.36	120.80	132.62
22	3	1500	ANP	O1G-PG-N3B	-2.87	107.54	111.77
22	5	802	ANP	C5-C6-N6	2.35	123.93	120.35

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	Atoms
22	2	1003	ANP	PB-N3B-PG-O1G
22	2	1003	ANP	PA-O3A-PB-O1B
22	2	1003	ANP	PA-O3A-PB-O2B
22	3	1500	ANP	PB-N3B-PG-O1G
22	3	1500	ANP	PA-O3A-PB-O1B

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
22	3	1500	ANP	3	0
22	2	1003	ANP	1	0



Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
22	5	802	ANP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.















5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-13375. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

Orthogonal projections (i) 6.1

6.1.1**Primary** map



6.1.2Raw map



The images above show the map projected in three orthogonal directions.



6.2 Central slices (i)

6.2.1 Primary map



X Index: 220



Y Index: 220



Z Index: 220

6.2.2 Raw map



X Index: 220

Y Index: 220



The images above show central slices of the map in three orthogonal directions.



6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 200



Y Index: 202



Z Index: 196

6.3.2 Raw map



X Index: 199

Y Index: 202



The images above show the largest variance slices of the map in three orthogonal directions.



6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.



6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0113. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.



Mask visualisation (i) 6.6

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency

A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure

$emd_{13375}msk_{1.map}$ (i) 6.6.1





7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)



The volume at the recommended contour level is 763 $\rm nm^3;$ this corresponds to an approximate mass of 689 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.312 ${\rm \AA^{-1}}$



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.312 ${\rm \AA^{-1}}$



8.2 Resolution estimates (i)

$\mathbf{Bosolution} \text{ ostimato } (\mathbf{\hat{\lambda}})$	Estimation criterion (FSC cut-off)		
Resolution estimate (A)	0.143	0.5	Half-bit
Reported by author	3.20	-	-
Author-provided FSC curve	3.18	3.62	3.22
Unmasked-calculated*	4.05	7.43	4.15

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.05 differs from the reported value 3.2 by more than 10 %



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-13375 and PDB model 7PFO. Per-residue inclusion information can be found in section 3 on page 14.

9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.0113 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.



9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0113).



9.4 Atom inclusion (i)



At the recommended contour level, 85% of all backbone atoms, 86% of all non-hydrogen atoms, are inside the map.



1.0

0.0 <0.0

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.0113) and Q-score for the entire model and for each chain.

\mathbf{Chain}	Atom inclusion	Q-score
All	0.8550	0.3990
2	0.8940	0.4650
3	0.9210	0.4690
4	0.9260	0.4440
5	0.9050	0.4930
6	0.9250	0.4650
7	0.9150	0.4240
A	0.8090	0.3140
В	0.6880	0.2300
С	0.9150	0.4820
D	0.9110	0.4700
Ε	0.9230	0.5120
F	0.9370	0.4900
G	0.9050	0.4760
Н	0.7360	0.2340
Ι	0.7870	0.3680
J	0.6880	0.2050
K	0.9140	0.4170
L	0.8940	0.4220
М	0.8820	0.3500
Ν	0.9040	0.2830
Q	0.7500	0.3280

