



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 17, 2022 – 04:16 pm BST

PDB ID : 7PGO  
Title : Crystal Structure of a Class D Carbapenemase\_R250A  
Authors : Zhou, Q.; He, Y.; Jin, Y.  
Deposited on : 2021-08-15  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.29  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.29


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2469 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

MolProbity failed to run properly - the sequence quality summary graphics cannot be shown.

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 16884 atoms, of which 7929 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	AAA	244	3978	1278	1973	350	370	7	50	4	0
1	BBB	244	3992	1282	1981	352	368	9	48	4	0
1	CCC	243	3988	1282	1977	353	369	7	50	4	0
1	DDD	244	3957	1273	1958	349	370	7	48	2	0

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	6	MET	-	initiating methionine	UNP Q6XEC0
AAA	7	HIS	-	expression tag	UNP Q6XEC0
AAA	8	HIS	-	expression tag	UNP Q6XEC0
AAA	9	HIS	-	expression tag	UNP Q6XEC0
AAA	10	HIS	-	expression tag	UNP Q6XEC0
AAA	11	HIS	-	expression tag	UNP Q6XEC0
AAA	12	HIS	-	expression tag	UNP Q6XEC0
AAA	13	SER	-	expression tag	UNP Q6XEC0
AAA	14	ALA	-	expression tag	UNP Q6XEC0
AAA	15	GLY	-	expression tag	UNP Q6XEC0
AAA	16	GLU	-	expression tag	UNP Q6XEC0
AAA	17	ASN	-	expression tag	UNP Q6XEC0
AAA	18	LEU	-	expression tag	UNP Q6XEC0
AAA	19	TYR	-	expression tag	UNP Q6XEC0
AAA	20	PHE	-	expression tag	UNP Q6XEC0
AAA	21	GLN	-	expression tag	UNP Q6XEC0
AAA	22	GLY	-	expression tag	UNP Q6XEC0
AAA	250	ALA	ARG	engineered mutation	UNP Q6XEC0
BBB	6	MET	-	initiating methionine	UNP Q6XEC0
BBB	7	HIS	-	expression tag	UNP Q6XEC0
BBB	8	HIS	-	expression tag	UNP Q6XEC0

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
BBB	9	HIS	-	expression tag	UNP Q6XEC0
BBB	10	HIS	-	expression tag	UNP Q6XEC0
BBB	11	HIS	-	expression tag	UNP Q6XEC0
BBB	12	HIS	-	expression tag	UNP Q6XEC0
BBB	13	SER	-	expression tag	UNP Q6XEC0
BBB	14	ALA	-	expression tag	UNP Q6XEC0
BBB	15	GLY	-	expression tag	UNP Q6XEC0
BBB	16	GLU	-	expression tag	UNP Q6XEC0
BBB	17	ASN	-	expression tag	UNP Q6XEC0
BBB	18	LEU	-	expression tag	UNP Q6XEC0
BBB	19	TYR	-	expression tag	UNP Q6XEC0
BBB	20	PHE	-	expression tag	UNP Q6XEC0
BBB	21	GLN	-	expression tag	UNP Q6XEC0
BBB	22	GLY	-	expression tag	UNP Q6XEC0
BBB	250	ALA	ARG	engineered mutation	UNP Q6XEC0
CCC	6	MET	-	initiating methionine	UNP Q6XEC0
CCC	7	HIS	-	expression tag	UNP Q6XEC0
CCC	8	HIS	-	expression tag	UNP Q6XEC0
CCC	9	HIS	-	expression tag	UNP Q6XEC0
CCC	10	HIS	-	expression tag	UNP Q6XEC0
CCC	11	HIS	-	expression tag	UNP Q6XEC0
CCC	12	HIS	-	expression tag	UNP Q6XEC0
CCC	13	SER	-	expression tag	UNP Q6XEC0
CCC	14	ALA	-	expression tag	UNP Q6XEC0
CCC	15	GLY	-	expression tag	UNP Q6XEC0
CCC	16	GLU	-	expression tag	UNP Q6XEC0
CCC	17	ASN	-	expression tag	UNP Q6XEC0
CCC	18	LEU	-	expression tag	UNP Q6XEC0
CCC	19	TYR	-	expression tag	UNP Q6XEC0
CCC	20	PHE	-	expression tag	UNP Q6XEC0
CCC	21	GLN	-	expression tag	UNP Q6XEC0
CCC	22	GLY	-	expression tag	UNP Q6XEC0
CCC	250	ALA	ARG	engineered mutation	UNP Q6XEC0
DDD	6	MET	-	initiating methionine	UNP Q6XEC0
DDD	7	HIS	-	expression tag	UNP Q6XEC0
DDD	8	HIS	-	expression tag	UNP Q6XEC0
DDD	9	HIS	-	expression tag	UNP Q6XEC0
DDD	10	HIS	-	expression tag	UNP Q6XEC0
DDD	11	HIS	-	expression tag	UNP Q6XEC0
DDD	12	HIS	-	expression tag	UNP Q6XEC0
DDD	13	SER	-	expression tag	UNP Q6XEC0
DDD	14	ALA	-	expression tag	UNP Q6XEC0

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
DDD	15	GLY	-	expression tag	UNP Q6XEC0
DDD	16	GLU	-	expression tag	UNP Q6XEC0
DDD	17	ASN	-	expression tag	UNP Q6XEC0
DDD	18	LEU	-	expression tag	UNP Q6XEC0
DDD	19	TYR	-	expression tag	UNP Q6XEC0
DDD	20	PHE	-	expression tag	UNP Q6XEC0
DDD	21	GLN	-	expression tag	UNP Q6XEC0
DDD	22	GLY	-	expression tag	UNP Q6XEC0
DDD	250	ALA	ARG	engineered mutation	UNP Q6XEC0

- Molecule 2 is 1-BUTANOL (three-letter code: 1BO) (formula: C<sub>4</sub>H<sub>10</sub>O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	AAA	1	Total	C	H	O	0	0
			15	4	10	1		
2	AAA	1	Total	C	H	O	0	0
			15	4	10	1		
2	BBB	1	Total	C	H	O	0	0
			15	4	10	1		
2	DDD	1	Total	C	H	O	0	0
			15	4	10	1		

- Molecule 3 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	AAA	2	Total 3	Br 3	0	1
3	BBB	3	Total 4	Br 4	0	1

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	227	Total 227	O 227	0	0
4	BBB	249	Total 249	O 249	0	0
4	CCC	224	Total 224	O 224	0	0
4	DDD	201	Total 202	O 202	0	1

MolProbity failed to run properly - this section is therefore empty.

### 3 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.55Å 108.20Å 124.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	58.75 – 1.85 58.75 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.9 (58.75-1.85) 99.9 (58.75-1.85)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.86 (at 1.86Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.205 , 0.234 0.211 , 0.241	Depositor DCC
$R_{free}$ test set	5022 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.2	Xtrriage
Anisotropy	0.890	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	16884	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 49.63 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.2588e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 4 Model quality [i](#)

### 4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.3 Torsion angles [i](#)

#### 4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	KCX	CCC	73	1	9,11,12	0.67	0	5,12,14	0.34	0
1	KCX	AAA	73	1	9,11,12	0.65	0	5,12,14	0.54	0
1	KCX	BBB	73	1	9,11,12	0.70	0	5,12,14	0.50	0
1	KCX	DDD	73	1	9,11,12	0.72	0	5,12,14	0.33	0



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	CCC	73	1	-	0/9/10/12	-
1	KCX	AAA	73	1	-	0/9/10/12	-
1	KCX	BBB	73	1	-	0/9/10/12	-
1	KCX	DDD	73	1	-	0/9/10/12	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

#### 4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

#### 4.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 7 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	1BO	DDD	301	-	4,4,4	0.18	0	3,3,3	0.08	0
2	1BO	BBB	301	-	4,4,4	0.31	0	3,3,3	0.35	0
2	1BO	AAA	301	-	4,4,4	0.22	0	3,3,3	0.41	0
2	1BO	AAA	302	-	4,4,4	0.27	0	3,3,3	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1BO	DDD	301	-	-	0/2/2/2	-
2	1BO	BBB	301	-	-	0/2/2/2	-
2	1BO	AAA	301	-	-	1/2/2/2	-
2	1BO	AAA	302	-	-	0/2/2/2	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	AAA	301	1BO	C2-C3-C4-OH

There are no ring outliers.

No monomer is involved in short contacts.

#### 4.7 Other polymers [i](#)

There are no such residues in this entry.

#### 4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 5 Fit of model and data

### 5.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	AAA	243/260 (93%)	-0.39	1 (0%) 92 92	11, 18, 34, 64	1 (0%)
1	BBB	243/260 (93%)	-0.37	3 (1%) 79 79	11, 17, 41, 71	1 (0%)
1	CCC	242/260 (93%)	-0.24	7 (2%) 51 50	11, 19, 46, 85	2 (0%)
1	DDD	243/260 (93%)	-0.15	9 (3%) 41 39	11, 19, 48, 102	6 (2%)
All	All	971/1040 (93%)	-0.29	20 (2%) 63 63	11, 18, 44, 102	10 (1%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	DDD	247	LEU	7.0
1	CCC	245	ASP	6.4
1	CCC	244	SER	6.0
1	DDD	242	PRO	5.9
1	DDD	245	ASP	5.8
1	CCC	247	LEU	5.5
1	DDD	244	SER	4.4
1	DDD	243	THR	4.2
1	DDD	246	GLY	3.1
1	DDD	23	LYS	2.8
1	DDD	25	TRP	2.7
1	CCC	246	GLY	2.6
1	CCC	242	PRO	2.5
1	BBB	247	LEU	2.3
1	CCC	23[A]	LYS	2.3
1	AAA	22	GLY	2.3
1	BBB	245	ASP	2.2
1	DDD	182	HIS	2.2
1	BBB	246	GLY	2.1
1	CCC	25	TRP	2.1

## 5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	KCX	CCC	73	12/13	0.95	0.10	15,19,21,23	1
1	KCX	BBB	73	12/13	0.96	0.09	12,14,23,25	9
1	KCX	AAA	73	12/13	0.96	0.09	11,15,20,20	8
1	KCX	DDD	73	12/13	0.96	0.11	15,20,23,25	1

## 5.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	1BO	AAA	302	5/5	0.73	0.26	25,27,28,29	15
2	1BO	BBB	301	5/5	0.75	0.20	31,35,37,37	0
2	1BO	AAA	301	5/5	0.81	0.21	29,30,30,32	15
2	1BO	DDD	301	5/5	0.81	0.16	29,31,34,35	15
3	BR	BBB	304	1/1	0.90	0.09	42,42,42,42	1
3	BR	AAA	304[B]	1/1	0.94	0.21	34,34,34,34	1
3	BR	AAA	304[A]	1/1	0.94	0.21	42,42,42,42	1
3	BR	BBB	303[B]	1/1	0.96	0.20	33,33,33,33	1
3	BR	BBB	303[A]	1/1	0.96	0.20	37,37,37,37	1
3	BR	BBB	302	1/1	0.99	0.06	21,21,21,21	0
3	BR	AAA	303	1/1	0.99	0.05	23,23,23,23	0

## 5.5 Other polymers [i](#)

There are no such residues in this entry.