



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 22, 2022 – 03:10 pm GMT

PDB ID : 7PH1
Title : Trypsin in complex with BPTI mutant (2S)-2-amino-4-monofluorobutanoic acid
Authors : Dimos, N.; Leppkes, J.; Kokschi, B.; Loll, B.
Deposited on : 2021-08-16
Resolution : 1.18 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.27
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.27


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.18 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1123 (1.20-1.16)
Clashscore	141614	1182 (1.20-1.16)
Ramachandran outliers	138981	1134 (1.20-1.16)
Sidechain outliers	138945	1134 (1.20-1.16)
RSRZ outliers	127900	1102 (1.20-1.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	223	
2	I	57	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	I	107[A]	-	X	-	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5269 atoms, of which 2475 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cationic trypsin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	E	223	3693	1141	1872	304	362	14	0	43	0


- Molecule 2 is a protein called Pancreatic trypsin inhibitor.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace	
			Total	C	F	H	N	O				S
2	I	57	1029	324	1	515	94	87	8	0	11	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
I	15	7OZ	LYS	conflict	UNP P00974

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	E	1	Total	O	S	0	0
			5	4	1		
3	E	1	Total	O	S	0	0
			5	4	1		
3	E	1	Total	O	S	0	0
			5	4	1		
3	E	1	Total	O	S	0	1
			10	8	2		
3	E	1	Total	O	S	0	0
			5	4	1		
3	E	1	Total	O	S	0	1
			5	4	1		
3	E	1	Total	O	S	0	0
			5	4	1		
3	E	1	Total	O	S	0	1
			5	4	1		
3	E	1	Total	O	S	0	0
			5	4	1		
3	I	1	Total	O	S	0	0
			5	4	1		
3	I	1	Total	O	S	0	0
			5	4	1		
3	I	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	E	1	Total	C	H	O	0	1
			28	6	16	6		
4	E	1	Total	C	H	O	0	0
			14	3	8	3		
4	E	1	Total	C	H	O	0	0
			14	3	8	3		
4	E	1	Total	C	H	O	0	0
			14	3	8	3		
4	E	1	Total	C	H	O	0	0
			14	3	8	3		
4	E	1	Total	C	H	O	0	1
			28	6	16	6		
4	I	1	Total	C	H	O	0	1
			14	3	8	3		
4	I	1	Total	C	H	O	0	0
			14	3	8	3		
4	I	1	Total	C	H	O	0	1
			14	3	8	3		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	E	1	Total	Ca	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	E	237	Total	O	0	27
			237	237		
6	I	85	Total	O	0	22
			85	85		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cationic trypsin



- Molecule 2: Pancreatic trypsin inhibitor



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	74.97Å 81.29Å 124.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.56 – 1.18 41.23 – 1.18	Depositor EDS
% Data completeness (in resolution range)	100.0 (27.56-1.18) 99.8 (41.23-1.18)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.06 (at 1.18Å)	Xtriage
Refinement program	PHENIX 1.19.1_4122	Depositor
R, R_{free}	0.149 , 0.166 0.150 , 0.166	Depositor DCC
R_{free} test set	1812 reflections (1.46%)	wwPDB-VP
Wilson B-factor (Å ²)	15.8	Xtriage
Anisotropy	0.370	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	5269	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, 7OZ, SO4, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	E	0.63	0/2009	0.80	1/2718 (0.0%)
2	I	0.85	2/536 (0.4%)	1.00	3/714 (0.4%)
All	All	0.68	2/2545 (0.1%)	0.85	4/3432 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	31[A]	GLN	CG-CD	5.34	1.63	1.51
2	I	31[B]	GLN	CG-CD	5.34	1.63	1.51

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	52[A]	MET	CG-SD-CE	-7.24	88.61	100.20
2	I	52[B]	MET	CG-SD-CE	-7.24	88.61	100.20
1	E	191	ASP	CB-CG-OD1	5.79	123.51	118.30
2	I	42	ARG	NE-CZ-NH1	5.02	122.81	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1821	1872	1818	5	0
2	I	514	515	494	2	0
3	E	50	0	0	0	0
3	I	20	0	0	0	0
4	E	48	64	60	3	0
4	I	18	24	23	0	0
5	E	1	0	0	0	0
6	E	237	0	0	2	0
6	I	85	0	0	0	0
All	All	2794	2475	2395	8	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (8) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:47:ILE:HD11	1:E:240:ILE:HD11	1.82	0.61
4:E:305:GOL:H32	6:E:403:HOH:O	2.01	0.59
1:E:197:SER:OG	2:I:15:7OZ:C	2.60	0.49
2:I:29[A]:LEU:CD1	2:I:31[A]:GLN:HG3	2.43	0.49
4:E:302[B]:GOL:H11	6:E:444:HOH:O	2.15	0.46
1:E:79:ASN:CG	1:E:117[B]:ARG:HD3	2.38	0.44
1:E:25:ASN:CB	1:E:117[A]:ARG:HD3	2.48	0.44
1:E:50[B]:GLN:HB2	4:E:307[B]:GOL:H31	2.01	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	E	266/223 (119%)	264 (99%)	2 (1%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	I	65/57 (114%)	63 (97%)	2 (3%)	0	100	100
All	All	331/280 (118%)	327 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	226/184 (123%)	226 (100%)	0	100	100
2	I	53/44 (120%)	51 (96%)	2 (4%)	33	4
All	All	279/228 (122%)	277 (99%)	2 (1%)	91	57

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	I	29[A]	LEU
2	I	29[B]	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	7OZ	I	15	2	5,6,7	0.68	0	1,6,8	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	7OZ	I	15	2	-	0/3/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	I	15	7OZ	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 26 ligands modelled in this entry, 1 is monoatomic - leaving 25 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	I	107[A]	-	5,5,5	2.32	2 (40%)	5,5,5	1.55	1 (20%)
4	GOL	E	303	-	5,5,5	1.12	1 (20%)	5,5,5	1.05	0
3	SO4	E	308	-	4,4,4	0.75	0	6,6,6	0.85	0
3	SO4	I	106	-	4,4,4	0.42	0	6,6,6	0.20	0
3	SO4	E	311	-	4,4,4	0.24	0	6,6,6	0.20	0
3	SO4	E	316	-	4,4,4	0.44	0	6,6,6	1.24	1 (16%)
3	SO4	I	105	-	4,4,4	0.60	0	6,6,6	0.84	0
4	GOL	E	307[B]	-	5,5,5	2.05	1 (20%)	5,5,5	0.54	0
4	GOL	E	302[B]	-	5,5,5	1.04	0	5,5,5	1.08	0
4	GOL	E	307[A]	-	5,5,5	1.48	2 (40%)	5,5,5	0.55	0
4	GOL	I	101[B]	-	5,5,5	1.05	0	5,5,5	1.46	1 (20%)
4	GOL	E	302[A]	-	5,5,5	1.78	1 (20%)	5,5,5	0.95	0
4	GOL	E	304	-	5,5,5	1.09	1 (20%)	5,5,5	1.49	1 (20%)
4	GOL	E	306	-	5,5,5	1.32	1 (20%)	5,5,5	0.57	0
3	SO4	E	315[A]	-	4,4,4	0.21	0	6,6,6	0.39	0
3	SO4	E	310[B]	-	4,4,4	0.15	0	6,6,6	0.21	0
3	SO4	E	310[A]	-	4,4,4	0.38	0	6,6,6	0.41	0
3	SO4	I	103	-	4,4,4	0.30	0	6,6,6	1.00	0
3	SO4	E	309	-	4,4,4	0.22	0	6,6,6	0.34	0
3	SO4	E	301	-	4,4,4	0.38	0	6,6,6	0.70	0
3	SO4	E	313[A]	-	4,4,4	0.18	0	6,6,6	0.37	0
3	SO4	E	314	-	4,4,4	0.20	0	6,6,6	0.66	0
4	GOL	E	305	-	5,5,5	1.56	1 (20%)	5,5,5	0.72	0
3	SO4	I	104	-	4,4,4	0.20	0	6,6,6	0.26	0
4	GOL	I	102	-	5,5,5	1.22	0	5,5,5	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	E	304	-	-	2/4/4/4	-
4	GOL	E	306	-	-	2/4/4/4	-
4	GOL	I	107[A]	-	-	4/4/4/4	-
4	GOL	E	305	-	-	0/4/4/4	-
4	GOL	E	303	-	-	2/4/4/4	-
4	GOL	E	307[B]	-	-	0/4/4/4	-
4	GOL	I	102	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	E	302[B]	-	-	2/4/4/4	-
4	GOL	E	307[A]	-	-	0/4/4/4	-
4	GOL	I	101[B]	-	-	2/4/4/4	-
4	GOL	E	302[A]	-	-	2/4/4/4	-

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	I	107[A]	GOL	C3-C2	4.62	1.70	1.51
4	E	302[A]	GOL	O2-C2	-3.69	1.32	1.43
4	E	307[B]	GOL	C3-C2	3.66	1.66	1.51
4	E	305	GOL	C3-C2	2.83	1.63	1.51
4	E	307[A]	GOL	C3-C2	2.29	1.61	1.51
4	E	306	GOL	O2-C2	-2.21	1.36	1.43
4	E	307[A]	GOL	C1-C2	2.12	1.60	1.51
4	E	303	GOL	C1-C2	2.11	1.60	1.51
4	E	304	GOL	C1-C2	2.10	1.60	1.51
4	I	107[A]	GOL	C1-C2	2.01	1.60	1.51

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	I	107[A]	GOL	O2-C2-C3	2.81	121.48	109.12
4	E	304	GOL	C3-C2-C1	-2.34	102.61	111.70
3	E	316	SO4	O3-S-O2	-2.25	97.55	109.31
4	I	101[B]	GOL	O2-C2-C1	2.09	118.32	109.12

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	E	302[A]	GOL	O1-C1-C2-O2
4	E	302[A]	GOL	O1-C1-C2-C3
4	E	302[B]	GOL	O1-C1-C2-C3
4	E	303	GOL	C1-C2-C3-O3
4	I	107[A]	GOL	C1-C2-C3-O3
4	E	306	GOL	C1-C2-C3-O3
4	I	101[B]	GOL	O1-C1-C2-C3
4	I	102	GOL	C1-C2-C3-O3
4	I	107[A]	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	E	303	GOL	O2-C2-C3-O3
4	E	306	GOL	O2-C2-C3-O3
4	I	107[A]	GOL	O2-C2-C3-O3
4	I	102	GOL	O2-C2-C3-O3
4	I	107[A]	GOL	O1-C1-C2-O2
4	E	302[B]	GOL	O1-C1-C2-O2
4	E	304	GOL	O1-C1-C2-O2
4	E	304	GOL	O1-C1-C2-C3
4	I	101[B]	GOL	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	307[B]	GOL	1	0
4	E	302[B]	GOL	1	0
4	E	305	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	E	223/223 (100%)	0.73	17 (7%) 13 14	14, 21, 28, 38	0
2	I	56/57 (98%)	1.17	9 (16%) 1 2	13, 16, 29, 39	0
All	All	279/280 (99%)	0.82	26 (9%) 8 10	13, 20, 29, 39	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	147[A]	SER	6.0
1	E	125[A]	THR	4.8
1	E	79	ASN	4.7
2	I	2	PRO	4.4
1	E	146[A]	SER	4.3
2	I	57	GLY	4.0
2	I	26[A]	LYS	3.9
1	E	77	GLU	3.5
1	E	78	GLY	3.5
1	E	145[A]	LYS	3.4
2	I	19[A]	ILE	3.2
1	E	76	VAL	3.1
1	E	117[A]	ARG	2.8
2	I	58	ALA	2.7
1	E	149	THR	2.5
1	E	148[A]	GLY	2.5
2	I	29[A]	LEU	2.5
1	E	127[A]	SER	2.4
1	E	220	LYS	2.3
2	I	52[A]	MET	2.3
1	E	135[A]	GLN	2.2
1	E	187	GLU	2.2
1	E	23	GLY	2.0
1	E	21	THR	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	I	48	ALA	2.0
2	I	18	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	7OZ	I	15	7/8	0.99	0.09	13,16,19,19	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	E	305	6/6	0.36	0.30	51,61,63,63	0
4	GOL	E	307[A]	6/6	0.50	0.29	32,39,41,42	14
4	GOL	E	307[B]	6/6	0.50	0.29	31,37,39,40	14
4	GOL	E	304	6/6	0.59	0.38	38,46,52,54	0
3	SO4	E	315[A]	5/5	0.72	0.29	44,44,44,45	5
4	GOL	E	302[A]	6/6	0.75	0.18	34,41,45,45	14
4	GOL	E	302[B]	6/6	0.75	0.18	35,43,46,47	14
3	SO4	E	308	5/5	0.77	0.21	35,37,38,39	5
3	SO4	E	301	5/5	0.78	0.36	19,20,22,23	5
4	GOL	E	306	6/6	0.78	0.28	42,51,55,56	0
4	GOL	I	101[B]	6/6	0.82	0.42	27,32,35,37	14
4	GOL	I	102	6/6	0.82	0.27	24,35,43,45	0
3	SO4	E	314	5/5	0.83	0.23	35,38,39,40	5
4	GOL	I	107[A]	6/6	0.83	0.23	23,28,33,34	14
3	SO4	E	313[A]	5/5	0.85	0.16	47,47,47,47	5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	E	311	5/5	0.87	0.21	48,49,49,49	5
4	GOL	E	303	6/6	0.91	0.20	35,42,46,47	0
3	SO4	I	103	5/5	0.92	0.15	38,39,40,41	0
3	SO4	I	105	5/5	0.93	0.11	21,21,26,27	5
3	SO4	E	316	5/5	0.94	0.26	41,42,44,44	0
3	SO4	E	309	5/5	0.97	0.26	51,51,51,51	0
3	SO4	E	310[A]	5/5	0.98	0.14	15,16,18,18	5
3	SO4	E	310[B]	5/5	0.98	0.14	55,55,55,55	5
3	SO4	I	104	5/5	0.98	0.07	19,20,23,24	0
3	SO4	I	106	5/5	0.99	0.08	17,18,18,20	0
5	CA	E	312	1/1	1.00	0.07	23,23,23,23	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.