

Full wwPDB X-ray Structure Validation Report (i)

Oct 3, 2023 – 03:02 AM EDT

PDB ID	:	6PI5
Title	:	The evolving story of AtzT, a periplasmic binding protein
Authors	:	Peat, T.S.; Newman, J.; Scott, C.; Esquirol, L.; Dennis, M.; Nebl, T.
Deposited on		
Resolution	:	1.67 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	FAILED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.67 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 11222 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	224	Total	С	Ν	0	\mathbf{S}	0	8	0
	A	334	2601	1663	442	485	11	0		
1	В	334	Total	С	Ν	0	S	0	3	0
	. D	- 334	2556	1635	434	476	11	0		
1	С	<u> </u>	Total	С	Ν	0	S	0	3	0
	C	333	2545	1626	431	477	11	0	J	0
1	П	220	Total	С	Ν	0	S	0	5	0
		332	2555	1631	434	478	12	0	0	0

• Molecule 1 is a protein called Atrazine periplasmic binding protein.

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	6	MET	-	initiating methionine	UNP Q936X6
А	7	GLY	-	expression tag	UNP Q936X6
А	8	SER	-	expression tag	UNP Q936X6
А	9	SER	-	expression tag	UNP Q936X6
А	10	HIS	-	expression tag	UNP Q936X6
А	11	HIS	-	expression tag	UNP Q936X6
А	12	HIS	-	expression tag	UNP Q936X6
А	13	HIS	-	expression tag	UNP Q936X6
А	14	HIS	-	expression tag	UNP Q936X6
А	15	HIS	-	expression tag	UNP Q936X6
А	16	SER	-	expression tag	UNP Q936X6
А	17	SER	-	expression tag	UNP Q936X6
А	18	GLY	-	expression tag	UNP Q936X6
А	19	LEU	-	expression tag	UNP Q936X6
А	20	VAL	-	expression tag	UNP Q936X6
А	21	PRO	-	expression tag	UNP Q936X6
А	22	ARG	-	expression tag	UNP Q936X6
А	23	GLY	-	expression tag	UNP Q936X6
А	24	SER	-	expression tag	UNP Q936X6
А	25	HIS	-	expression tag	UNP Q936X6
А	26	MET	-	expression tag	UNP Q936X6

Continued on next page...



Chain	Residue	Modelled	Actual	Comment	Reference
В	6	MET	-	initiating methionine	UNP Q936X6
В	7	GLY	-	expression tag	UNP Q936X6
В	8	SER	-	expression tag	UNP Q936X6
В	9	SER	-	expression tag	UNP Q936X6
В	10	HIS	-	expression tag	UNP Q936X6
В	11	HIS	-	expression tag	UNP Q936X6
В	12	HIS	-	expression tag	UNP Q936X6
В	13	HIS	-	expression tag	UNP Q936X6
В	14	HIS	-	expression tag	UNP Q936X6
В	15	HIS	-	expression tag	UNP Q936X6
В	16	SER	-	expression tag	UNP Q936X6
В	17	SER	-	expression tag	UNP Q936X6
В	18	GLY	-	expression tag	UNP Q936X6
В	19	LEU	-	expression tag	UNP Q936X6
В	20	VAL	-	expression tag	UNP Q936X6
В	21	PRO	-	expression tag	UNP Q936X6
В	22	ARG	-	expression tag	UNP Q936X6
В	23	GLY	-	expression tag	UNP Q936X6
В	24	SER	-	expression tag	UNP Q936X6
В	25	HIS	-	expression tag	UNP Q936X6
В	26	MET	-	expression tag	UNP Q936X6
С	6	MET	_	initiating methionine	UNP Q936X6
С	7	GLY	-	expression tag	UNP Q936X6
С	8	SER	-	expression tag	UNP Q936X6
С	9	SER	-	expression tag	UNP Q936X6
С	10	HIS	-	expression tag	UNP Q936X6
С	11	HIS	-	expression tag	UNP Q936X6
С	12	HIS	-	expression tag	UNP Q936X6
С	13	HIS	-	expression tag	UNP Q936X6
С	14	HIS	-	expression tag	UNP Q936X6
С	15	HIS	-	expression tag	UNP Q936X6
С	16	SER	-	expression tag	UNP Q936X6
С	17	SER	-	expression tag	UNP Q936X6
С	18	GLY	-	expression tag	UNP Q936X6
С	19	LEU	-	expression tag	UNP Q936X6
С	20	VAL	-	expression tag	UNP Q936X6
С	21	PRO	-	expression tag	UNP Q936X6
С	22	ARG	-	expression tag	UNP Q936X6
С	23	GLY	-	expression tag	UNP Q936X6
С	24	SER	-	expression tag	UNP Q936X6
С	25	HIS	-	expression tag	UNP Q936X6
С	26	MET	-	expression tag	UNP Q936X6

Continued from previous page...


Continued on next page...



Chain	Residue	Modelled	Actual	Comment	Reference
D	6	MET	-	initiating methionine	UNP Q936X6
D	7	GLY	-	expression tag	UNP Q936X6
D	8	SER	-	expression tag	UNP Q936X6
D	9	SER	-	expression tag	UNP Q936X6
D	10	HIS	-	expression tag	UNP Q936X6
D	11	HIS	-	expression tag	UNP Q936X6
D	12	HIS	-	expression tag	UNP Q936X6
D	13	HIS	-	expression tag	UNP Q936X6
D	14	HIS	-	expression tag	UNP Q936X6
D	15	HIS	-	expression tag	UNP Q936X6
D	16	SER	-	expression tag	UNP Q936X6
D	17	SER	-	expression tag	UNP Q936X6
D	18	GLY	-	expression tag	UNP Q936X6
D	19	LEU	-	expression tag	UNP Q936X6
D	20	VAL	-	expression tag	UNP Q936X6
D	21	PRO	-	expression tag	UNP Q936X6
D	22	ARG	-	expression tag	UNP Q936X6
D	23	GLY	-	expression tag	UNP Q936X6
D	24	SER	-	expression tag	UNP Q936X6
D	25	HIS	-	expression tag	UNP Q936X6
D	26	MET	-	expression tag	UNP Q936X6

Continued from previous page...


• Molecule 2 is GUANINE (three-letter code: GUN) (formula: $C_5H_5N_5O$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	$\begin{array}{cccc} \text{Total} \text{C} \text{N} \text{O} \\ 11 5 5 1 \end{array}$	0	0
2	В	1	Total C N O 11 5 5 1	0	0
2	С	1	Total C N O 11 5 5 1	0	0
2	D	1	Total C N O 11 5 5 1	0	0

• Molecule 3 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C₂H₆OS).



Μ	ol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf
e J	}	С	1	Total 4	$\begin{array}{c} \mathrm{C} \\ \mathrm{2} \end{array}$	0 1	S 1	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	254	Total O 254 254	0	0
4	В	264	Total O 264 264	0	0
4	С	233	Total O 233 233	0	0
4	D	166	Total O 166 166	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	99.01Å 63.14Å 120.37Å	Depositor
a, b, c, α , β , γ	90.00° 93.55° 90.00°	Depositor
Resolution (Å)	49.30 - 1.67	Depositor
% Data completeness	99.3 (49.30-1.67)	Depositor
(in resolution range)		-
R _{merge}	0.08	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.07 (at 1.67 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.156 , 0.184	Depositor
Wilson B-factor $(Å^2)$	21.6	Xtriage
Anisotropy	0.049	Xtriage
L-test for twinning ²	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	11222	wwPDB-VP
Average B, all atoms $(Å^2)$	26.0	wwPDB-VP

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



6PI5

Mol	Type	Chain	Res	Link	В	ond leng	gths	Bond angles		
IVIOI					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
2	GUN	В	401	-	7,12,12	1.91	2 (28%)	8,17,17	1.37	2 (25%)
2	GUN	А	401	-	7,12,12	1.99	2 (28%)	8,17,17	1.31	1 (12%)
2	GUN	D	401	-	7,12,12	1.49	2 (28%)	8,17,17	1.46	2 (25%)
3	DMS	С	402	-	3,3,3	0.35	0	3,3,3	0.31	0
2	GUN	С	401	-	7,12,12	1.83	2 (28%)	8,17,17	1.52	1 (12%)

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GUN	В	401	-	-	-	0/2/2/2
2	GUN	С	401	-	-	-	0/2/2/2
2	GUN	D	401	-	-	-	0/2/2/2
2	GUN	А	401	-	-	-	0/2/2/2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms		Observed(Å)	Ideal(Å)
2	А	401	GUN	C5-C6	-4.07	1.39	1.47
2	В	401	GUN	C5-C6	-3.82	1.39	1.47
2	С	401	GUN	C5-C6	-3.45	1.40	1.47
2	В	401	GUN	C6-N1	2.86	1.42	1.37
2	С	401	GUN	C6-N1	2.77	1.42	1.37
2	А	401	GUN	C6-N1	2.65	1.41	1.37
2	D	401	GUN	C5-C6	-2.59	1.42	1.47
2	D	401	GUN	C6-N1	2.56	1.41	1.37

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	С	401	GUN	N2-C2-N3	-2.73	114.42	119.74
2	А	401	GUN	N2-C2-N3	-2.31	115.23	119.74
2	D	401	GUN	O6-C6-C5	2.14	128.55	124.37
2	В	401	GUN	C2-N1-C6	-2.11	121.20	125.10
2	В	401	GUN	N2-C2-N3	-2.07	115.71	119.74

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	D	401	GUN	N2-C2-N3	-2.02	115.80	119.74

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.























4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

