



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 25, 2020 – 05:28 PM BST

PDB ID : 4PJ2
Title : Crystal structure of *Aeromonas hydrophila* PliI in complex with Meretrix lusoria lysozyme
Authors : Leysen, S.; Van Herreweghe, J.M.; Yoneda, K.; Ogata, M.; Usui, T.; Michiels, C.W.; Araki, T.; Strelkov, S.V.
Deposited on : 2014-05-10
Resolution : 1.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2024 (1.28-1.20)
Clashscore	141614	1007 (1.26-1.22)
Ramachandran outliers	138981	2053 (1.28-1.20)
Sidechain outliers	138945	2051 (1.28-1.20)
RSRZ outliers	127900	1987 (1.28-1.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	135	 8% (poor fit), 83% (0-1 outliers), 6% (2 outliers), 10% (3+ outliers or not modelled)
1	B	135	 7% (poor fit), 83% (0-1 outliers), 7% (2 outliers), 9% (3+ outliers or not modelled)
2	C	122	 2% (poor fit), 92% (0-1 outliers), 8% (2+ outliers or not modelled)
2	D	122	 11% (poor fit), 89% (0-1 outliers), 11% (2+ outliers or not modelled)

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 8388 atoms, of which 3889 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative exported protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
1	A	121	1923	609	960	158	196	0	10	0
1	B	123	1946	617	982	157	190	0	7	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	19	MET	-	initiating methionine	UNP A0KHJ5
A	146	LEU	-	expression tag	UNP A0KHJ5
A	147	GLU	-	expression tag	UNP A0KHJ5
A	148	HIS	-	expression tag	UNP A0KHJ5
A	149	HIS	-	expression tag	UNP A0KHJ5
A	150	HIS	-	expression tag	UNP A0KHJ5
A	151	HIS	-	expression tag	UNP A0KHJ5
A	152	HIS	-	expression tag	UNP A0KHJ5
A	153	HIS	-	expression tag	UNP A0KHJ5
B	19	MET	-	initiating methionine	UNP A0KHJ5
B	146	LEU	-	expression tag	UNP A0KHJ5
B	147	GLU	-	expression tag	UNP A0KHJ5
B	148	HIS	-	expression tag	UNP A0KHJ5
B	149	HIS	-	expression tag	UNP A0KHJ5
B	150	HIS	-	expression tag	UNP A0KHJ5
B	151	HIS	-	expression tag	UNP A0KHJ5
B	152	HIS	-	expression tag	UNP A0KHJ5
B	153	HIS	-	expression tag	UNP A0KHJ5

- Molecule 2 is a protein called Lysozyme.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	C	122	1961	611	976	182	173	19	8	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	D	122	1923	600	955	177	172	19	10	0	

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	A	1	14	3	8	3	0	0
3	A	1	14	3	8	3	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
4	B	1	1	1	0	0
4	A	1	1	1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	156	156	156	0	0
5	B	162	162	162	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	172	Total	O	0	0
			172	172		
5	D	115	Total	O	0	0
			115	115		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative exported protein





- Molecule 1: Putative exported protein



- Molecule 2: Lysozyme



- Molecule 2: Lysozyme



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	52.93Å 77.67Å 69.38Å 90.00° 108.12° 90.00°	Depositor
Resolution (Å)	47.80 – 1.24 47.80 – 1.24	Depositor EDS
% Data completeness (in resolution range)	95.5 (47.80-1.24) 95.5 (47.80-1.24)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.64 (at 1.24Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.148 , 0.168 0.152 , 0.172	Depositor DCC
R_{free} test set	1457 reflections (1.01%)	wwPDB-VP
Wilson B-factor (Å ²)	15.6	Xtriage
Anisotropy	0.448	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 48.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	8388	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/1007	0.67	0/1365
1	B	0.47	0/1002	0.66	0/1360
2	C	0.44	0/1036	0.61	0/1382
2	D	0.43	0/1016	0.58	0/1356
All	All	0.45	0/4061	0.63	0/5463

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	963	960	952	7	0
1	B	964	982	980	11	0
2	C	985	976	973	8	0
2	D	968	955	954	15	0
3	A	12	16	16	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	156	0	0	2	3
5	B	162	0	0	9	3

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	172	0	0	6	2
5	D	115	0	0	8	4
All	All	4499	3889	3875	39	6

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:118:CYS:SG	5:D:302:HOH:O	2.18	1.01
2:D:13:CYS:SG	5:D:301:HOH:O	2.20	0.98
2:D:16:LYS:O	5:D:201:HOH:O	1.98	0.81
2:C:85[A]:ARG:NH1	5:C:201:HOH:O	2.13	0.81
2:D:13:CYS:N	5:D:301:HOH:O	2.14	0.80
2:D:13:CYS:SG	5:D:302:HOH:O	2.39	0.80
1:B:22:GLY:N	5:B:458:HOH:O	2.14	0.78
1:B:91:LYS:O	5:B:301:HOH:O	2.07	0.72
1:A:26:GLN:OE1	5:A:400:HOH:O	2.07	0.72
2:C:50:ARG:NH1	5:C:337:HOH:O	2.22	0.71
2:C:30:MET:HB2	5:C:342:HOH:O	1.89	0.71
1:A:22:GLY:O	5:A:405:HOH:O	2.08	0.70
2:D:9:ARG:O	5:D:301:HOH:O	2.09	0.69
1:B:92:GLN:N	1:B:92:GLN:OE1	2.27	0.68
1:B:71[B]:LYS:NZ	5:B:302:HOH:O	2.28	0.64
2:C:29:ASP:HB3	5:C:207:HOH:O	2.03	0.59
1:A:132:PRO:O	2:D:30:MET:SD	2.60	0.59
1:B:122:ASP:OD1	5:B:302:HOH:O	2.17	0.59
2:D:82[B]:CYS:SG	2:D:88:GLY:HA3	2.50	0.51
2:C:29:ASP:N	5:C:207:HOH:O	2.43	0.51
1:A:35[B]:VAL:HG11	1:A:115:LEU:CD2	2.42	0.50
2:D:111:ARG:HA	2:D:114:LYS:HE3	1.97	0.47
1:B:133:ASN:C	5:B:444:HOH:O	2.53	0.46
2:C:26:CYS:HB3	2:C:33[B]:LEU:HG	1.97	0.46
1:B:114[B]:THR:HG22	5:B:307:HOH:O	2.16	0.46
1:B:81:GLU:OE1	5:B:368:HOH:O	2.21	0.45
2:C:82[B]:CYS:SG	2:C:88:GLY:HA3	2.57	0.44
1:A:115:LEU:HA	1:A:120:GLY:O	2.17	0.44
2:C:23:ASN:OD1	5:C:298:HOH:O	2.21	0.44
1:B:134:GLU:O	5:B:455:HOH:O	2.22	0.43
1:B:92:GLN:HG2	5:B:377:HOH:O	2.17	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:28:MET:HG2	2:D:33:LEU:CD2	2.48	0.43
1:A:35[B]:VAL:HG11	1:A:115:LEU:HD22	2.01	0.42
2:D:28:MET:HG2	2:D:33:LEU:HD23	2.02	0.41
2:D:13:CYS:HA	5:D:302:HOH:O	2.21	0.41
2:D:82[B]:CYS:SG	2:D:88:GLY:CA	3.09	0.41
1:B:33:GLN:OE1	1:B:117:PRO:HA	2.20	0.41
1:A:100[B]:GLU:CD	2:D:30:MET:HE3	2.41	0.40
2:D:20:GLY:N	5:D:201:HOH:O	2.54	0.40

All (6) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:338:HOH:O	5:D:239:HOH:O[2_456]	1.94	0.26
5:A:332:HOH:O	5:D:212:HOH:O[2_456]	1.95	0.25
5:A:303:HOH:O	5:D:223:HOH:O[1_455]	1.96	0.24
5:B:338:HOH:O	5:C:221:HOH:O[2_355]	2.12	0.08
5:A:303:HOH:O	5:D:240:HOH:O[1_455]	2.15	0.05
5:B:337:HOH:O	5:C:325:HOH:O[1_655]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	128/135 (95%)	127 (99%)	1 (1%)	0	100	100
1	B	126/135 (93%)	124 (98%)	2 (2%)	0	100	100
2	C	131/122 (107%)	130 (99%)	1 (1%)	0	100	100
2	D	130/122 (107%)	129 (99%)	1 (1%)	0	100	100
All	All	515/514 (100%)	510 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	111/114 (97%)	107 (96%)	4 (4%)	35	4
1	B	110/114 (96%)	109 (99%)	1 (1%)	78	49
2	C	108/97 (111%)	107 (99%)	1 (1%)	78	49
2	D	107/97 (110%)	106 (99%)	1 (1%)	78	49
All	All	436/422 (103%)	429 (98%)	7 (2%)	65	26

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	LEU
1	A	115	LEU
1	A	123[A]	SER
1	A	123[B]	SER
1	B	134	GLU
2	C	21	CYS
2	D	21	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	202	-	5,5,5	0.29	0	5,5,5	0.45	0
3	GOL	A	201	-	5,5,5	0.40	0	5,5,5	0.98	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	202	-	-	2/4/4/4	-
3	GOL	A	201	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	202	GOL	O1-C1-C2-C3
3	A	202	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	121/135 (89%)	0.65	11 (9%) 9 7	11, 18, 36, 42	0
1	B	123/135 (91%)	0.76	9 (7%) 15 12	11, 16, 38, 46	0
2	C	122/122 (100%)	0.24	2 (1%) 72 67	11, 17, 24, 31	0
2	D	122/122 (100%)	0.73	13 (10%) 6 5	14, 25, 38, 44	0
All	All	488/514 (94%)	0.59	35 (7%) 15 12	11, 19, 37, 46	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	146	LEU	8.9
1	B	133	ASN	6.9
1	A	124	PHE	6.3
1	B	143	PRO	5.7
2	D	30	MET	5.5
1	B	145	ASP	5.1
1	B	144	ARG	5.0
2	D	120	GLY	4.6
1	B	92	GLN	4.4
1	A	118	GLN	4.1
1	B	91	LYS	3.9
1	A	142[A]	THR	3.8
2	D	31	GLY	3.8
2	C	30	MET	3.3
1	A	92	GLN	3.3
2	D	116	SER	3.2
2	D	27	LYS	3.2
2	D	106	ILE	3.2
2	D	9	ARG	3.2
1	A	89	GLY	3.0
2	D	24	VAL	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	90	ASP	2.6
1	A	91	LYS	2.6
1	A	122	ASP	2.6
1	A	123[A]	SER	2.5
1	A	119	GLU	2.4
2	D	21	CYS	2.3
2	D	28	MET	2.3
1	A	85	LEU	2.2
1	A	121	LEU	2.2
2	D	119	HIS	2.2
2	D	32	SER	2.2
2	D	60	ALA	2.1
1	B	142	THR	2.1
2	C	31	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	202	6/6	0.86	0.18	33,40,41,44	0
3	GOL	A	201	6/6	0.90	0.12	18,21,24,26	0
4	MG	A	203	1/1	0.98	0.06	28,28,28,28	0
4	MG	B	201	1/1	0.99	0.07	25,25,25,25	1

6.5 Other polymers [i](#)

There are no such residues in this entry.