



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 16, 2023 – 05:10 PM EST

PDB ID : 4PJH
Title : Structure of human MR1-Ac-6-FP in complex with human MAIT B-G8 TCR
Authors : Birkinshaw, R.W.; Rossjohn, J.
Deposited on : 2014-05-12
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	 2% 92% 5%
1	C	271	 4% 89% 6% 6%
2	B	100	 95%
2	D	100	 4% 95%
3	E	205	 10% 89% 5% 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	205	 2% 93% • •
4	F	244	 2% 93% 5% • •
4	H	244	 % 95% • •

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 14126 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major histocompatibility complex class I-related gene protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	262	Total	C	N	O	S	0	2	0
			2175	1394	377	394	10			
1	C	256	Total	C	N	O	S	0	1	0
			2084	1331	363	379	11			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP Q95460
A	261	SER	CYS	engineered mutation	UNP Q95460
C	0	MET	-	initiating methionine	UNP Q95460
C	261	SER	CYS	engineered mutation	UNP Q95460

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	99	Total	C	N	O	S	0	0	0
			794	508	133	150	3			
2	D	98	Total	C	N	O	S	0	0	0
			779	501	132	143	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769
D	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called TCR-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	192	Total	C	N	O	S	0	0	0
			1456	925	231	290	10			

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	G	198	Total 1534	C 968	N 243	O 313	S 10	0	1	0

- Molecule 4 is a protein called TCR-beta.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	F	239	Total 1852	C 1164	N 322	O 357	S 9	0	0	0
4	H	239	Total 1851	C 1164	N 324	O 354	S 9	0	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	A	1	Total 6	C 3	O 3	0	0
5	A	1	Total 6	C 3	O 3	0	0
5	C	1	Total 6	C 3	O 3	0	0
5	C	1	Total 6	C 3	O 3	0	0

- Molecule 6 is N-(6-formyl-4-oxo-3,4-dihydropteridin-2-yl)acetamide (three-letter code: 30W) (formula: C₉H₇N₅O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			16	9	5	2		
6	C	1	Total	C	N	O	0	0
			16	9	5	2		

- Molecule 7 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	1	Total	Na	0	0
			1	1		
7	E	1	Total	Na	0	0
			1	1		
7	F	1	Total	Na	0	0
			1	1		
7	H	1	Total	Na	0	0
			1	1		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	304	Total	O	0	0
			304	304		
8	B	122	Total	O	0	0
			122	122		
8	C	230	Total	O	0	0
			230	230		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	D	68	Total O 68 68	0	0
8	E	154	Total O 154 154	0	0
8	F	241	Total O 241 241	0	0
8	G	191	Total O 191 191	0	0
8	H	231	Total O 231 231	0	0



3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Major histocompatibility complex class I-related gene protein




- Molecule 1: Major histocompatibility complex class I-related gene protein





- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin



- Molecule 3: TCR-alpha




GLU
SER
SER

- Molecule 3: TCR-alpha



HIS MET MET GLY Q2 Q28 V80 L51 R66 S83 Q112 L121 Q145 S149 C157 D179 S199 PRO GLU SER SER

- Molecule 4: TCR-beta



HIS MET ASN ALA G3 N65 R77 L144 Y149 F150 D151 H152 P174 R185 R191 P202 Y213 N218 Q223 D224 A241 ASP

- Molecule 4: TCR-beta



HIS MET ASN ALA G3 V125 P128 L141 A180 L181 R191 T197 Q209 I232 A235 A241 ASP

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	217.74Å 71.48Å 143.80Å 90.00° 104.59° 90.00°	Depositor
Resolution (Å)	29.28 – 2.00 28.76 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (29.28-2.00) 99.9 (28.76-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.75 (at 2.00Å)	Xtrriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.169 , 0.198 0.171 , 0.201	Depositor DCC
R_{free} test set	7250 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	26.4	Xtrriage
Anisotropy	0.673	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 58.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	14126	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 30W, NA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/2240	0.61	0/3043
1	C	0.52	0/2142	0.60	0/2907
2	B	0.51	0/817	0.60	0/1113
2	D	0.44	0/802	0.62	0/1093
3	E	0.50	0/1487	0.64	0/2021
3	G	0.50	0/1568	0.63	0/2129
4	F	0.52	0/1901	0.65	0/2590
4	H	0.50	0/1900	0.64	0/2588
All	All	0.51	0/12857	0.62	0/17484

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2175	0	2060	8	0
1	C	2084	0	1943	11	0
2	B	794	0	732	2	0
2	D	779	0	722	2	0
3	E	1456	0	1337	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	G	1534	0	1422	3	0
4	F	1852	0	1725	5	0
4	H	1851	0	1729	4	0
5	A	12	0	16	1	0
5	C	12	0	16	0	0
6	A	16	0	6	0	0
6	C	16	0	6	0	0
7	B	1	0	0	0	0
7	E	1	0	0	0	0
7	F	1	0	0	0	0
7	H	1	0	0	0	0
8	A	304	0	0	2	0
8	B	122	0	0	2	0
8	C	230	0	0	0	0
8	D	68	0	0	1	0
8	E	154	0	0	1	0
8	F	241	0	0	0	0
8	G	191	0	0	0	0
8	H	231	0	0	1	0
All	All	14126	0	11714	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:G:28:GLY:HA3	3:G:93[A]:SER:OG	1.81	0.81
1:C:154:LYS:HD3	3:G:51:LEU:HD11	1.74	0.70
3:E:95:TYR:OH	8:E:554:HOH:O	2.07	0.69
1:A:221:ILE:HD13	1:A:223:GLN:O	1.93	0.68
1:C:254:TYR:CB	1:C:254:TYR:C	2.63	0.67
4:H:125:VAL:HG23	4:H:235:ALA:HB3	1.79	0.65
1:C:254:TYR:CB	1:C:254:TYR:N	2.60	0.65
5:A:302:GOL:H12	8:A:534:HOH:O	1.97	0.64
1:C:254:TYR:C	1:C:254:TYR:N	2.52	0.64
4:F:65:ASN:HD21	4:F:77:ARG:HE	1.46	0.63
2:B:3:ARG:HD2	8:B:229:HOH:O	2.03	0.58
1:C:77:LEU:HD13	1:C:92:TYR:HB2	1.85	0.57
1:C:215:MET:HG3	1:C:257:HIS:CD2	2.40	0.56
4:H:128:PRO:HD3	4:H:141:LEU:HG	1.88	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:186:VAL:HG11	1:A:269:VAL:HG22	1.88	0.54
3:E:28:GLY:HA3	3:E:93:SER:HB3	1.90	0.53
4:H:197:THR:HG23	8:H:431:HOH:O	2.09	0.52
1:A:154:LYS:HD3	3:E:51:LEU:HD11	1.93	0.51
1:C:186:VAL:HG11	1:C:269:VAL:HG22	1.93	0.49
1:A:1:ARG:HD3	8:A:698:HOH:O	2.13	0.49
3:E:131:VAL:CG1	4:F:144:LEU:HD11	2.44	0.47
3:E:50:VAL:O	3:E:66:ARG:HD3	2.15	0.47
1:C:113:ALA:HB2	2:D:60:TRP:CE2	2.50	0.47
4:F:151:ASP:CG	4:F:174:PRO:HG3	2.37	0.45
4:F:152:HIS:HB3	4:F:213:TYR:HB2	1.97	0.45
4:H:209:GLN:HG3	4:H:232:ILE:HG23	1.99	0.44
3:G:50:VAL:O	3:G:66:ARG:HD3	2.17	0.44
1:A:113:ALA:HB2	2:B:60:TRP:CE2	2.53	0.44
1:A:77:LEU:HD13	1:A:92:TYR:HB2	2.01	0.42
1:A:4:SER:HB3	1:A:99:GLU:HG2	2.02	0.42
1:A:203:HIS:HD2	8:B:255:HOH:O	2.03	0.41
1:C:19:VAL:HG11	8:D:165:HOH:O	2.18	0.41
1:C:4:SER:HB3	1:C:99:GLU:HG2	2.03	0.41
3:E:121:LEU:O	3:E:130:SER:HB2	2.21	0.41
1:C:116:GLY:O	2:D:3:ARG:NH2	2.54	0.41
4:F:149:TYR:HB2	4:F:185:ARG:HG2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	258/271 (95%)	253 (98%)	5 (2%)	0	100 100
1	C	248/271 (92%)	242 (98%)	5 (2%)	1 (0%)	34 30
2	B	97/100 (97%)	97 (100%)	0	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	D	96/100 (96%)	96 (100%)	0	0	100	100
3	E	186/205 (91%)	183 (98%)	3 (2%)	0	100	100
3	G	197/205 (96%)	195 (99%)	2 (1%)	0	100	100
4	F	237/244 (97%)	234 (99%)	3 (1%)	0	100	100
4	H	237/244 (97%)	233 (98%)	4 (2%)	0	100	100
All	All	1556/1640 (95%)	1533 (98%)	22 (1%)	1 (0%)	51	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	17	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/241 (95%)	229 (100%)	1 (0%)	91	93
1	C	215/241 (89%)	212 (99%)	3 (1%)	67	72
2	B	86/95 (90%)	84 (98%)	2 (2%)	50	53
2	D	83/95 (87%)	82 (99%)	1 (1%)	71	76
3	E	154/181 (85%)	153 (99%)	1 (1%)	86	90
3	G	169/181 (93%)	167 (99%)	2 (1%)	71	76
4	F	197/210 (94%)	195 (99%)	2 (1%)	76	81
4	H	196/210 (93%)	194 (99%)	2 (1%)	76	81
All	All	1330/1454 (92%)	1316 (99%)	14 (1%)	73	78

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	73	PHE
2	B	19	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	70	PHE
1	C	73	PHE
1	C	82	ARG
1	C	215	MET
2	D	70	PHE
3	E	157	CYS
4	F	191	ARG
4	F	223	GLN
3	G	121	LEU
3	G	157	CYS
4	H	128	PRO
4	H	191	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	203	HIS
1	C	42	GLN
1	C	83	HIS
1	C	111	GLN
1	C	203	HIS
4	F	11	GLN
4	F	65	ASN
3	G	94	ASN
4	H	30	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	C	302	-	5,5,5	0.13	0	5,5,5	0.43	0
5	GOL	A	301	-	5,5,5	0.16	0	5,5,5	0.29	0
6	30W	A	303	1	16,17,18	0.66	1 (6%)	18,24,25	2.42	2 (11%)
6	30W	C	303	1	16,17,18	0.74	1 (6%)	18,24,25	2.44	3 (16%)
5	GOL	C	301	-	5,5,5	0.17	0	5,5,5	0.17	0
5	GOL	A	302	-	5,5,5	0.16	0	5,5,5	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	C	302	-	-	2/4/4/4	-
5	GOL	A	301	-	-	0/4/4/4	-
6	30W	A	303	1	-	0/4/4/6	0/2/2/2
6	30W	C	303	1	-	0/4/4/6	0/2/2/2
5	GOL	C	301	-	-	3/4/4/4	-
5	GOL	A	302	-	-	2/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	303	30W	C2-N2	-2.30	1.32	1.37
6	A	303	30W	C2-N2	-2.26	1.33	1.37

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	303	30W	N2-C2-N3	8.36	130.74	117.98

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	303	30W	N2-C2-N3	8.26	130.59	117.98
6	C	303	30W	N2-C2-N1	-5.13	110.76	117.72
6	A	303	30W	N2-C2-N1	-4.70	111.34	117.72
6	C	303	30W	C4A-C8A-N1	-2.15	120.11	123.18

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	302	GOL	O1-C1-C2-C3
5	C	301	GOL	C1-C2-C3-O3
5	C	302	GOL	O1-C1-C2-O2
5	A	302	GOL	O1-C1-C2-O2
5	C	301	GOL	O2-C2-C3-O3
5	A	302	GOL	O1-C1-C2-C3
5	C	301	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	302	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	262/271 (96%)	-0.41	5 (1%) 66 65	17, 26, 54, 98	0
1	C	256/271 (94%)	-0.22	10 (3%) 39 38	22, 32, 61, 101	0
2	B	99/100 (99%)	-0.37	0 100 100	20, 34, 58, 68	0
2	D	98/100 (98%)	-0.01	4 (4%) 37 36	27, 50, 78, 94	0
3	E	192/205 (93%)	0.17	21 (10%) 5 5	19, 37, 82, 108	0
3	G	198/205 (96%)	-0.35	4 (2%) 65 63	22, 31, 60, 77	0
4	F	239/244 (97%)	-0.33	4 (1%) 70 68	18, 31, 69, 217	0
4	H	239/244 (97%)	-0.37	3 (1%) 77 76	21, 34, 54, 83	0
All	All	1583/1640 (96%)	-0.26	51 (3%) 47 46	17, 32, 69, 217	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	H	180	ALA	5.8
1	A	222	VAL	5.1
1	A	17	HIS	5.1
3	E	190	ILE	4.8
1	C	221	ILE	4.7
3	E	180	PHE	4.5
4	H	181	LEU	4.5
4	F	241	ALA	4.3
3	E	149	SER	4.2
3	E	147	LYS	3.9
1	C	18	GLY	3.9
1	C	196	THR	3.7
4	F	202	PRO	3.6
3	E	198	PRO	3.3
3	E	146	SER	3.3
3	E	129	LYS	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	G	179	ASP	3.1
3	E	148	ASP	3.0
4	H	241	ALA	3.0
3	G	145	GLN	3.0
1	A	221	ILE	2.8
1	C	17	HIS	2.8
4	F	224	ASP	2.8
3	E	184	ASN	2.8
1	C	190	GLU	2.8
1	C	96	ILE	2.7
3	E	194	ASP	2.7
1	C	223	GLN	2.7
1	C	254	TYR	2.6
4	F	218	ASN	2.6
3	E	188	ASN	2.5
1	A	217	ASN	2.5
3	E	163	SER	2.5
3	E	112	GLN	2.4
3	E	113	ASN	2.4
3	E	199	SER	2.3
3	E	151	VAL	2.3
3	E	181	ALA	2.3
3	E	124	SER	2.3
3	E	189	SER	2.3
3	G	112	GLN	2.3
2	D	44	GLU	2.2
3	G	149	SER	2.2
1	C	188	ARG	2.2
3	E	182	CYS	2.1
2	D	0	MET	2.1
1	A	194	GLY	2.1
1	C	16	ILE	2.1
3	E	191	ILE	2.0
2	D	75	LYS	2.0
2	D	97	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	A	302	6/6	0.87	0.14	40,48,50,53	0
5	GOL	A	301	6/6	0.90	0.21	34,45,47,51	0
5	GOL	C	301	6/6	0.91	0.16	40,43,44,45	0
5	GOL	C	302	6/6	0.91	0.26	43,44,50,52	0
7	NA	E	301	1/1	0.95	0.23	39,39,39,39	0
7	NA	H	301	1/1	0.96	0.07	38,38,38,38	0
6	30W	C	303	16/17	0.97	0.20	26,28,29,31	0
6	30W	A	303	16/17	0.98	0.10	20,22,26,26	0
7	NA	F	301	1/1	0.98	0.14	27,27,27,27	0
7	NA	B	101	1/1	0.98	0.18	40,40,40,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.