

wwPDB EM Validation Summary Report (i)

Sep 21, 2024 – 10:59 am BST

PDB ID : 8PJ2

EMDB ID : EMD-17697

Title: Structure of human 48S translation initiation complex in AUG recognition

state after eIF5-induced GTP hydrolysis by eIF2 (48S-2)

Authors: Petrychenko, V.; Yi, S.-H.; Liedtke, D.; Peng, B.Z.; Rodnina, M.V.; Fischer,

N.

Deposited on : 2023-06-22

Resolution : 3.40 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev112

Mogul : 1.8.4, CSD as541be (2020)

MolProbity : FAILED

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

 $MapQ \quad : \quad 1.9.13$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.38.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.40 Å.

There are no overall percentile quality scores available for this entry.

MolProbity failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 55 unique types of molecules in this entry. The entry contains 118707 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Eukaryotic translation initiation factor 3 subunit B.

Mol	Chain	Residues		Ato	AltConf	Trace			
1	1	588	Total 3258	C 1986	N 633	O 634	S 5	0	0

• Molecule 2 is a protein called Eukaryotic translation initiation factor 3 subunit I.

Mol	Chain	Residues		Ato	ms	AltConf	Trace	
2	2	304	Total 1493	C 885	N 304	O 304	0	0

• Molecule 3 is a protein called Eukaryotic translation initiation factor 3 subunit K.

Mol	Chain	Residues		Ato	ms	AltConf	Trace	
3	3	213	Total 1057		N 213	O 213	0	0

• Molecule 4 is a protein called Eukaryotic translation initiation factor 3 subunit F.

Mol	Chain	Residues		Ato	ms	AltConf	Trace	
4	4	257	Total 1272	C 757	N 257	O 258	0	0

• Molecule 5 is a protein called Eukaryotic translation initiation factor 3 subunit L.

Mol	Chain	Residues		At	AltConf	Trace			
5	5	520	Total 4347	C 2814	N 721	O 793	S 19	0	0

• Molecule 6 is a protein called Eukaryotic translation initiation factor 3 subunit M.

Mol	Chain	Residues		Ato	AltConf	Trace			
6	6	362	Total 2196	C 1348	N 414	O 427	S 7	0	0



• Molecule 7 is a RNA chain called mRNA.

Mol	Chain	Residues		A	toms	AltConf	Trace		
7	7	57	Total 1218	C 547	N 231	O 383	P 57	0	0

• Molecule 8 is a protein called Eukaryotic translation initiation factor 3 subunit H.

Mol	Chain	Residues		Ato	ms	AltConf	Trace	
8	8	317	Total 1574	C 937	N 318	O 319	0	0

• Molecule 9 is a protein called 60S ribosomal protein L41.

\mathbf{Mol}	Chain	Residues		Ato	$\mathbf{m}\mathbf{s}$	AltConf	Trace		
0	0	24	Total	С	N	О	S	0	0
9	9	24	230	139	62	26	3	0	U

• Molecule 10 is a RNA chain called 18S rRNA.

Mol	Chain	Residues		I	AltConf	Trace			
10	A	1754	Total 37429	C 16718	N 6714	O 12244	P 1753	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Α	1248	B8N	U	$\operatorname{conflict}$	GB NR_046235.3

• Molecule 11 is a protein called 40S ribosomal protein S11.

Mol	Chain	Residues		At	oms	AltConf	Trace		
11	В	142	Total 1166	C 743	N 218	O 199	S 6	0	0

• Molecule 12 is a protein called 40S ribosomal protein S4, X isoform.

Mol	Chain	Residues		Ato	AltConf	Trace			
12	С	256	Total 2035	C 1302	N 378	O 347	S 8	0	0

• Molecule 13 is a protein called 40S ribosomal protein S9.



Mol	Chain	Residues		At	oms	AltConf	Trace		
13	D	177	Total 1477	C 941	N 295	O 239	S 2	0	0

• Molecule 14 is a protein called 40S ribosomal protein S23.

Mol	Chain	Residues		At	oms	AltConf	Trace		
1/1	E	140	Total	С	N	О	S	0	0
14	ப	140	1087	687	215	182	3		

• Molecule 15 is a protein called 40S ribosomal protein S30.

Mol	Chain	Residues		Ato	oms	AltConf	Trace		
15	F	59	Total 468	C 290	N 102	O 75	S 1	0	0

• Molecule 16 is a protein called 40S ribosomal protein S7.

Mol	Chain	Residues		At	oms	AltConf	Trace		
16	G	177	Total	C	N	0	S	0	0
			1430	917	260	252	1		

• Molecule 17 is a protein called 40S ribosomal protein S27.

Mol	Chain	Residues		At	oms	AltConf	Trace		
17	Н	81	Total 631	_	N 116	O 111	S 7	0	0

• Molecule 18 is a protein called 40S ribosomal protein S13.

Mol	Chain	Residues		At	oms	AltConf	Trace		
10	Т	150	Total	С	N	О	S	0	0
10	1	150	1208	773	229	205	1	0	U

• Molecule 19 is a protein called 40S ribosomal protein S15a.

Mol	Chain	Residues		At	oms	AltConf	Trace		
19	J	129	Total 1034	C 659	N 193	O 176	S 6	0	0

• Molecule 20 is a protein called 40S ribosomal protein S21.



Mol	Chain	Residues		At	oms			AltConf	Trace
20	I/	01	Total	С	N	О	S	0	0
20	IX.	01	617	380	114	118	5	0	U

• Molecule 21 is a protein called 40S ribosomal protein S2.

Mol	Chain	Residues		At	AltConf	Trace			
21	L	220	Total 1707	C 1104	N 292	O 301	S 10	0	0

• Molecule 22 is a protein called 40S ribosomal protein S17.

Mol	Chain	Residues		At	oms	AltConf	Trace		
22	M	131	Total	С	N	О	S	0	0
22	IVI	191	1064	668	198	194	4	0	U

• Molecule 23 is a protein called 40S ribosomal protein SA.

Mol	Chain	Residues		Ato	AltConf	Trace			
23	N	207	Total	С	N	О	S	0	0
	1,		1633	1040	288	297	8		

• Molecule 24 is a protein called 40S ribosomal protein S3a.

Mol	Chain	Residues		At	AltConf	Trace			
24	О	211	Total 1715	C 1088	N 307	O 306	S 14	0	0

• Molecule 25 is a protein called 40S ribosomal protein S14.

Mol	Chain	Residues		At	oms	AltConf	Trace		
25	Р	133	Total 997	C 610	N 196	O 185	S 6	0	0

• Molecule 26 is a protein called 40S ribosomal protein S26.

Mol	Chain	Residues		At	oms	AltConf	Trace		
26	Q	99	Total 792	C 492	N 165	O 130	S 5	0	0

• Molecule 27 is a protein called 40S ribosomal protein S8.



Mol	Chain	Residues		\mathbf{At}	oms		AltConf	Trace	
27	R	198	Total 1627	C 1021	N 322	O 279	S 5	0	0

• Molecule 28 is a protein called 40S ribosomal protein S6.

Mo	ol	Chain	Residues		Ato	AltConf	Trace			
28		S	230	Total 1862	C 1164	N 371	O 320	S 7	0	0

• Molecule 29 is a protein called 40S ribosomal protein S24.

Mol	Chain	Residues		At	oms		AltConf	Trace	
29	Т	125	Total 1015	C 642	N 199	O 169	S 5	0	0

• Molecule 30 is a protein called 40S ribosomal protein S5.

Mol	Chain	Residues		At	oms	AltConf	Trace		
30	V	184	Total 1461	C 914	N 276	O 264	S 7	0	0

• Molecule 31 is a protein called 40S ribosomal protein S16.

Mol	Chain	Residues		At	oms	AltConf	Trace		
31	Y	141	Total 1124	C 715	N 212	O 194	S	0	0

• Molecule 32 is a protein called 40S ribosomal protein S3.

Mol	Chain	Residues		Ato	AltConf	Trace			
32	Z	227	Total 1765	C 1125	N 317	O 315	S 8	0	0

• Molecule 33 is a protein called 40S ribosomal protein S10.

Mol	Chain	Residues		At	oms	AltConf	Trace		
33	a	99	Total 834	C 544	N 149	O 135	S 6	0	0

 \bullet Molecule 34 is a protein called 40S ribosomal protein S15.



Mol	Chain	Residues		At	oms	AltConf	Trace		
34	b	131	Total 1072	C 682	N 201	O 182	S 7	0	0

• Molecule 35 is a protein called Receptor of activated protein C kinase 1.

Mol	Chain	Residues		At	AltConf	Trace			
35	c	313	Total 2436	C 1535	N 424	O 465	S 12	0	0

• Molecule 36 is a protein called 40S ribosomal protein S19.

Mol	Chain	Residues		At	oms			AltConf	Trace
36	d	142	Total 1105	C 692	N 213	O 197	S 3	0	0

• Molecule 37 is a protein called 40S ribosomal protein S25.

Mol	Chain	Residues		At	oms	AltConf	Trace		
37	e	81	Total	С	N	О	S	0	0
	C	01	649	420	119	109	1		

• Molecule 38 is a protein called 40S ribosomal protein S18.

Mol	Chain	Residues		At	oms	AltConf	Trace		
38	f	149	Total 1227	C 770	N 249	O 207	S 1	0	0

• Molecule 39 is a protein called 40S ribosomal protein S20.

Mol	Chain	Residues		At	oms	AltConf	Trace		
39	h	103	Total	С	N	О	S	0	0
	11	100	817	511	155	147	4		

• Molecule 40 is a protein called 40S ribosomal protein S29.

Mol	Chain	Residues		Ato	$\mathbf{m}\mathbf{s}$	AltConf	Trace		
40	· .	50	Total	С	N	О	S	0	0
40	1	30	419	262	85	67	5	0	U

• Molecule 41 is a protein called Ubiquitin.



Mol	Chain	Residues		Ato	oms	AltConf	Trace		
41	k	68	Total 554	C 349	N 103	O 95	S	0	0

• Molecule 42 is a protein called 40S ribosomal protein S12.

Mol	Chain	Residues		At	oms	AltConf	Trace		
49	***	122	Total	С	N	О	S	0	0
42	m	122	950	596	168	177	9	U	U

• Molecule 43 is a protein called 40S ribosomal protein S28.

Mol	Chain	Residues		Ato	oms	Atoms					
43	n	63	Total 498	C 302	N 101	O 93	S 2	0	0		

• Molecule 44 is a protein called Eukaryotic translation initiation factor 3 subunit G.

Mol	Chain	Residues		Ato	ms		AltConf	Trace
44	О	77	Total 616	C 389	N 111	O 116	0	0

• Molecule 45 is a protein called Eukaryotic translation initiation factor 1A, X-chromosomal.

Mol	Chain	Residues		At	oms	AltConf	Trace		
45	q	112	Total 902	_	N 173	O 165	S 4	0	0

• Molecule 46 is a protein called Eukaryotic translation initiation factor 2 subunit 1.

Mol	Chain	Residues		Ato	AltConf	Trace			
46	r	296	Total 2138	C 1342	N 384	O 404	S 8	0	0

• Molecule 47 is a protein called Eukaryotic translation initiation factor 2 subunit 3.

Mol	Chain	Residues		At	AltConf	Trace			
47	t	455	Total 3439	C 2179	N 599	O 643	S 18	0	0

• Molecule 48 is a protein called Eukaryotic translation initiation factor 3 subunit A.



Mol	Chain	Residues		At	AltConf	Trace			
48	u	706	Total 5383	C 3379	N 982	O 999	S 23	1	0

• Molecule 49 is a protein called Eukaryotic translation initiation factor 3 subunit E.

Mol	Chain	Residues		At	oms			AltConf	Trace
49	V	405	Total 2740	C 1720	N 498	O 510	S 12	0	0

• Molecule 50 is a RNA chain called Initiator Met-tRNA-i.

Mol	Chain	Residues		A	toms			AltConf	Trace
50	***	75	Total	С	N	О	Р	0	0
30	50 w	75	1604	717	298	515	74	0	U

• Molecule 51 is a protein called Eukaryotic translation initiation factor 3 subunit D.

Mol	Chain	Residues		At	oms			AltConf	Trace
51	X	422	Total 2837	C 1749	N 522	O 556	S 10	0	0

• Molecule 52 is a protein called Eukaryotic translation initiation factor 3 subunit C.

Mol	Chain	Residues		At	oms			AltConf	Trace
52	у	656	Total 5263	C 3312	N 939	O 977	S 35	0	0

• Molecule 53 is a protein called Eukaryotic translation initiation factor 5.

Mol	Chain	Residues		\mathbf{A}	toms	AltConf	Trace		
53	Z	145	Total 1146	C 723	N 203	O 209	S 11	0	0

• Molecule 54 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	AltConf
54	A	83	Total Mg 83 83	0
54	V	1	Total Mg 1 1	0
54	f	1	Total Mg 1 1	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	AltConf
54	i	2	$\begin{array}{cc} \text{Total} & \text{Mg} \\ 2 & 2 \end{array}$	0

• Molecule 55 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	AltConf
55	Q	1	Total Zn 1 1	0
55	k	1	Total Zn 1 1	0

MolProbity failed to run properly - this section is therefore empty.



3 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	61742	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	45	Depositor
Minimum defocus (nm)	200	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	59000	Depositor
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	23.548	Depositor
Minimum map value	-8.712	Depositor
Average map value	0.000	Depositor
Map value standard deviation	1.000	Depositor
Recommended contour level	3	Depositor
Map size (Å)	417.74402, 417.74402, 417.74402	wwPDB
Map dimensions	432, 432, 432	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.96700007, 0.96700007, 0.96700007	Depositor



4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

29 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trmo	Chain	in Res	Link	Bond lengths				Bond angles			
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2		
10	A2M	A	484	10	18,25,26	4.23	9 (50%)	18,36,39	3.91	5 (27%)		
10	OMG	A	509	54,10	18,26,27	2.65	9 (50%)	19,38,41	2.55	9 (47%)		
10	OMC	A	517	10	19,22,23	2.95	8 (42%)	26,31,34	3.28	8 (30%)		



N.C. 1	TD.	aı ·	D	т. 1	В	ond leng	$_{ m gths}$	В	ond ang	gles
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
10	A2M	A	1031	10	18,25,26	4.37	7 (38%)	18,36,39	3.81	4 (22%)
10	A2M	A	27	54,10	18,25,26	4.27	8 (44%)	18,36,39	3.75	5 (27%)
10	PSU	A	119	10	18,21,22	1.00	1 (5%)	22,30,33	1.64	4 (18%)
10	OMC	A	174	54,10	19,22,23	3.20	8 (42%)	26,31,34	2.79	9 (34%)
10	A2M	A	166	10	18,25,26	4.27	8 (44%)	18,36,39	3.73	4 (22%)
10	JMH	A	1219	10	18,22,23	2.87	6 (33%)	21,32,35	1.62	4 (19%)
10	A2M	A	668	54,10	18,25,26	4.20	7 (38%)	18,36,39	3.84	5 (27%)
10	OMC	A	1703	10	19,22,23	3.23	8 (42%)	26,31,34	2.83	9 (34%)
10	PSU	A	823	10	18,21,22	1.12	1 (5%)	22,30,33	1.78	5 (22%)
10	MA6	A	1851	10	18,26,27	1.33	2 (11%)	19,38,41	3.27	2 (10%)
10	OMU	A	121	10	19,22,23	2.97	6 (31%)	26,31,34	1.73	5 (19%)
10	OMG	A	644	10	18,26,27	2.35	7 (38%)	19,38,41	2.58	9 (47%)
10	OMG	A	683	10	18,26,27	1.03	1 (5%)	19,38,41	1.27	2 (10%)
10	PSU	A	822	10	18,21,22	1.04	1 (5%)	22,30,33	1.84	5 (22%)
10	A2M	A	159	10	18,25,26	4.28	8 (44%)	18,36,39	3.81	4 (22%)
10	5MU	A	814	10	19,22,23	7.25	10 (52%)	28,32,35	3.91	19 (67%)
10	PSU	A	1081	10	18,21,22	1.00	1 (5%)	22,30,33	1.74	3 (13%)
10	B8N	A	1248	10	24,29,30	3.02	6 (25%)	29,42,45	2.29	7 (24%)
10	MA6	A	1850	10	18,26,27	1.37	2 (11%)	19,38,41	3.07	2 (10%)
10	OMU	A	116	10	19,22,23	2.98	6 (31%)	26,31,34	1.64	4 (15%)
10	PSU	A	1243	10	18,21,22	1.07	1 (5%)	22,30,33	1.81	4 (18%)
10	PSU	A	612	10	18,21,22	1.00	1 (5%)	22,30,33	1.82	5 (22%)
10	A2M	A	1678	54,10	18,25,26	4.34	8 (44%)	18,36,39	3.82	4 (22%)
10	6MZ	A	1832	54,10	18,25,26	1.75	4 (22%)	16,36,39	2.23	3 (18%)
10	UR3	A	1830	10	19,22,23	2.77	8 (42%)	26,32,35	1.45	4 (15%)
10	5MC	A	1374	10	18,22,23	0.62	0	26,32,35	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	A2M	A	484	10	-	1/5/27/28	0/3/3/3
10	OMG	A	509	54,10	-	3/5/27/28	0/3/3/3
10	OMC	A	517	10	-	3/9/27/28	0/2/2/2

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	A2M	A	1031	10	-	1/5/27/28	0/3/3/3
10	A2M	A	27	54,10	-	1/5/27/28	0/3/3/3
10	PSU	A	119	10	-	0/7/25/26	0/2/2/2
10	OMC	A	174	54,10	-	1/9/27/28	0/2/2/2
10	A2M	A	166	10	-	1/5/27/28	0/3/3/3
10	JMH	A	1219	10	-	2/7/25/26	0/2/2/2
10	A2M	A	668	54,10	-	2/5/27/28	0/3/3/3
10	OMC	A	1703	10	-	2/9/27/28	0/2/2/2
10	PSU	A	823	10	-	2/7/25/26	0/2/2/2
10	MA6	A	1851	10	-	3/7/29/30	0/3/3/3
10	OMU	A	121	10	-	0/9/27/28	0/2/2/2
10	OMG	A	644	10	-	4/5/27/28	0/3/3/3
10	OMG	A	683	10	-	2/5/27/28	0/3/3/3
10	PSU	A	822	10	-	2/7/25/26	0/2/2/2
10	A2M	A	159	10	-	3/5/27/28	0/3/3/3
10	5MU	A	814	10	-	0/7/25/26	0/2/2/2
10	PSU	A	1081	10	-	1/7/25/26	0/2/2/2
10	B8N	A	1248	10	-	5/16/34/35	0/2/2/2
10	MA6	A	1850	10	-	1/7/29/30	0/3/3/3
10	OMU	A	116	10	-	1/9/27/28	0/2/2/2
10	PSU	A	1243	10	-	0/7/25/26	0/2/2/2
10	PSU	A	612	10	-	0/7/25/26	0/2/2/2
10	A2M	A	1678	54,10	-	1/5/27/28	0/3/3/3
10	6MZ	A	1832	54,10	-	2/5/27/28	0/3/3/3
10	UR3	A	1830	10	-	2/7/25/26	0/2/2/2
10	5MC	A	1374	10	_	0/7/25/26	0/2/2/2

The worst 5 of 152 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$Ideal(\AA)$
10	A	814	5MU	C4-C5	20.88	1.79	1.44
10	A	814	5MU	C6-N1	15.31	1.64	1.38
10	A	1031	A2M	C3'-C2'	-13.11	1.23	1.52
10	A	1678	A2M	C3'-C2'	-12.88	1.24	1.52
10	A	27	A2M	C3'-C2'	-12.84	1.24	1.52

The worst 5 of 153 bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
10	A	1851	MA6	N1-C6-N6	-12.87	103.51	117.06
10	A	1850	MA6	N1-C6-N6	-12.03	104.40	117.06
10	A	484	A2M	C1'-N9-C4	10.54	145.17	126.64
10	A	814	5MU	C5-C4-N3	10.31	124.11	115.31
10	A	1678	A2M	C1'-N9-C4	10.23	144.61	126.64

There are no chirality outliers.

5 of 46 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
10	A	27	A2M	C1'-C2'-O2'-CM'
10	A	116	OMU	C1'-C2'-O2'-CM2
10	A	159	A2M	C1'-C2'-O2'-CM'
10	A	166	A2M	C1'-C2'-O2'-CM'
10	A	174	OMC	C1'-C2'-O2'-CM2

There are no ring outliers.

No monomer is involved in short contacts.

4.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

4.6 Ligand geometry (i)

Of 89 ligands modelled in this entry, 89 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers (i)

There are no such residues in this entry.



4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




5 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-17697. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

5.1 Orthogonal projections (i)

5.1.1 Primary map



5.1.2 Raw map





The images above show the map projected in three orthogonal directions.




5.2 Central slices (i)


5.2.1 Primary map










X Index: 216












Z Index: 180

The images above show central slices of the map in three orthogonal directions.




5.3 Largest variance slices (i)


5.3.1 Primary map








Y Index: 222




Z Index: 214


5.3.2 Raw map



X Index: 147



Y Index: 186




Z Index: 173

The images above show the largest variance slices of the map in three orthogonal directions.




5.4 Orthogonal standard-deviation projections (False-color) (i)

5.4.1 Primary map



5.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




5.5 Orthogonal surface views (i)

5.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 3.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

5.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.




5.6 Mask visualisation (i)


This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure

$5.6.1 \quad \mathrm{emd}_17697_\mathrm{msk}_1.\mathrm{map}\ \ \mathbf{\mathring{1}}$








6 Map analysis (i)

This section contains the results of statistical analysis of the map.


6.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



6.2 Volume estimate (i)




The volume at the recommended contour level is $1196~\mathrm{nm^3}$; this corresponds to an approximate mass of $1081~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



6.3 Rotationally averaged power spectrum (i)




^{*}Reported resolution corresponds to spatial frequency of 0.294 $\rm \mathring{A}^{-1}$



7 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

7.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.294 $\rm \mathring{A}^{-1}$



7.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)			
rtesolution estimate (A)	0.143	0.5	Half-bit	
Reported by author	3.40	-	-	
Author-provided FSC curve	3.53	4.70	3.67	
Unmasked-calculated*	4.22	8.15	4.37	


^{*}Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.22 differs from the reported value 3.4 by more than 10 %



8 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-17697 and PDB model 8PJ2. Per-residue inclusion information can be found in section ?? on page ??.


8.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 3.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




8.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


8.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (3).



8.4 Atom inclusion (i)



At the recommended contour level, 73% of all backbone atoms, 73% of all non-hydrogen atoms, are inside the map.



8.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (3) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7290	0.3200
1	0.0660	0.0560
2	0.0030	0.0270
3	0.2680	0.0420
4	0.4680	0.0680
5	0.2880	0.0140
6	0.4120	0.0450
7	0.4450	0.1710
8	0.4440	0.0950
9	0.8130	0.3960
A	0.9550	0.4200
В	0.8770	0.4630
С	0.9170	0.4800
D	0.8830	0.4620
E	0.9000	0.4850
F	0.7810	0.4060
G	0.7900	0.3880
Н	0.8420	0.4310
I	0.8670	0.4410
J	0.8900	0.4870
K	0.8230	0.4450
L	0.8350	0.4600
M	0.7810	0.3970
N	0.8590	0.4560
О	0.8670	0.4260
Р	0.8470	0.4250
Q	0.9070	0.4720
R	0.9160	0.4400
S	0.9250	0.4050
Т	0.9210	0.4370
V	0.8450	0.4470
Y	0.9070	0.4670
Z	0.8020	0.4310
a	0.8630	0.4190
b	0.8600	0.4060



Continued on next page...



$Continued\ from\ previous\ page...$

Chain	Atom inclusion	Q-score
c	0.8720	0.4070
d	0.9250	0.4570
e	0.7370	0.3520
f	0.8570	0.4140
h	0.8220	0.4070
i	0.9350	0.4950
k	0.7930	0.2220
m	0.7210	0.2450
n	0.7590	0.4100
О	0.4240	0.1700
q	0.7200	0.3750
r	0.5210	0.2030
t	0.0590	0.0420
u	0.5390	0.1820
v	0.5470	0.1060
W	0.8120	0.2070
X	0.5380	0.2200
У	0.5820	0.2390
Z	0.3610	0.2520

