

# Full wwPDB X-ray Structure Validation Report (i)

#### Oct 4, 2023 – 06:51 PM EDT

PDB ID : 6PK1

Title : Alanine-glyoxylate aminotransferase 1 (AGT1) from Arabidopsis thaliana in

presence of serine

Authors : Liepman, A.H.; Saper, M.A.

Deposited on : 2019-06-28

Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.orgA user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity : FAILED

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : FAILED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



## 2 Entry composition (i)


There are 4 unique types of molecules in this entry. The entry contains 13007 atoms, of which 6322 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Serine–glyoxylate aminotransferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	399	Total 6255	C 1993	H 3148	N 535	O 567	S 12	0	4	0
1	В	399	Total 6272	C 1999	H 3155	N 538	O 568	S 12	0	6	0


• Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C<sub>8</sub>H<sub>10</sub>NO<sub>6</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf		
2	Λ	1	Total	С	Н	N	О	Р	0	0	
2	$\mathcal{L}$ $\mathcal{A}$	1	24	8	8	1	6	1	0		
2	D	1	Total	С	Н	N	О	Р	0	0	
	2 B	1	24	8	8	1	6	1	0	U	

• Molecule 3 is 3-HYDROXYPYRUVIC ACID (three-letter code: 3PY) (formula: C<sub>3</sub>H<sub>4</sub>O<sub>4</sub>).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total 10	C 3	Н 3	O 4	0	0

#### • Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	223	Total O 223 223	0	0
4	В	199	Total O 199 199	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



## 3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants	139.18Å 62.38Å 96.98Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	22.16 - 2.10	Depositor
% Data completeness	90.6 (22.16-2.10)	Depositor
(in resolution range)	, , ,	
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	3.29  (at  2.10Å)	Xtriage
Refinement program	CNS 1.1, PHENIX 1.16rc1	Depositor
$R, R_{free}$	0.115 , 0.166	Depositor
Wilson B-factor (Å <sup>2</sup> )	19.6	Xtriage
Anisotropy	0.881	Xtriage
L-test for twinning <sup>2</sup>	$ < L > = 0.50, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	13007	wwPDB-VP
Average B, all atoms $(\mathring{A}^2)$	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.51% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $< L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 4 Model quality (i)

### 4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3 Torsion angles (i)

#### 4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

### 4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 4.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Chain Res Li		Bo	ond leng	$ ag{ths}$	Bond angles			
MIOI	Type	Chain	nes	Link	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2	
2	PLP	В	501	-	16,16,16	1.35	3 (18%)	20,23,23	1.54	3 (15%)	
2	PLP	A	501	-	16,16,16	1.30	3 (18%)	20,23,23	1.24	4 (20%)	
3	3PY	A	502	-	5,6,6	2.34	3 (60%)	6,7,7	1.65	2 (33%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	В	501	-	-	2/8/8/8	0/1/1/1
2	PLP	A	501	-	-	2/8/8/8	0/1/1/1
3	3PY	A	502	-	-	0/5/6/6	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(\text{\AA})$
3	A	502	3PY	O2-C1	-3.47	1.20	1.30
3	A	502	3PY	O3-C2	-2.77	1.17	1.23
2	A	501	PLP	C4-C4A	2.73	1.52	1.46
2	В	501	PLP	C4-C5	-2.66	1.38	1.42
2	В	501	PLP	C4-C4A	2.52	1.52	1.46
3	A	502	3PY	C2-C1	-2.46	1.50	1.53
2	В	501	PLP	C6-N1	2.27	1.39	1.34
2	A	501	PLP	P-O1P	2.20	1.57	1.50
2	A	501	PLP	C2-N1	2.02	1.37	1.33

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$Ideal(^{o})$
2	В	501	PLP	C5-C6-N1	-3.08	118.68	123.82
3	A	502	3PY	O2-C1-O1	-2.94	116.88	123.61
2	В	501	PLP	O4A-C4A-C4	-2.85	118.69	124.91
2	В	501	PLP	O4P-C5A-C5	-2.67	104.27	109.35
3	A	502	3PY	O2-C1-C2	2.48	120.75	113.97
2	A	501	PLP	O3-C3-C2	2.43	122.80	117.49
2	A	501	PLP	O4P-C5A-C5	-2.37	104.84	109.35

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	A	501	PLP	O4A-C4A-C4	-2.29	119.92	124.91
2	A	501	PLP	C2A-C2-C3	2.20	123.60	120.89

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	PLP	C3-C4-C4A-O4A
2	A	501	PLP	C5-C4-C4A-O4A
2	В	501	PLP	C3-C4-C4A-O4A
2	В	501	PLP	C5-C4-C4A-O4A

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	3PY	0	1

## 4.7 Other polymers (i)

There are no such residues in this entry.

## 4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 5 Fit of model and data (i)

#### 5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

#### 5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

#### 5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

#### 5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

### 5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

