

Full wwPDB X-ray Structure Validation Report (i)

Oct 3, 2023 – 01:57 AM EDT

PDB ID : 6PKD

Title: Myocilin OLF mutant N428D/D478H

Authors: Lieberman, R.L.; Hill, S.E.

Deposited on : 2019-06-29

Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : FAILED Xtriage (Phenix) : 1.13 EDS : FAILED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 8039 atoms, of which 3481 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Myocilin.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace			
1	A	268	Total 3861	C 1368	H 1727	N 344	O 415	S 7	0	1	0
1	В	258	Total 3838	C 1340	H 1754	N 336	O 402	S 6	0	2	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	428	ASP	ASN	engineered mutation	UNP Q99972
A	478	HIS	ASP	engineered mutation	UNP Q99972
В	428	ASP	ASN	engineered mutation	UNP Q99972
В	478	HIS	ASP	engineered mutation	UNP Q99972

• Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Na 1 1	0	0
2	В	1	Total Na 1 1	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	163	Total O 163 163	0	0
3	В	175	Total O 175 175	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source	
Space group	P 1 21 1	Depositor	
Cell constants	48.55Å 47.34Å 97.08Å	Depositor	
a, b, c, α , β , γ	90.00° 100.20° 90.00°	Depositor	
Resolution (Å)	33.63 - 1.90	Depositor	
% Data completeness	98.1 (33.63-1.90)	Depositor	
(in resolution range)	, , ,	_	
R_{merge}	0.14	Depositor	
R_{sym}	(Not available)	Depositor	
$< I/\sigma(I) > 1$	3.80 (at 1.91Å)	Xtriage	
Refinement program	PHENIX 1.15.2_3472	Depositor	
R, R_{free}	0.182 , 0.208	Depositor	
Wilson B-factor (\mathring{A}^2)	19.3	Xtriage	
Anisotropy	0.219	Xtriage	
L-test for twinning ²	$ < L > = 0.49, < L^2> = 0.32$	Xtriage	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	8039	wwPDB-VP	
Average B, all atoms $(Å^2)$	25.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 54.01 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.8840e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.



There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

