



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 30, 2022 – 04:48 pm BST

PDB ID : 7PKL  
Title : Mechanistic understanding of antibody masking with anti-idiotypic antibody fragments  
Authors : Hargreaves, D.  
Deposited on : 2021-08-25  
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.30  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.30


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	226	 78% 19% ..
2	L	368	 2% 72% 18% • 9%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4497 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called trastuzumab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	221	1656	1045	281	324	6	0	0	0

- Molecule 2 is a protein called trastuzumab Light Chain VHH fusion.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	335	2568	1603	437	518	10	0	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
3	L	1	5	4	1	0	0
3	L	1	5	4	1	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	L	1	Total O S 5 4 1	0	0
3	L	1	Total O S 5 4 1	0	0


- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	H	98	Total O 98 98	0	0
4	L	155	Total O 155 155	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: trastuzumab Heavy Chain

Chain H: 



- Molecule 2: trastuzumab Light Chain VHH fusion

Chain L: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	160.74Å 59.97Å 63.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	80.37 – 2.35 80.37 – 2.34	Depositor EDS
% Data completeness (in resolution range)	60.3 (80.37-2.35) 60.3 (80.37-2.34)	Depositor EDS
$R_{merge}$	0.30	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.33 (at 2.34Å)	Xtrriage
Refinement program	BUSTER 2.11.7	Depositor
R, $R_{free}$	0.180 , 0.265 0.183 , 0.268	Depositor DCC
$R_{free}$ test set	831 reflections (5.20%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.8	Xtrriage
Anisotropy	0.095	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4497	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.96% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	H	0.53	0/1698	0.78	0/2316
2	L	0.51	0/2624	0.77	0/3561
All	All	0.51	0/4322	0.78	0/5877

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1656	0	1610	18	0
2	L	2568	0	2461	28	0
3	L	20	0	0	0	0
4	H	98	0	0	0	0
4	L	155	0	0	2	0
All	All	4497	0	4071	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:5:VAL:HA	2:L:114:GLN:HE22	1.43	0.83
2:L:12:VAL:HG13	2:L:16:ASP:HB2	1.76	0.68
2:L:5:VAL:HA	2:L:114:GLN:NE2	2.10	0.65
2:L:208:LEU:HD11	2:L:216:PHE:O	1.98	0.64
2:L:29:PHE:HE1	2:L:74:ASN:HA	1.63	0.63
2:L:57:SER:HB3	2:L:185:THR:HG21	1.86	0.56
1:H:11:LEU:HB2	1:H:154:PRO:HG3	1.89	0.54
1:H:83:MET:HB3	1:H:86:LEU:HD21	1.89	0.54
2:L:6:GLU:OE1	2:L:113:GLY:HA3	2.08	0.52
2:L:220:ARG:HG3	2:L:225:PHE:CE2	2.45	0.52
2:L:36:TRP:HD1	2:L:70:ILE:HD12	1.74	0.52
1:H:36:TRP:HD1	1:H:70:ILE:HD12	1.76	0.51
2:L:29:PHE:CE1	2:L:74:ASN:HA	2.44	0.51
1:H:123:THR:HG22	1:H:210:SER:HB3	1.93	0.50
1:H:47:TRP:CD2	2:L:250:PRO:HD2	2.47	0.49
2:L:225:PHE:HB3	4:L:521:HOH:O	2.13	0.49
2:L:175:ILE:HD12	2:L:227:LEU:HD23	1.94	0.49
1:H:2:VAL:HG13	1:H:27:PHE:CD1	2.48	0.48
1:H:40:ALA:HB3	1:H:43:LYS:HB2	1.95	0.48
1:H:38:ARG:HG2	1:H:48:VAL:HG22	1.95	0.47
1:H:107:MET:HE3	1:H:110:TRP:CZ2	2.49	0.47
1:H:175:ALA:HA	1:H:185:LEU:HB3	1.97	0.47
1:H:30:LYS:HG2	1:H:54:THR:OG1	2.16	0.46
1:H:98:ARG:O	1:H:107:MET:HA	2.14	0.46
2:L:244:GLN:O	2:L:250:PRO:HA	2.15	0.46
1:H:12:VAL:HG11	1:H:86:LEU:HD13	1.98	0.45
1:H:172:THR:HA	1:H:187:SER:HA	1.98	0.45
2:L:189:TRP:HB2	2:L:202:ILE:HB	1.98	0.45
2:L:367:GLU:HB3	4:L:568:HOH:O	2.16	0.45
1:H:99:TRP:CZ2	2:L:245:HIS:HB2	2.53	0.44
2:L:39:GLN:HB3	2:L:95:TYR:HE2	1.81	0.44
2:L:309:GLN:HB3	2:L:312:ASN:HD21	1.82	0.43
2:L:91:THR:HG23	2:L:119:THR:HA	2.01	0.43
1:H:104:PHE:HZ	2:L:102:TRP:CG	2.36	0.43
2:L:301:GLN:HB3	2:L:308:LEU:HD11	1.99	0.43
2:L:313:SER:HA	2:L:332:THR:O	2.20	0.41
2:L:299:LYS:HG2	2:L:301:GLN:NE2	2.36	0.41
1:H:154:PRO:HD2	1:H:209:PRO:CB	2.51	0.41
1:H:159:VAL:HG22	1:H:205:VAL:HG22	2.03	0.41
2:L:248:THR:HA	2:L:249:PRO:HA	1.85	0.41
2:L:340:TYR:HA	2:L:346:TYR:OH	2.21	0.41
2:L:237:PHE:CZ	2:L:319:GLU:HG3	2.56	0.41

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:201:LEU:HA	2:L:212:VAL:HG21	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	219/226 (97%)	208 (95%)	11 (5%)	0	100	100
2	L	329/368 (89%)	311 (94%)	16 (5%)	2 (1%)	25	27
All	All	548/594 (92%)	519 (95%)	27 (5%)	2 (0%)	34	38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	L	254	GLN
2	L	292	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	181/187 (97%)	165 (91%)	16 (9%)	10	8
2	L	285/306 (93%)	257 (90%)	28 (10%)	8	7
All	All	466/493 (94%)	422 (91%)	44 (9%)	8	8

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	7	SER
1	H	13	GLN
1	H	17	SER
1	H	30	LYS
1	H	46	GLU
1	H	50	ARG
1	H	69	THR
1	H	123	THR
1	H	134	SER
1	H	138	THR
1	H	142	THR
1	H	180	SER
1	H	186	SER
1	H	196	LEU
1	H	211	ASN
1	H	213	LYS
2	L	7	SER
2	L	12	VAL
2	L	13	GLN
2	L	18	LEU
2	L	29	PHE
2	L	32	VAL
2	L	71	SER
2	L	78	THR
2	L	84	SER
2	L	108	ARG
2	L	157	GLN
2	L	164	SER
2	L	168	SER
2	L	181	GLN
2	L	191	GLN
2	L	206	SER
2	L	226	THR
2	L	243	GLN
2	L	257	LYS
2	L	259	GLU
2	L	301	GLN
2	L	306	ASN
2	L	315	GLU
2	L	321	ASP
2	L	335	LEU
2	L	341	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	L	345	VAL
2	L	360	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	28	ASN
1	H	199	GLN
2	L	114	GLN
2	L	181	GLN
2	L	301	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	L	403	-	4,4,4	0.21	0	6,6,6	0.20	0
3	SO4	L	404	-	4,4,4	0.20	0	6,6,6	0.19	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	L	401	-	4,4,4	0.21	0	6,6,6	0.25	0
3	SO4	L	402	-	4,4,4	0.12	0	6,6,6	0.14	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	H	221/226 (97%)	-0.18	1 (0%) 91 95	17, 33, 58, 99	0
2	L	335/368 (91%)	-0.09	7 (2%) 63 74	18, 32, 57, 95	0
All	All	556/594 (93%)	-0.13	8 (1%) 75 83	17, 33, 58, 99	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	136	LYS	5.9
2	L	366	GLY	3.7
2	L	2	VAL	3.1
2	L	308	LEU	2.8
2	L	26	GLY	2.5
2	L	123	GLY	2.3
2	L	344	LYS	2.2
2	L	367	GLU	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	L	404	5/5	0.92	0.14	83,85,87,88	0
3	SO4	L	402	5/5	0.93	0.13	81,83,84,86	0
3	SO4	L	403	5/5	0.98	0.21	66,66,70,71	0
3	SO4	L	401	5/5	0.99	0.11	44,45,46,49	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.