



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 29, 2023 – 11:50 PM EDT

PDB ID : 3PQR  
Title : Crystal structure of Metarhodopsin II in complex with a C-terminal peptide derived from the Galpha subunit of transducin  
Authors : Choe, H.-W.; Kim, Y.J.; Park, J.H.; Morizumi, T.; Pai, E.F.; Krauss, N.; Hofmann, K.P.; Scheerer, P.; Ernst, O.P.  
Deposited on : 2010-11-26  
Resolution : 2.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3168 (2.90-2.82)
Clashscore	141614	3438 (2.90-2.82)
Ramachandran outliers	138981	3348 (2.90-2.82)
Sidechain outliers	138945	3351 (2.90-2.82)
RSRZ outliers	127900	3103 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	348	 4% 70% 23% 6%
2	B	11	 82% 18%
3	C	4	 75% 25%
4	D	2	 50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
6	NAG	A	501	-	-	-	X
7	BOG	A	602	-	-	-	X

## 2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 2904 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rhodopsin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	326	2615	1741	404	444	26	0	2	0

- Molecule 2 is a protein called Guanine nucleotide-binding protein G(t) subunit alpha-1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	11	89	59	13	17	0	0	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	341	LEU	LYS	engineered mutation	UNP P04695
B	347	VAL	CYS	engineered mutation	UNP P04695

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	4	50	28	2	20	0	0	0

- Molecule 4 is an oligosaccharide called alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace	
4	D	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 5 is RETINAL (three-letter code: RET) (formula: C<sub>20</sub>H<sub>28</sub>O).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	C	0	0
			20	20		

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
6	A	1	14	8	1	5	0	0

- Molecule 7 is octyl beta-D-glucopyranoside (three-letter code: BOG) (formula:  $C_{14}H_{28}O_6$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
7	A	1	20	14	6	0	0
7	A	1	20	14	6	0	0

- Molecule 8 is PALMITIC ACID (three-letter code: PLM) (formula:  $C_{16}H_{32}O_2$ ).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	C O	0	0
			17	16 1		

- Molecule 9 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	1	Total	O S	0	0
			5	4 1		

- Molecule 10 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	1	Total C O 4 2 2	0	0
10	A	1	Total C O 4 2 2	0	0

- Molecule 11 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	A	19	Total O 19 19	0	0
11	B	4	Total O 4 4	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Rhodopsin



- Molecule 2: Guanine nucleotide-binding protein G(t) subunit alpha-1



- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	241.55Å 241.55Å 109.87Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	34.86 – 2.85 34.86 – 2.85	Depositor EDS
% Data completeness (in resolution range)	100.0 (34.86-2.85) 99.0 (34.86-2.85)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.87 (at 2.85Å)	Xtrriage
Refinement program	REFMAC 5.5.0072	Depositor
R, $R_{free}$	0.217 , 0.250 0.217 , 0.247	Depositor DCC
$R_{free}$ test set	1421 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	64.7	Xtrriage
Anisotropy	0.114	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 67.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	2904	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	62.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.75% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BOG, RET, BMA, GLC, MAN, SO4, PLM, ACT, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.46	0/2699	0.52	0/3678
2	B	0.42	0/89	0.48	0/117
All	All	0.46	0/2788	0.52	0/3795

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2615	0	2581	64	0
2	B	89	0	93	2	0
3	C	50	0	43	1	0
4	D	23	0	21	0	0
5	A	20	0	27	0	0
6	A	14	0	13	2	0
7	A	40	0	56	2	0
8	A	17	0	31	0	0
9	A	5	0	0	1	0
10	A	8	0	6	0	0
11	A	19	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
11	B	4	0	0	0	0
All	All	2904	0	2871	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:135:ARG:HH12	1:A:257:MET:HE1	1.32	0.93
1:A:135:ARG:HH12	1:A:257:MET:CE	1.83	0.92
1:A:183:MET:HE1	7:A:602:BOG:H7'2	1.56	0.86
1:A:62:THR:HG21	1:A:77:LEU:HD11	1.55	0.86
1:A:214:ILE:HB	1:A:215:PRO:HD3	1.70	0.74
1:A:319:THR:HG23	1:A:325:LYS:HA	1.72	0.72
1:A:135:ARG:NH1	1:A:257:MET:CE	2.53	0.72
1:A:183:MET:CE	7:A:602:BOG:H7'2	2.21	0.71
1:A:135:ARG:NH1	1:A:257:MET:HE1	2.07	0.69
1:A:183:MET:HE1	1:A:286:ILE:HD13	1.73	0.69
1:A:200:ASN:O	1:A:204:VAL:HG23	1.99	0.63
1:A:39:MET:HA	1:A:39:MET:HE2	1.80	0.62
3:C:3:BMA:O2	3:C:4:MAN:C1	2.48	0.60
1:A:39:MET:HA	1:A:39:MET:CE	2.31	0.60
1:A:6:GLY:HA3	1:A:9:PHE:CZ	2.37	0.60
1:A:227:VAL:O	1:A:231:LYS:HG2	2.02	0.58
1:A:44:MET:O	1:A:48:ILE:HG13	2.04	0.58
1:A:183:MET:CE	1:A:286:ILE:HD13	2.35	0.57
1:A:135:ARG:HH12	1:A:257:MET:HE3	1.68	0.55
1:A:302:ASN:ND2	11:A:350:HOH:O	2.37	0.55
1:A:75:ILE:CG2	1:A:131:LEU:HD13	2.37	0.54
1:A:135:ARG:HG2	2:B:347:VAL:HB	1.89	0.54
1:A:214:ILE:HB	1:A:215:PRO:CD	2.39	0.53
1:A:177:ARG:HH21	1:A:193:THR:HG21	1.74	0.53
1:A:50:LEU:O	1:A:54:ILE:HG12	2.09	0.52
1:A:28:GLN:NE2	1:A:30:TYR:OH	2.43	0.52
1:A:31:LEU:HD12	1:A:31:LEU:C	2.30	0.51
1:A:170:PRO:HB2	1:A:171:PRO:HD3	1.93	0.51
1:A:31:LEU:HB2	1:A:285:PRO:HG3	1.93	0.50
1:A:225:GLN:HA	1:A:225:GLN:OE1	2.13	0.49
1:A:233:ALA:HB3	2:B:340:ILE:HD13	1.95	0.49
1:A:282:ASP:OD2	6:A:501:NAG:H61	2.13	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:300:VAL:O	1:A:303:PRO:HD2	2.12	0.48
1:A:93:THR:HG21	1:A:113:GLU:HG3	1.95	0.48
1:A:135:ARG:NH1	1:A:257:MET:HE3	2.26	0.47
1:A:230:VAL:CG1	1:A:247:GLU:HA	2.44	0.47
1:A:40:LEU:HD23	1:A:98:SER:HB3	1.95	0.47
1:A:66:LYS:HA	1:A:69:ARG:NH2	2.30	0.47
1:A:64[B]:GLN:NE2	1:A:320:THR:OG1	2.47	0.47
1:A:180:PRO:HB2	1:A:184:GLN:NE2	2.30	0.47
1:A:75:ILE:HG21	1:A:131:LEU:HD13	1.98	0.46
1:A:125:LEU:HD13	1:A:261:PHE:CE1	2.51	0.45
1:A:64[B]:GLN:HG3	1:A:65:HIS:CD2	2.52	0.45
1:A:171:PRO:HA	1:A:175:TRP:O	2.17	0.45
1:A:9:PHE:C	1:A:9:PHE:CD1	2.90	0.45
1:A:193:THR:HG22	1:A:195:HIS:HB2	1.98	0.44
1:A:52:PHE:HB3	1:A:53:PRO:HD3	2.00	0.44
1:A:266:LEU:HB3	1:A:267:PRO:HD3	1.98	0.44
1:A:287:PHE:CD2	1:A:287:PHE:C	2.90	0.44
1:A:27:PRO:HB3	1:A:29:TYR:CE2	2.53	0.44
1:A:55:ASN:HD21	1:A:303:PRO:HG3	1.83	0.43
1:A:177:ARG:NH2	1:A:193:THR:HG21	2.33	0.43
1:A:238:GLN:HA	1:A:244:GLN:NE2	2.34	0.43
1:A:102:TYR:CZ	1:A:104:VAL:HG12	2.55	0.42
1:A:41:ALA:HB1	1:A:95:LEU:HD12	2.01	0.42
1:A:23:PRO:HB3	1:A:184:GLN:HB3	2.02	0.41
1:A:302:ASN:HB2	1:A:303:PRO:HD3	2.02	0.41
1:A:16:LYS:HB3	1:A:16:LYS:HE2	1.84	0.41
1:A:285:PRO:HD2	9:A:801:SO4:O3	2.20	0.41
1:A:31:LEU:C	1:A:31:LEU:CD1	2.89	0.41
1:A:257:MET:HG2	1:A:306:TYR:CZ	2.56	0.41
1:A:20:VAL:HA	1:A:30:TYR:CZ	2.56	0.41
1:A:282:ASP:HB2	6:A:501:NAG:H4	2.03	0.41
1:A:89:GLY:O	1:A:113:GLU:HG2	2.22	0.40
1:A:198:THR:O	1:A:199:ASN:HB2	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/348 (94%)	304 (93%)	20 (6%)	2 (1%)	25	53
2	B	9/11 (82%)	9 (100%)	0	0	100	100
All	All	335/359 (93%)	313 (93%)	20 (6%)	2 (1%)	25	53

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	176	SER
1	A	280	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	280/296 (95%)	273 (98%)	7 (2%)	47	76
2	B	10/10 (100%)	10 (100%)	0	100	100
All	All	290/306 (95%)	283 (98%)	7 (2%)	49	77

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	SER
1	A	31	LEU
1	A	61	VAL
1	A	193	THR

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type
1	A	283	PHE
1	A	293	PHE
1	A	326	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	GLN
1	A	55	ASN
1	A	184	GLN
1	A	244	GLN
1	A	312	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	C	1	1,3	14,14,15	0.61	0	17,19,21	1.15	1 (5%)
3	NAG	C	2	3	14,14,15	0.78	0	17,19,21	1.02	1 (5%)
3	BMA	C	3	3	11,11,12	1.00	0	15,15,17	1.82	4 (26%)
3	MAN	C	4	3	11,11,12	0.94	0	15,15,17	0.82	0
4	GLC	D	1	4	11,11,12	1.10	0	15,15,17	1.10	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GLC	D	2	4	12,12,12	1.01	0	17,17,17	0.88	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	BMA	C	3	3	-	2/2/19/22	0/1/1/1
3	MAN	C	4	3	-	2/2/19/22	0/1/1/1
4	GLC	D	1	4	-	2/2/19/22	0/1/1/1
4	GLC	D	2	4	-	1/2/22/22	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	3	BMA	O5-C1-C2	-4.07	104.50	110.77
4	D	1	GLC	O5-C5-C6	3.18	112.19	107.20
3	C	2	NAG	O5-C5-C6	2.93	111.80	107.20
3	C	3	BMA	C3-C4-C5	2.67	115.01	110.24
3	C	3	BMA	C1-O5-C5	-2.57	108.71	112.19
3	C	3	BMA	O5-C5-C4	2.56	117.07	110.83
3	C	1	NAG	C6-C5-C4	-2.17	107.93	113.00

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	1	GLC	O5-C5-C6-O6
3	C	3	BMA	O5-C5-C6-O6
3	C	4	MAN	O5-C5-C6-O6
4	D	1	GLC	C4-C5-C6-O6
3	C	4	MAN	C4-C5-C6-O6
3	C	3	BMA	C4-C5-C6-O6
4	D	2	GLC	O5-C5-C6-O6


There are no ring outliers.




2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	4	MAN	1	0
3	C	3	BMA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
8	PLM	A	701	1	16,16,17	0.49	0	15,15,17	0.70	0
7	BOG	A	602	-	20,20,20	1.06	2 (10%)	25,25,25	1.03	2 (8%)
6	NAG	A	501	1	14,14,15	1.17	1 (7%)	17,19,21	2.40	4 (23%)
5	RET	A	401	1	20,20,21	1.28	2 (10%)	27,27,28	0.95	1 (3%)
10	ACT	A	902	-	3,3,3	0.84	0	3,3,3	1.49	0
9	SO4	A	801	-	4,4,4	0.14	0	6,6,6	0.10	0
10	ACT	A	901	-	3,3,3	0.76	0	3,3,3	1.21	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
7	BOG	A	601	-	20,20,20	0.90	1 (5%)	25,25,25	1.35	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	PLM	A	701	1	-	8/13/14/15	-
7	BOG	A	602	-	-	8/11/31/31	0/1/1/1
6	NAG	A	501	1	-	2/6/23/26	0/1/1/1
5	RET	A	401	1	-	0/13/30/31	0/1/1/1
7	BOG	A	601	-	-	8/11/31/31	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	602	BOG	O1-C1	3.44	1.46	1.40
5	A	401	RET	C2-C3	-3.27	1.44	1.52
5	A	401	RET	C14-C13	3.12	1.36	1.33
6	A	501	NAG	O5-C1	2.77	1.48	1.43
7	A	601	BOG	O1-C1	2.71	1.44	1.40
7	A	602	BOG	O5-C1	2.23	1.47	1.41

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	501	NAG	O5-C1-C2	7.83	123.65	111.29
6	A	501	NAG	C1-O5-C5	4.07	117.71	112.19
7	A	601	BOG	C4-C3-C2	3.34	116.65	110.82
7	A	602	BOG	O5-C5-C4	3.09	115.30	109.69
6	A	501	NAG	O5-C5-C6	2.79	111.58	107.20
7	A	601	BOG	C1-C2-C3	2.77	115.77	110.00
5	A	401	RET	C17-C1-C2	-2.71	98.07	108.91
7	A	601	BOG	O5-C5-C4	-2.42	105.31	109.69
7	A	602	BOG	C1-O5-C5	2.39	118.38	113.69
6	A	501	NAG	C6-C5-C4	-2.25	107.73	113.00
7	A	601	BOG	C1-O5-C5	-2.22	109.32	113.69

There are no chirality outliers.

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	602	BOG	O5-C1-O1-C1'
7	A	602	BOG	C2'-C1'-O1-C1
7	A	602	BOG	O5-C5-C6-O6
7	A	601	BOG	C4-C5-C6-O6
7	A	601	BOG	O5-C5-C6-O6
8	A	701	PLM	C5-C6-C7-C8
7	A	601	BOG	C3'-C4'-C5'-C6'
8	A	701	PLM	CB-CC-CD-CE
7	A	602	BOG	C3'-C4'-C5'-C6'
7	A	602	BOG	C4-C5-C6-O6
7	A	602	BOG	C1'-C2'-C3'-C4'
6	A	501	NAG	C4-C5-C6-O6
7	A	602	BOG	C2'-C3'-C4'-C5'
8	A	701	PLM	C9-CA-CB-CC
8	A	701	PLM	C2-C3-C4-C5
7	A	602	BOG	C5'-C6'-C7'-C8'
8	A	701	PLM	CD-CE-CF-CG
8	A	701	PLM	C6-C7-C8-C9
7	A	601	BOG	C4'-C5'-C6'-C7'
7	A	601	BOG	C5'-C6'-C7'-C8'
8	A	701	PLM	C1-C2-C3-C4
6	A	501	NAG	O5-C5-C6-O6
7	A	601	BOG	C2-C1-O1-C1'
7	A	601	BOG	C2'-C3'-C4'-C5'
7	A	601	BOG	O5-C1-O1-C1'
8	A	701	PLM	CA-CB-CC-CD


There are no ring outliers.


3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	602	BOG	2	0
6	A	501	NAG	2	0
9	A	801	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	326/348 (93%)	-0.22	13 (3%) 38 32	25, 57, 89, 122	1 (0%)
2	B	11/11 (100%)	-0.12	0 100 100	54, 60, 75, 75	0
All	All	337/359 (93%)	-0.21	13 (3%) 39 34	25, 57, 89, 122	1 (0%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	279	GLN	5.5
1	A	145	ASN	4.2
1	A	147	ARG	4.0
1	A	277	THR	3.8
1	A	144	SER	3.2
1	A	143	MET	3.0
1	A	142	PRO	2.5
1	A	280	GLY	2.5
1	A	1	MET	2.3
1	A	324	GLY	2.3
1	A	282	ASP	2.2
1	A	199	ASN	2.2
1	A	283	PHE	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,


median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GLC	D	1	11/12	0.69	0.30	131,143,146,146	0
3	MAN	C	4	11/12	0.81	0.31	120,122,126,128	0
4	GLC	D	2	12/12	0.81	0.17	138,143,144,145	0
3	BMA	C	3	11/12	0.88	0.23	81,96,101,111	0
3	NAG	C	2	14/15	0.97	0.16	30,42,57,66	0
3	NAG	C	1	14/15	0.98	0.10	41,45,52,63	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	NAG	A	501	14/15	0.73	0.45	109,117,124,126	0
7	BOG	A	602	20/20	0.78	0.42	68,116,126,127	0
10	ACT	A	902	4/4	0.78	0.29	77,80,81,83	0
8	PLM	A	701	17/18	0.88	0.21	60,68,85,91	0
7	BOG	A	601	20/20	0.88	0.26	63,86,102,103	0
10	ACT	A	901	4/4	0.93	0.19	73,75,76,76	0
5	RET	A	401	20/21	0.93	0.43	61,76,83,84	0
9	SO4	A	801	5/5	0.97	0.21	99,101,103,103	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around BOG A 602:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around PLM A 701:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around BOG A 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around RET A 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.