



wwPDB X-ray Structure Validation Summary Report ⓘ

Jul 31, 2023 – 10:50 PM EDT

PDB ID : 3PRC
Title : PHOTOSYNTHETIC REACTION CENTER FROM RHODOPSEUDOMONAS VIRIDIS (QB-DEPLETED)
Authors : Lancaster, C.R.D.; Michel, H.
Deposited on : 1997-07-29
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.34
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.34


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	336	 82% 15% ..
2	L	273	 85% 13% .
3	M	323	 84% 16% .
4	H	258	 74% 24% .

2 Entry composition

There are 13 unique types of molecules in this entry. The entry contains 10606 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOTOSYNTHETIC REACTION CENTER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	C	332	2607	1642	467	480	18	38	1	0

- Molecule 2 is a protein called PHOTOSYNTHETIC REACTION CENTER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	273	2193	1471	358	357	7	14	2	0

- Molecule 3 is a protein called PHOTOSYNTHETIC REACTION CENTER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	M	323	2566	1711	420	424	11	14	1	0

- Molecule 4 is a protein called PHOTOSYNTHETIC REACTION CENTER.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	H	258	2028	1298	345	382	3	129	1	0

- Molecule 5 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
5	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
5	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
5	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
5	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 6 is BACTERIOCHLOROPHYLL B (three-letter code: BCB) (formula: $C_{55}H_{72}MgN_4O_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
6	L	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
6	L	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
6	M	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
6	M	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		

- Molecule 7 is BACTERIOPHEOPHYTIN B (three-letter code: BPB) (formula: $C_{55}H_{74}N_4O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	L	1	Total	C	N	O	0	0
			65	55	4	6		
7	M	1	Total	C	N	O	7	0
			65	55	4	6		

- Molecule 8 is LAURYL DIMETHYLAMINE-N-OXIDE (three-letter code: LDA) (formula: $C_{14}H_{31}NO$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
8	L	1	16	14	1	1	0	0
8	L	1	16	14	1	1	0	0
8	M	1	16	14	1	1	3	0
8	M	1	16	14	1	1	5	0
8	M	1	16	14	1	1	4	0
8	H	1	16	14	1	1	0	0
8	H	1	16	14	1	1	1	0

- Molecule 9 is FE (II) ION (three-letter code: FE2) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Fe		
9	M	1	1	1	0	0

- Molecule 10 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	M	1	Total	O	S	0	0
			5	4	1		
10	M	1	Total	O	S	0	0
			5	4	1		
10	M	1	Total	O	S	0	0
			5	4	1		
10	H	1	Total	O	S	0	0
			5	4	1		

- Molecule 11 is MENAQUINONE-7 (three-letter code: MQ7) (formula: $C_{46}H_{64}O_2$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	M	1	Total	C O	0	0
			48	46 2		

- Molecule 12 is 15-cis-1,2-dihydroneurosporene (three-letter code: NS5) (formula: C₄₀H₆₀).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	M	1	Total	C	9	0
			40	40		


- Molecule 13 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	C	163	Total	O	0	0
			163	163		
13	L	65	Total	O	0	0
			65	65		
13	M	97	Total	O	0	0
			97	97		
13	H	100	Total	O	0	0
			100	100		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PHOTOSYNTHETIC REACTION CENTER




- Molecule 2: PHOTOSYNTHETIC REACTION CENTER



- Molecule 3: PHOTOSYNTHETIC REACTION CENTER



- Molecule 4: PHOTOSYNTHETIC REACTION CENTER





4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	223.50Å 223.50Å 113.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.40 27.62 – 2.40	Depositor EDS
% Data completeness (in resolution range)	79.5 (10.00-2.40) 79.5 (27.62-2.40)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.69 (at 2.39Å)	Xtrriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.178 , 0.215 0.170 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	30.3	Xtrriage
Anisotropy	0.030	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 89.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10606	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MQ7, FME, BPB, HEM, FE2, SO4, BCB, NS5, LDA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.51	0/2674	0.60	0/3645
2	L	0.53	0/2281	0.57	0/3112
3	M	0.52	0/2671	0.58	0/3653
4	H	0.53	0/2055	0.71	3/2807 (0.1%)
All	All	0.52	0/9681	0.61	3/13217 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	5
2	L	0	1
3	M	0	1
All	All	0	7

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	53	ALA	CB-CA-C	-11.82	92.36	110.10
4	H	45	GLU	CB-CA-C	-7.85	94.70	110.40
4	H	53	ALA	N-CA-C	7.52	131.30	111.00

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	147	TYR	Sidechain
1	C	182	TYR	Sidechain
1	C	190	PHE	Sidechain
1	C	227	TYR	Sidechain
1	C	89	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2607	0	2575	33	0
2	L	2193	0	2122	31	0
3	M	2566	0	2460	37	0
4	H	2028	0	2029	38	0
5	C	172	0	120	1	0
6	L	132	0	144	9	0
6	M	132	0	144	13	0
7	L	65	0	74	5	0
7	M	65	0	74	12	0
8	H	32	0	62	0	0
8	L	32	0	62	10	0
8	M	48	0	93	2	0
9	M	1	0	0	0	0
10	H	5	0	0	0	0
10	M	15	0	0	0	0
11	M	48	0	64	0	0
12	M	40	0	60	0	0
13	C	163	0	0	5	0
13	H	100	0	0	2	0
13	L	65	0	0	2	0
13	M	97	0	0	0	0
All	All	10606	0	10083	157	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 157 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:L:402:BPB:HBBB	7:L:402:BPB:HHC	1.53	0.89
7:M:401:BPB:HBBB	7:M:401:BPB:HHC	1.50	0.89
6:M:805:BCB:HHC	6:M:805:BCB:HBB2	1.58	0.82
1:C:121:TRP:HA	1:C:123[A]:GLN:HE21	1.43	0.81
7:L:402:BPB:HHC	7:L:402:BPB:CBB	2.10	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	331/336 (98%)	320 (97%)	11 (3%)	0	100	100
2	L	273/273 (100%)	264 (97%)	9 (3%)	0	100	100
3	M	322/323 (100%)	308 (96%)	13 (4%)	1 (0%)	41	55
4	H	256/258 (99%)	247 (96%)	9 (4%)	0	100	100
All	All	1182/1190 (99%)	1139 (96%)	42 (4%)	1 (0%)	51	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	M	322	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	281/282 (100%)	264 (94%)	17 (6%)	19	31
2	L	220/218 (101%)	213 (97%)	7 (3%)	39	59
3	M	250/249 (100%)	245 (98%)	5 (2%)	55	74
4	H	212/212 (100%)	189 (89%)	23 (11%)	6	9
All	All	963/961 (100%)	911 (95%)	52 (5%)	22	36

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	M	290	ASP
4	H	89	LYS
4	H	236	ASP
4	H	9	HIS
4	H	42	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
4	H	102	GLN
3	M	147	ASN
2	L	239	ASN
2	L	214	GLN
3	M	16	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	FME	H	1[A]	4	8,9,10	0.68	0	7,9,11	2.23	2 (28%)
4	FME	H	1[B]	4	8,9,10	0.60	0	7,9,11	2.95	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FME	H	1[A]	4	-	3/7/9/11	-
4	FME	H	1[B]	4	-	4/7/9/11	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	1[B]	FME	CA-N-CN	-5.33	114.62	122.82
4	H	1[B]	FME	O1-CN-N	-5.01	112.09	125.27
4	H	1[A]	FME	CA-N-CN	-3.81	116.96	122.82
4	H	1[A]	FME	O1-CN-N	-3.81	115.25	125.27

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	H	1[A]	FME	O1-CN-N-CA
4	H	1[B]	FME	O1-CN-N-CA
4	H	1[B]	FME	CB-CG-SD-CE
4	H	1[A]	FME	N-CA-CB-CG
4	H	1[B]	FME	CA-CB-CG-SD

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 24 ligands modelled in this entry, 1 is monoatomic - leaving 23 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	LDA	L	702	-	12,15,15	2.14	1 (8%)	14,17,17	0.69	0
10	SO4	M	803	-	4,4,4	0.63	0	6,6,6	0.48	0
5	HEM	C	340	1	41,50,50	1.49	5 (12%)	45,82,82	1.16	5 (11%)
6	BCB	M	806	3	54,74,74	1.44	7 (12%)	52,115,115	2.48	13 (25%)
6	BCB	M	805	3	54,74,74	1.60	9 (16%)	52,115,115	2.31	13 (25%)
8	LDA	H	701	-	12,15,15	2.51	1 (8%)	14,17,17	0.75	0
10	SO4	H	801	-	4,4,4	0.64	0	6,6,6	0.40	0
7	BPB	M	401	-	49,70,70	1.37	9 (18%)	47,101,101	2.16	7 (14%)
8	LDA	M	706	-	12,15,15	2.23	1 (8%)	14,17,17	0.51	0
5	HEM	C	338	1	41,50,50	1.40	5 (12%)	45,82,82	1.25	5 (11%)
8	LDA	H	703	-	12,15,15	2.29	1 (8%)	14,17,17	0.59	0
5	HEM	C	339	1	41,50,50	1.59	5 (12%)	45,82,82	1.12	3 (6%)
12	NS5	M	600	-	39,39,39	0.73	0	44,46,46	1.15	5 (11%)
10	SO4	M	802	-	4,4,4	1.02	0	6,6,6	0.85	0
8	LDA	L	707	-	12,15,15	1.97	1 (8%)	14,17,17	0.60	0
10	SO4	M	804	-	4,4,4	1.03	0	6,6,6	0.68	0
6	BCB	L	304	2	54,74,74	1.55	8 (14%)	52,115,115	2.35	11 (21%)
6	BCB	L	302	2	54,74,74	1.48	6 (11%)	52,115,115	2.00	11 (21%)
7	BPB	L	402	-	49,70,70	1.32	6 (12%)	47,101,101	2.17	10 (21%)
11	MQ7	M	501	-	49,49,49	1.57	9 (18%)	60,63,63	1.37	9 (15%)
8	LDA	M	705	-	12,15,15	2.12	1 (8%)	14,17,17	0.53	0
5	HEM	C	337	1	41,50,50	1.47	5 (12%)	45,82,82	1.50	8 (17%)
8	LDA	M	704	-	12,15,15	2.24	1 (8%)	14,17,17	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	LDA	L	702	-	-	3/13/13/13	-
5	HEM	C	340	1	-	3/12/54/54	-
6	BCB	M	806	3	-	3/37/177/177	-
6	BCB	M	805	3	-	7/37/177/177	-
8	LDA	H	701	-	-	4/13/13/13	-
7	BPB	M	401	-	-	7/37/105/105	0/5/6/6
8	LDA	M	706	-	-	5/13/13/13	-
5	HEM	C	338	1	-	5/12/54/54	-
8	LDA	H	703	-	-	5/13/13/13	-
12	NS5	M	600	-	-	11/43/43/43	-
5	HEM	C	339	1	-	2/12/54/54	-
8	LDA	L	707	-	-	5/13/13/13	-
6	BCB	L	304	2	-	7/37/177/177	-
6	BCB	L	302	2	-	4/37/177/177	-
7	BPB	L	402	-	-	5/37/105/105	0/5/6/6
11	MQ7	M	501	-	-	1/41/61/61	0/2/2/2
8	LDA	M	705	-	-	2/13/13/13	-
5	HEM	C	337	1	-	5/12/54/54	-
8	LDA	M	704	-	-	5/13/13/13	-

The worst 5 of 81 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	H	701	LDA	O1-N1	-8.65	1.21	1.42
8	H	703	LDA	O1-N1	-7.89	1.23	1.42
8	M	704	LDA	O1-N1	-7.70	1.24	1.42
8	M	706	LDA	O1-N1	-7.68	1.24	1.42
8	L	702	LDA	O1-N1	-7.38	1.24	1.42

The worst 5 of 100 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	M	401	BPB	O2D-CGD-CBD	10.39	124.16	111.00
6	L	304	BCB	O2D-CGD-CBD	9.98	123.64	111.00
6	M	805	BCB	O2D-CGD-CBD	9.68	123.26	111.00
7	L	402	BPB	O2D-CGD-CBD	9.14	122.57	111.00
6	M	806	BCB	O2D-CGD-CBD	8.80	122.14	111.00

There are no chirality outliers.

5 of 89 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
6	L	302	BCB	C4-C3-C5-C6
6	M	806	BCB	CAD-CBD-CGD-O1D
6	M	806	BCB	CAD-CBD-CGD-O2D
7	L	402	BPB	O2A-C1-C2-C3
12	M	600	NS5	C3-C4-C5-C7



There are no ring outliers.


10 monomers are involved in 49 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	L	702	LDA	8	0
6	M	806	BCB	6	0
6	M	805	BCB	7	0
7	M	401	BPB	12	0
8	M	706	LDA	2	0
5	C	339	HEM	1	0
8	L	707	LDA	2	0
6	L	304	BCB	4	0
6	L	302	BCB	5	0
7	L	402	BPB	5	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.















5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	332/336 (98%)	-0.86	0 100 100	11, 25, 47, 67	15 (4%)
2	L	273/273 (100%)	-1.02	1 (0%) 92 91	9, 19, 41, 50	7 (2%)
3	M	323/323 (100%)	-0.97	1 (0%) 94 93	9, 22, 50, 61	8 (2%)
4	H	249/258 (96%)	-0.83	0 100 100	12, 30, 54, 64	22 (8%)
All	All	1177/1190 (98%)	-0.92	2 (0%) 95 94	9, 24, 49, 67	52 (4%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	59	TRP	2.1
3	M	319	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	FME	H	1[A]	10/11	0.97	0.09	22,25,27,27	10
4	FME	H	1[B]	10/11	0.97	0.09	27,30,30,30	10

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	LDA	L	707	16/16	0.64	0.27	54,64,65,66	0
8	LDA	M	706	16/16	0.74	0.28	69,70,72,72	4
8	LDA	L	702	16/16	0.81	0.25	19,44,66,66	0
8	LDA	M	704	16/16	0.83	0.22	53,59,61,62	3
12	NS5	M	600	40/40	0.88	0.19	31,49,68,68	9
8	LDA	M	705	16/16	0.90	0.15	51,55,59,59	5
8	LDA	H	703	16/16	0.94	0.16	30,41,52,52	1
11	MQ7	M	501	48/48	0.95	0.11	11,17,43,49	0
10	SO4	M	803	5/5	0.96	0.11	85,85,86,86	0
8	LDA	H	701	16/16	0.96	0.10	28,30,31,32	0
6	BCB	M	805	66/66	0.96	0.12	8,18,37,38	20
7	BPB	M	401	65/65	0.97	0.10	10,24,61,62	7
7	BPB	L	402	65/65	0.98	0.08	7,12,18,20	0
5	HEM	C	338	43/43	0.98	0.13	16,25,32,38	0
5	HEM	C	340	43/43	0.98	0.10	11,19,34,42	0
6	BCB	L	304	66/66	0.98	0.12	6,13,26,36	0
10	SO4	H	801	5/5	0.98	0.09	64,65,65,67	0
5	HEM	C	337	43/43	0.98	0.08	21,27,38,43	0
6	BCB	M	806	66/66	0.98	0.10	4,12,30,31	0
10	SO4	M	804	5/5	0.99	0.14	40,41,42,43	0
6	BCB	L	302	66/66	0.99	0.10	7,11,18,19	0
10	SO4	M	802	5/5	0.99	0.08	30,30,32,39	0
5	HEM	C	339	43/43	0.99	0.06	8,14,18,25	0
9	FE2	M	500	1/1	1.00	0.04	14,14,14,14	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NS5 M 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around MQ7 M 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BCB M 805:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around BPB M 401:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around BPB L 402:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)








Electron density around HEM C 340:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BCB L 304:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM C 337:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around BCB M 806:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around BCB L 302:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.