

# Full wwPDB X-ray Structure Validation Report (i)

#### Mar 10, 2024 – 01:59 PM EDT

PDB ID : 4PRK

Title: Crystal structure of D-lactate dehydrogenase (D-LDH) from Lactobacillus

jensenii

Authors: Kim, S.; Kim, K.J.

Deposited on : 2014-03-06

Resolution : 2.13 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.orgA user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467 Xtriage (Phenix) : 1.13

EDS : 2.36

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{\rm A})}) \end{array}$
$R_{free}$	130704	2523 (2.16-2.12)
Clashscore	141614	2653 (2.16-2.12)
Ramachandran outliers	138981	2618 (2.16-2.12)
Sidechain outliers	138945	2617 (2.16-2.12)
RSRZ outliers	127900	2485 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	336	79%	17%	
1	В	336	7%	19%	



# 2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 5168 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called 4-phosphoerythronate dehydrogenase.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	329	Total 2535	C 1619	N 415	O 490	S 11	0	0	0
1	В	329	Total 2535	C 1619	N 415	O 490	S 11	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	331	HIS	-	expression tag	UNP C5G4U0
A	332	HIS	-	expression tag	UNP C5G4U0
A	333	HIS	-	expression tag	UNP C5G4U0
A	334	HIS	-	expression tag	UNP C5G4U0
A	335	HIS	-	expression tag	UNP C5G4U0
A	336	HIS	-	expression tag	UNP C5G4U0
В	331	HIS	-	expression tag	UNP C5G4U0
В	332	HIS	-	expression tag	UNP C5G4U0
В	333	HIS	-	expression tag	UNP C5G4U0
В	334	HIS	-	expression tag	UNP C5G4U0
В	335	HIS	_	expression tag	UNP C5G4U0
В	336	HIS	_	expression tag	UNP C5G4U0

• Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	52	Total O 52 52	0	0
2	В	46	Total O 46 46	0	0




# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 4-phosphoerythronate dehydrogenase



• Molecule 1: 4-phosphoerythronate dehydrogenase





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants	90.53Å 90.53Å 157.78Å	Donogitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $120.00^{\circ}$	Depositor
Resolution (Å)	38.04 - 2.13	Depositor
Resolution (A)	38.04 - 2.13	EDS
% Data completeness	98.3 (38.04-2.13)	Depositor
(in resolution range)	98.4 (38.04-2.13)	EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.58  (at  2.14Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
D.D.	0.215 , $0.274$	Depositor
$R, R_{free}$	0.215 , $0.274$	DCC
$R_{free}$ test set	2110 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.1	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.38, 50.6	EDS
L-test for twinning <sup>2</sup>	$< L > = 0.47, < L^2> = 0.30$	Xtriage
Estimated twinning fraction	0.073 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5168	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.84% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain		nd lengths	Bond angles		
IVIOI		RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.75	$1/2577 \ (0.0\%)$	0.83	4/3489 (0.1%)	
1	В	0.71	0/2577	0.79	2/3489 (0.1%)	
All	All	0.73	1/5154 (0.0%)	0.81	6/6978 (0.1%)	

#### All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}( ext{\AA})$
1	A	8	PRO	N-CD	5.17	1.55	1.47

#### All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	115	MET	CG-SD-CE	-7.89	87.57	100.20
1	A	119	ARG	NE-CZ-NH1	6.13	123.36	120.30
1	В	51	LYS	C-N-CD	6.04	141.08	128.40
1	В	115	MET	CG-SD-CE	-5.83	90.88	100.20
1	A	51	LYS	C-N-CD	5.82	140.61	128.40
1	A	101	TYR	CB-CA-C	-5.38	99.64	110.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

$\mathbf{Mol}$	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2535	0	2575	37	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	В	2535	0	2575	46	0
2	A	52	0	0	0	0
2	В	46	0	0	1	0
All	All	5168	0	5150	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A., -1	A., 0	Interatomic	Clash
Atom-1	Atom-2	${\rm distance} \ (\rm \mathring{A})$	overlap (Å)
1:A:35:SER:O	1:A:39:VAL:HG23	1.75	0.85
1:B:52:PRO:HG2	1:B:82:THR:HG21	1.69	0.74
1:B:84:ASP:CG	1:B:87:LEU:HD13	2.09	0.73
1:B:99:PRO:HG3	1:B:326:PHE:CD1	2.32	0.63
1:A:80:THR:HG21	1:A:328:VAL:HG11	1.81	0.63
1:B:1:MET:HE2	1:B:316:ILE:HG23	1.82	0.60
1:B:84:ASP:OD2	1:B:87:LEU:HD13	2.03	0.59
1:B:264:GLU:OE2	1:B:295:HIS:HD2	1.86	0.58
1:A:111:VAL:O	1:A:115:MET:HG3	2.04	0.58
1:A:227:ASP:HA	1:A:254:ALA:HB2	1.86	0.58
1:A:55:ASN:HA	1:A:82:THR:HB	1.86	0.57
1:B:86:ASP:O	1:B:90:LYS:HG2	2.04	0.57
1:A:264:GLU:OE1	1:A:295:HIS:HD2	1.87	0.57
1:B:52:PRO:HG2	1:B:82:THR:CG2	2.35	0.56
1:B:191:ASP:HB2	2:B:427:HOH:O	2.06	0.56
1:A:75:LEU:CD2	1:A:80:THR:HG22	2.37	0.55
1:B:1:MET:CE	1:B:316:ILE:HG23	2.35	0.55
1:A:34:LEU:HD11	1:A:39:VAL:HG22	1.89	0.55
1:B:226:ASN:HA	1:B:252:GLU:O	2.07	0.54
1:A:36:ALA:HB2	1:A:60:ILE:HD11	1.89	0.54
1:A:100:VAL:HG13	1:A:103:PRO:HD3	1.91	0.52
1:B:226:ASN:OD1	1:B:252:GLU:OE2	2.29	0.51
1:B:1:MET:HA	1:B:46:ASP:OD2	2.10	0.51
1:B:7:SER:N	1:B:8:PRO:CD	2.74	0.50
1:A:1:MET:HE2	1:A:316:ILE:HG12	1.93	0.50
1:A:36:ALA:CB	1:A:60:ILE:HD11	2.42	0.50
1:B:281:TYR:CD1	1:B:281:TYR:C	2.86	0.49
1:B:176:ASP:OD1	1:B:177:VAL:N	2.45	0.49
1:B:177:VAL:HG23	1:B:178:ILE:HG12	1.95	0.49
1:A:39:VAL:O	1:A:42:VAL:HG23	2.13	0.48



 $Continued\ from\ previous\ page...$ 

Continued from pret		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:B:96:THR:HG21	1:B:314:ASP:HB3	1.95	0.48
1:A:39:VAL:HG21	1:A:61:VAL:HG22	1.95	0.48
1:B:106:ILE:HA	1:B:297:ALA:HB1	1.96	0.47
1:B:85:PHE:HE2	1:B:328:VAL:O	1.97	0.47
1:A:12:GLU:HG2	1:A:308:ILE:HD11	1.97	0.47
1:A:42:VAL:O	1:A:45:CYS:HB2	2.14	0.47
1:A:34:LEU:CD1	1:A:39:VAL:HG22	2.45	0.47
1:B:187:LEU:HD12	1:B:187:LEU:C	2.35	0.47
1:A:75:LEU:HD23	1:A:80:THR:HG22	1.96	0.46
1:B:227:ASP:HA	1:B:254:ALA:HB2	1.98	0.46
1:A:180:ASN:O	1:A:183:VAL:HG22	2.15	0.46
1:A:258:LEU:HD13	1:A:261:LEU:HD21	1.96	0.46
1:B:272:VAL:HB	1:B:276:GLN:HB2	1.97	0.45
1:B:184:GLU:N	1:B:185:PRO:CD	2.79	0.45
1:B:192:PHE:CE1	1:B:196:LEU:HD22	2.52	0.45
1:A:118:ASN:OD1	1:A:147:LYS:HE3	2.18	0.44
1:B:318:LEU:HD13	1:B:324:ALA:HB2	1.99	0.44
1:A:16:VAL:HG21	1:A:29:LEU:HD21	1.98	0.44
1:A:97:ASN:OD1	1:A:99:PRO:HD3	2.16	0.44
1:B:180:ASN:OD1	1:B:182:GLU:HG2	2.17	0.44
1:B:267:TYR:CD1	1:B:267:TYR:N	2.84	0.44
1:A:159:SER:O	1:A:163:GLN:HG3	2.17	0.44
1:B:246:GLU:HG2	1:B:250:LYS:HE3	2.00	0.44
1:B:154:VAL:HG12	1:B:155:GLY:N	2.32	0.44
1:B:87:LEU:HD12	1:B:87:LEU:N	2.32	0.43
1:A:42:VAL:O	1:A:42:VAL:HG12	2.18	0.43
1:A:75:LEU:HD21	1:A:80:THR:HG22	2.01	0.43
1:B:226:ASN:OD1	1:B:252:GLU:CD	2.57	0.43
1:B:111:VAL:O	1:B:115:MET:CG	2.67	0.43
1:A:300:THR:HB	1:B:142:ASN:HA	2.00	0.43
1:A:81:ASN:OD1	1:A:82:THR:HG23	2.19	0.42
1:B:78:VAL:HG13	1:B:79:GLY:N	2.35	0.42
1:B:280:ASP:OD1	1:B:281:TYR:N	2.52	0.42
1:A:184:GLU:N	1:A:185:PRO:CD	2.82	0.42
1:B:52:PRO:CG	1:B:82:THR:HG21	2.42	0.42
1:B:89:LYS:HD3	1:B:89:LYS:HA	1.71	0.42
1:B:6:PHE:HB3	1:B:33:SER:HA	2.02	0.42
1:A:6:PHE:CE1	1:A:48:VAL:HG13	2.55	0.42
1:A:57:ASP:OD1	1:A:57:ASP:N	2.45	0.41
1:A:101:TYR:HB2	1:A:102:SER:H	1.66	0.41
1:A:7:SER:HA	1:A:31:ASP:O	2.21	0.41



Continued from previous page...

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:B:56:LEU:HD12	1:B:56:LEU:HA	1.95	0.41
1:A:38:ASN:O	1:A:41:GLN:HG3	2.20	0.41
1:B:258:LEU:O	1:B:292:VAL:HA	2.21	0.41
1:B:99:PRO:CG	1:B:326:PHE:CD1	3.02	0.41
1:B:64:LYS:HG2	1:B:68:TYR:CE2	2.56	0.41
1:B:208:LEU:HD22	1:B:237:GLU:HB2	2.02	0.41
1:B:256:ALA:O	1:B:290:VAL:HA	2.20	0.41
1:A:64:LYS:HA	1:A:67:ASP:HB2	2.02	0.41
1:B:111:VAL:O	1:B:115:MET:HG2	2.21	0.40
1:A:1:MET:HE1	1:A:46:ASP:HB3	2.03	0.40
1:A:267:TYR:CD1	1:A:267:TYR:N	2.89	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	327/336 (97%)	315 (96%)	12 (4%)	0	100	100
1	В	327/336 (97%)	314 (96%)	11 (3%)	2 (1%)	25	17
All	All	$654/672 \ (97\%)$	629 (96%)	23 (4%)	2 (0%)	41	36

#### All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	92	HIS
1	В	321	GLY



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	274/281 (98%)	266 (97%)	8 (3%)	42 40
1	В	274/281 (98%)	261 (95%)	13 (5%)	26 21
All	All	548/562 (98%)	527 (96%)	21 (4%)	33 30

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	53	LEU
1	A	57	ASP
1	A	60	ILE
1	A	75	LEU
1	A	87	LEU
1	A	197	LYS
1	A	313	ARG
1	A	322	GLU
1	В	1	MET
1	В	63	LYS
1	В	80	THR
1	В	100	VAL
1	В	115	MET
1	В	191	ASP
1	В	227	ASP
1	В	252	GLU
1	В	267	TYR
1	В	274	ASP
1	В	295	HIS
1	В	318	LEU
1	В	322	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	135	ASN



Continued from previous page...

Mol	Chain	Res	Type
1	A	163	GLN
1	A	295	HIS
1	В	251	HIS
1	В	295	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

There are no ligands in this entry.

## 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2	$OWAB(A^2)$	Q < 0.9
1	A	329/336~(97%)	0.40	38 (11%) 4 6	22, 38, 78, 88	0
1	В	329/336~(97%)	0.27	22 (6%) 17 22	23, 43, 77, 90	0
All	All	658/672 (97%)	0.34	60 (9%) 9 12	22, 41, 78, 90	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	34	LEU	6.0
1	A	53	LEU	5.6
1	A	54	GLY	5.5
1	A	56	LEU	5.1
1	A	329	SER	4.7
1	A	62	TYR	4.4
1	A	86	ASP	4.2
1	A	58	ASP	4.2
1	A	60	ILE	4.2
1	В	321	GLY	4.1
1	В	57	ASP	4.1
1	В	323	ARG	4.0
1	A	36	ALA	3.9
1	A	35	SER	3.9
1	A	57	ASP	3.8
1	A	91	TYR	3.7
1	A	69	GLY	3.7
1	В	329	SER	3.6
1	A	61	VAL	3.6
1	В	60	ILE	3.6
1	В	92	HIS	3.5
1	A	85	PHE	3.5
1	A	39	VAL	3.4
1	A	92	HIS	3.4



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	В	328	VAL	3.3
1	A	52	PRO	3.0
1	A	83	ILE	3.0
1	В	65	LEU	3.0
1	A	66	ALA	2.9
1	A	323	ARG	2.9
1	A	38	ASN	2.9
1	В	318	LEU	2.8
1	A	328	VAL	2.8
1	В	90	LYS	2.8
1	A	87	LEU	2.8
1	A	89	LYS	2.7
1	В	56	LEU	2.7
1	В	211	SER	2.6
1	A	67	ASP	2.6
1	В	66	ALA	2.5
1	В	322	GLU	2.4
1	A	88	ALA	2.4
1	В	88	ALA	2.4
1	A	47	GLY	2.3
1	В	85	PHE	2.3
1	A	94	THR	2.2
1	A	55	ASN	2.2
1	A	40	ASP	2.2
1	В	324	ALA	2.2
1	В	35	SER	2.2
1	В	63	LYS	2.2
1	A	82	THR	2.2
1	A	93	LEU	2.1
1	A	37	GLU	2.1
1	A	110	ALA	2.1
1	A	59	GLU	2.0
1	В	87	LEU	2.0
1	A	24	GLY	2.0
1	В	86	ASP	2.0
1	В	53	LEU	2.0

# 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.4 Ligands (i)

There are no ligands in this entry.

## 6.5 Other polymers (i)

There are no such residues in this entry.

