



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 26, 2024 – 01:32 PM EST

PDB ID : 5PRL
Title : PanDDA analysis group deposition – Crystal Structure of BRD1 after initial refinement with no ligand modelled (structure 93)
Authors : Pearce, N.M.; Krojer, T.; Talon, R.; Bradley, A.R.; Fairhead, M.; Sethi, R.; Wright, N.; MacLean, E.; Collins, P.; Brandao-Neto, J.; Douangamath, A.; Renjie, Z.; Dias, A.; Ng, J.; Brennan, P.E.; Cox, O.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.; von Delft, F.
Deposited on : 2017-02-07
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	156	
1	B	156	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2048 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bromodomain-containing protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	121	989	621	175	186	7	0	5	0
1	B	123	1038	651	187	194	6	0	6	0

There are 50 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP O95696
A	2	HIS	-	expression tag	UNP O95696
A	3	HIS	-	expression tag	UNP O95696
A	4	HIS	-	expression tag	UNP O95696
A	5	HIS	-	expression tag	UNP O95696
A	6	HIS	-	expression tag	UNP O95696
A	7	HIS	-	expression tag	UNP O95696
A	8	SER	-	expression tag	UNP O95696
A	9	SER	-	expression tag	UNP O95696
A	10	GLY	-	expression tag	UNP O95696
A	11	VAL	-	expression tag	UNP O95696
A	12	ASP	-	expression tag	UNP O95696
A	13	LEU	-	expression tag	UNP O95696
A	14	GLY	-	expression tag	UNP O95696
A	15	THR	-	expression tag	UNP O95696
A	16	GLU	-	expression tag	UNP O95696
A	17	ASN	-	expression tag	UNP O95696
A	18	LEU	-	expression tag	UNP O95696
A	19	TYR	-	expression tag	UNP O95696
A	20	PHE	-	expression tag	UNP O95696
A	21	GLN	-	expression tag	UNP O95696
A	22	SER	-	expression tag	UNP O95696
A	23	MET	VAL	engineered mutation	UNP O95696
A	34	GLU	PRO	engineered mutation	UNP O95696
A	37	ARG	VAL	engineered mutation	UNP O95696

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	MET	-	expression tag	UNP O95696
B	2	HIS	-	expression tag	UNP O95696
B	3	HIS	-	expression tag	UNP O95696
B	4	HIS	-	expression tag	UNP O95696
B	5	HIS	-	expression tag	UNP O95696
B	6	HIS	-	expression tag	UNP O95696
B	7	HIS	-	expression tag	UNP O95696
B	8	SER	-	expression tag	UNP O95696
B	9	SER	-	expression tag	UNP O95696
B	10	GLY	-	expression tag	UNP O95696
B	11	VAL	-	expression tag	UNP O95696
B	12	ASP	-	expression tag	UNP O95696
B	13	LEU	-	expression tag	UNP O95696
B	14	GLY	-	expression tag	UNP O95696
B	15	THR	-	expression tag	UNP O95696
B	16	GLU	-	expression tag	UNP O95696
B	17	ASN	-	expression tag	UNP O95696
B	18	LEU	-	expression tag	UNP O95696
B	19	TYR	-	expression tag	UNP O95696
B	20	PHE	-	expression tag	UNP O95696
B	21	GLN	-	expression tag	UNP O95696
B	22	SER	-	expression tag	UNP O95696
B	23	MET	VAL	engineered mutation	UNP O95696
B	34	GLU	PRO	engineered mutation	UNP O95696
B	37	ARG	VAL	engineered mutation	UNP O95696

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	4	0
2	A	1	Total C O 8 4 4	8	1
2	B	1	Total C O 4 2 2	4	0
2	B	1	Total C O 4 2 2	4	1


- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Na 1 1	1	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bromodomain-containing protein 1



- Molecule 1: Bromodomain-containing protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	55.52Å 56.70Å 101.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.09 – 1.75 29.09 – 1.75	Depositor EDS
% Data completeness (in resolution range)	96.8 (29.09-1.75) 96.9 (29.09-1.75)	Depositor EDS
R_{merge}	0.68	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.99 (at 1.75Å)	Xtrriage
Refinement program	REFMAC 5.8.0131	Depositor
R, R_{free}	0.479 , 0.511 0.478 , 0.511	Depositor DCC
R_{free} test set	1511 reflections (4.74%)	wwPDB-VP
Wilson B-factor (Å ²)	19.7	Xtrriage
Anisotropy	0.006	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 57.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	0.042 for k,h,-l	Xtrriage
F_o, F_c correlation	0.58	EDS
Total number of atoms	2048	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/1016	0.69	0/1371
1	B	0.46	0/1068	0.74	0/1437
All	All	0.46	0/2084	0.72	0/2808

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	989	0	959	8	0
1	B	1038	0	1033	6	2
2	A	12	0	18	0	2
2	B	8	0	12	0	0
3	A	1	0	0	0	0
All	All	2048	0	2022	14	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:103[B]:ILE:HD13	1:A:124[B]:ARG:HB2	1.57	0.85
1:A:38:LEU:HD11	1:A:137:GLU:HB3	1.75	0.67
1:B:23:MET:O	1:B:25:GLN:N	2.39	0.56
1:A:54:ILE:HG22	1:A:55:PHE:CD1	2.46	0.51
1:A:103[B]:ILE:HD13	1:A:124[B]:ARG:CB	2.38	0.50
1:B:23:MET:O	1:B:26:VAL:N	2.46	0.48
1:A:90:LYS:N	1:A:94:GLU:OE1	2.47	0.48
1:B:76:ASP:OD2	1:B:78:ALA:N	2.50	0.44
1:B:32:LEU:HD23	1:B:32:LEU:C	2.37	0.44
1:A:92:LEU:CD2	1:A:135:ARG:HA	2.48	0.44
1:A:40:ARG:HA	1:A:84:LEU:HD21	1.99	0.43
1:B:27:ALA:HA	1:B:30:LEU:HG	2.02	0.41
1:B:137:GLU:O	1:B:141:ILE:HG13	2.21	0.40
1:A:73:HIS:O	1:A:105:ASN:ND2	2.50	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:23:MET:CG	2:A:201:EDO:C2[2_555]	1.82	0.38
1:B:23:MET:SD	2:A:202[A]:EDO:O1[2_555]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	124/156 (80%)	119 (96%)	4 (3%)	1 (1%)	19 6
1	B	127/156 (81%)	122 (96%)	3 (2%)	2 (2%)	9 1
All	All	251/312 (80%)	241 (96%)	7 (3%)	3 (1%)	12 3

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	139	ASP
1	B	24	GLU
1	B	87	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	103/138 (75%)	103 (100%)	0	100	100
1	B	112/138 (81%)	111 (99%)	1 (1%)	78	67
All	All	215/276 (78%)	214 (100%)	1 (0%)	88	83

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	23	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	A	201	-	3,3,3	0.48	0	2,2,2	0.43	0
2	EDO	B	201	-	3,3,3	0.46	0	2,2,2	0.31	0
2	EDO	B	202[B]	-	3,3,3	0.47	0	2,2,2	0.31	0
2	EDO	A	202[A]	-	3,3,3	0.45	0	2,2,2	0.40	0
2	EDO	A	202[B]	-	3,3,3	0.47	0	2,2,2	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	201	-	-	0/1/1/1	-
2	EDO	B	201	-	-	1/1/1/1	-
2	EDO	B	202[B]	-	-	0/1/1/1	-
2	EDO	A	202[A]	-	-	1/1/1/1	-
2	EDO	A	202[B]	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	201	EDO	O1-C1-C2-O2
2	A	202[B]	EDO	O1-C1-C2-O2
2	A	202[A]	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	201	EDO	0	1
2	A	202[A]	EDO	0	1

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	121/156 (77%)	4.28	101 (83%) 0 0	12, 18, 31, 49	0
1	B	123/156 (78%)	3.86	107 (86%) 0 0	9, 15, 21, 24	0
All	All	244/312 (78%)	4.07	208 (85%) 0 0	9, 17, 27, 49	0

All (208) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	141	ILE	18.6
1	A	142	GLY	12.6
1	A	140	SER	11.2
1	A	130	VAL	9.7
1	B	61	LEU	9.6
1	A	121	VAL	9.5
1	A	35	LEU	9.1
1	B	101	LEU	8.8
1	B	143	LEU	8.6
1	B	138	VAL	8.5
1	A	138	VAL	8.4
1	A	84	LEU	8.3
1	B	55	PHE	7.9
1	B	71	ILE	7.9
1	B	64	VAL	7.8
1	B	130	VAL	7.8
1	A	60	SER	7.3
1	A	54	ILE	7.0
1	B	139	ASP	7.0
1	A	43	LEU	7.0
1	A	134	ALA	7.0
1	A	95	PHE	7.0
1	B	68	LEU	6.9
1	A	129	VAL	6.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	64	VAL	6.7
1	B	88	GLY	6.7
1	A	128	GLY	6.7
1	A	116	PHE	6.7
1	A	77	PHE	6.7
1	B	121	VAL	6.7
1	A	71	ILE	6.6
1	B	84	LEU	6.6
1	B	86	ALA	6.5
1	A	39	LEU	6.5
1	A	51	PRO	6.3
1	B	52	ALA	6.1
1	A	32	LEU	6.0
1	A	68	LEU	6.0
1	B	25	GLN	6.0
1	A	131	LEU	6.0
1	B	32	LEU	5.9
1	A	26	VAL	5.9
1	A	102	ILE	5.9
1	A	117	TYR	5.8
1	A	42	VAL	5.8
1	A	103[A]	ILE	5.7
1	A	82	LYS	5.7
1	B	129	VAL	5.7
1	B	35	LEU	5.6
1	B	109	TYR	5.6
1	A	61	LEU	5.6
1	B	99	PHE	5.6
1	B	92	LEU	5.4
1	B	141	ILE	5.3
1	B	87	GLN	5.2
1	A	92	LEU	5.2
1	B	43	LEU	5.2
1	A	99	PHE	5.1
1	A	136	ARG	5.1
1	B	53[A]	ARG	5.1
1	B	46	LEU	5.1
1	B	48[A]	ASP	5.0
1	B	103	ILE	4.9
1	A	55	PHE	4.9
1	A	45	GLN	4.9
1	A	22	SER	4.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	26	VAL	4.9
1	A	52	ALA	4.9
1	A	93	HIS	4.8
1	A	78	ALA	4.8
1	B	27	ALA	4.8
1	A	62	LYS	4.8
1	A	23[A]	MET	4.8
1	A	123	LEU	4.8
1	A	36	THR	4.7
1	A	115	VAL	4.7
1	B	54	ILE	4.7
1	B	59	VAL	4.7
1	A	133	GLN	4.7
1	A	50	ASP	4.6
1	A	28	MET	4.6
1	B	117	TYR	4.6
1	A	81	ARG	4.6
1	B	75	MET	4.6
1	B	28	MET	4.6
1	B	91	ASN	4.5
1	A	139	ASP	4.5
1	A	75	MET	4.4
1	B	30	LEU	4.4
1	A	58	PRO	4.4
1	B	100	ASP	4.4
1	A	101	LEU	4.4
1	A	97[A]	GLU	4.4
1	B	24	GLU	4.3
1	A	33	THR	4.3
1	B	67	TYR	4.3
1	B	111	ALA	4.3
1	A	108[A]	LYS	4.2
1	B	77	PHE	4.2
1	B	72	LYS	4.2
1	A	83	ARG	4.1
1	B	123	LEU	4.1
1	B	38	LEU	4.1
1	A	122	ARG	4.1
1	B	128	GLY	4.1
1	A	137	GLU	4.1
1	A	135	ARG	4.0
1	A	65	PRO	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	87	GLN	3.9
1	A	89	TYR	3.9
1	A	127	GLY	3.9
1	B	31	ARG	3.9
1	B	62	LYS	3.9
1	A	114	THR	3.8
1	B	113	ASP	3.8
1	B	63	GLU	3.8
1	B	140	SER	3.8
1	B	114	THR	3.8
1	A	109	TYR	3.7
1	A	74	PRO	3.7
1	B	90	LYS	3.7
1	A	79	THR	3.7
1	B	132	ARG	3.6
1	A	120	ALA	3.6
1	A	46	LEU	3.6
1	B	95	PHE	3.6
1	B	134	ALA	3.6
1	B	39	LEU	3.5
1	A	132	ARG	3.5
1	A	25	GLN	3.5
1	B	79	THR	3.5
1	B	94	GLU	3.5
1	A	38	LEU	3.4
1	B	37[A]	ARG	3.4
1	B	144	GLU	3.4
1	B	102	ILE	3.4
1	B	108[A]	LYS	3.4
1	A	107	MET	3.4
1	A	30	LEU	3.4
1	B	42	VAL	3.4
1	B	22	SER	3.4
1	A	124[A]	ARG	3.3
1	B	127	GLY	3.3
1	A	73	HIS	3.3
1	B	50	ASP	3.3
1	B	89	TYR	3.3
1	B	44[A]	ASP	3.2
1	B	36	THR	3.2
1	B	60	SER	3.2
1	B	34	GLU	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	40	ARG	3.1
1	B	83	ARG	3.1
1	B	74	PRO	3.1
1	B	125	ASP	3.1
1	A	126	GLN	3.1
1	B	33	THR	3.1
1	A	59	VAL	3.1
1	B	93	HIS	3.1
1	B	56	ALA	3.0
1	B	82	LYS	3.0
1	B	96[A]	GLU	3.0
1	B	45	GLN	3.0
1	B	57	GLN	2.9
1	A	27	ALA	2.9
1	B	58	PRO	2.9
1	B	81	ARG	2.9
1	A	110	ASN	2.9
1	B	112	ARG	2.9
1	B	116	PHE	2.9
1	B	70	HIS	2.9
1	A	119	ALA	2.8
1	B	65	PRO	2.8
1	A	86	ALA	2.8
1	A	112	ARG	2.7
1	B	131	LEU	2.7
1	B	51	PRO	2.7
1	A	67	TYR	2.7
1	B	78	ALA	2.7
1	A	104	ASP	2.7
1	A	47	GLN	2.6
1	A	70	HIS	2.6
1	B	23	MET	2.6
1	A	41	SER	2.6
1	A	63	GLU	2.5
1	B	136	ARG	2.5
1	B	126	GLN	2.5
1	B	29	GLU	2.5
1	A	37	ARG	2.4
1	A	40	ARG	2.4
1	B	142	GLY	2.3
1	A	49	LYS	2.3
1	A	56	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	98	ASP	2.3
1	A	118	ARG	2.3
1	B	115	VAL	2.2
1	A	85	GLU	2.2
1	B	66	ASP	2.2
1	A	48	ASP	2.2
1	B	135	ARG	2.2
1	A	106	CYS	2.1
1	B	119	ALA	2.1
1	A	94	GLU	2.1
1	B	124	ARG	2.0
1	A	111	ALA	2.0
1	B	69	ASP	2.0
1	B	104	ASP	2.0
1	A	105	ASN	2.0
1	B	120	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

LIGAND-RSR INFOmissingINFO

6.5 Other polymers [i](#)

There are no such residues in this entry.